

COVID-19 in Cuba: Assessing the National Response.

Source: MEDICC Review . Oct2020, Vol. 22 Issue 4, p29-34. 6p.

Author(s): Galbán-García, Enrique; Más-Bermejo, Pedro

Abstract:

The COVID-19 pandemic exhibits different characteristics in each country, related to the extent of SARS-CoV-2 local transmission, as well as the speed and effectiveness of epidemic response implemented by authorities. This study presents a descriptive epidemiological analysis of the daily and cumulative incidence of confirmed cases and deaths in Cuba from COVID-19 in the first 110 days after first-case confirmation on March 11, 2020. During this period, 2340 cases (20.7 x 100,000 population) were confirmed, of which 86 patients died (case fatality 3.67%; 52 men and 34 women). Mean age of the deceased was 73.6 years (with a minimum of 35 years and a maximum of 101), with the average age of men lower than that of women. More than 70% of all deceased had associated noncommunicable diseases. The incidence curve ascended for five weeks and then descended steadily. The average number of confirmed cases and deaths for the last week included (June 23--28, 2020) were 25 and 1 respectively; the curve always moved within the most favorable forecast zone of available mathematical models and the effective reproductive number fell below 1 after the fifth week following the onset of the epidemic. We present the prevention and control measures implemented during this period--some unique to Cuba--and assess their effectiveness using two analytical tools: comparison of observed deaths and confirmed cases with those predicted by mathematical models; and estimation of the effective reproductive rate of SARS-CoV-2. Some distinctive features of this strategy include nationwide door-to-door active screening for individuals with fever and/or symptoms of respiratory distress, isolation of cases and quarantine of contacts of confirmed cases and of persons suspected of having the virus. During this period, Cuba's response to the epidemic was successful in flattening the curve and limiting transmission, resulting in fewer cases and a lower number of subsequent deaths.

Copyright of MEDICC Review is the property of Medical Education Cooperation with Cuba and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use. This abstract may be abridged. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material for the full abstract.

For access to this entire article and additional high quality information, please check with your college/university library, local public library, or affiliated institution.



Important User Information: Remote access to EBSCO's databases is permitted to patrons of subscribing institutions accessing from remote locations for personal, non-commercial use. However, remote access to EBSCO's databases from non-subscribing institutions is not allowed if the purpose of the use is for commercial gain through cost reduction or avoidance for a non-subscribing institution.

[Privacy Policy](#) | [A/B Testing](#) | [Terms of Use](#) | [Copyright](#) | [Cookie Policy](#)

© 2021 EBSCO Industries, Inc. All rights reserved.

