

Journal of the
MARINE BIOLOGICAL ASSOCIATION
of the United Kingdom



CAMBRIDGE
UNIVERSITY PRESS

**Abundance distribution patterns of intertidal bivalves
Macoma balthica and *Cerastoderma edule* at the Murman
coast tidal flats (the Barents Sea)**

Journal:	<i>Journal of the Marine Biological Association of the United Kingdom</i>
Manuscript ID:	JMBA-11-14-SIEMBS-0452.R3
Manuscript Type:	Special Issue - EMBS European Marine Biology Symposium
Date Submitted by the Author:	12-Apr-2015
Complete List of Authors:	Nazarova, Sophia; St-Petersburg State University, Ichthyology and Hydrobiology ; Russian State Pedagogical University, Zoology Shunkina, Ksenia; Zoological Institute RAS, Laboratory of Evolutional Morphology Genelt-Yanovskiy, Evgeny; Saint-Petersburg State University, Ichthyology and hydrobiology; Baltic Fund for Nature,
Keywords:	<i>Cerastoderma edule</i>, cockle, <i>Macoma balthica</i>, Barents sea, Geographic abundance distribution, latitudinal variation, species ranges
Abstract:	Density distribution of the common infaunal bivalves, <i>Macoma balthica</i> and <i>Cerastoderma edule</i> , was studied along the Murman Coast of the Barents Sea during 2002–2010. In both species, abundance was generally higher in West Murman in contrast to East Murman. Highest density of <i>Macoma balthica</i> reaching 1535 ind. m ⁻² was observed in the Kola Inlet. <i>Cerastoderma edule</i> was less abundant; its density rarely exceeded 10 ind. m ⁻² in all but one site, where 282 ind. m ⁻² was registered. Reconstruction of abundance distribution across <i>Macoma balthica</i> European geographic range revealed that it does not match “abundant-centre” pattern, having features of ramped north. On the other hand, distribution of <i>Cerastoderma edule</i> abundance across the range generally follows “abundant-centre” pattern but southern edge populations show relatively higher abundances as compared to those at the north edge (the Barents Sea). Nazarova_et_al_abstract.doc

SCHOLARONE™
Manuscripts

For Review Only

INTRODUCTION

Patterns of species abundance at the biogeographic extremes may provide insights into such essential issues in ecology as identification of borders between adjacent biogeographic regions and causes of species range limits. Analysis of large-scale distribution of species abundance is also a key element of such applications as planning of protected areas and designation of species into regional Red Lists. A widespread paradigm known as “abundant centre hypothesis” (ACH) is widely used to test whether the species abundance is the highest in the range centre and declines towards range edges (Sagarin *et al.*, 2006). Intertidal fauna is an ideal test system for studying latitudinal gradients. In Western Europe, many intertidal and high subtidal marine invertebrates have extended ranges with distinct northern (or north-eastern) and southern (or south-western) limits characterized by diametrically opposed combination of environmental conditions. However, recent studies on wide-spread marine coastal invertebrates have shown that most species do not equally demonstrate the “abundance-centre” pattern, and even closely related species could show different spatial patterns (Sagarin & Gaines, 2002; Rivadeneira *et al.*, 2010).

Empirical approaches to quantify abundance in different parts of the range vary widely. One of the main challenges for examining the abundant centre pattern is a logistical difficulty of sampling throughout the whole species range. Particularly, it could be a problem to get adequate samples from the range edge, where local populations are not stable in time and can spontaneously become extinct and later reoccupy suitable habitats (Sagarin & Gaines, 2002; Sexton *et al.*, 2009).

Species ranges are not stable in time, often shifting, expanding and contracting (Gaston, 2009). Fluctuations of environmental variables, such as climate oscillations, may alter population fitness at the distribution limits, and range shifts can lag behind environmental changes (Pfenninger *et al.*, 2007; Svenning *et al.*, 2008). Marginal populations exhibit greater temporal variability in abundance being near the species' limit of environmental tolerance, especially in extreme years (Gaston, 2009). Since temporal variation can introduce errors in estimation of spatial abundance distribution, it is highly important to apply monitoring data series to improve the understanding of

differences in abundance between central and edge populations (Holt & Keitt, 2000; Johnstone & Chapin, 2003). The spatial distribution and patchy structure of population density are also likely to determine range limits. Edge populations are usually considered to be smaller and more spatially fragmented than populations in the centre of the range (Brown *et al.*, 1995). They may exhibit reductions in diversity and in number of immigrants due to spatial arrangement only, thus resulting in an isolation-by-distance effect (Sexton *et al.*, 2009).

Here, we present the results of the abundance distribution study on *Macoma balthica* and *Cerastoderma edule*, two key bivalve species at the tidal flats of the Barents Sea. *Macoma balthica* is currently regarded as a complex of Pacific *Macoma balthica balthica* and Atlantic *Macoma balthica rubra* subspecies (Nikula *et al.*, 2007). In Europe, this complex spreads from Bay of Biscay up north to the western Kara Sea (Figure 1 A). Southern distribution range limit of *M. balthica* has shifted several hundreds of kilometres to the north during last decades (Beukema *et al.*, 2009). Pure *M. balthica balthica* northern distribution limit belongs to the Varanger Peninsula, while populations in the White and Barents seas are proved to form a broad “hybrid swarm” – their genetic composition is intermediate between Atlantic and Pacific forms (Strelkov *et al.*, 2007). *Cerastoderma edule* range is shaped into at least two genetically different groups of populations along the North Atlantic coast, but separation between these groups does not have a subspecies level (Krakau *et al.*, 2012). Earlier we suggested that low density of individuals is typical for *C. edule* in the Barents Sea, a North-Eastern distribution limit for the species (Genelt-Yanovskiy *et al.*, 2010)(Figure 1A). To study latitudinal clines in geographical abundance distributions, we put together published records of mean *M. balthica* and *C. edule* density across species' European ranges with our own observations from the Barents Sea.

MATERIALS AND METHODS

Study area

Murman Coast is a northern border of the Kola Peninsula with a coastal line of about 700 km,

including numerous bays and fjords. The longest among them, Kola Inlet, divide Murman Coast into two regions: Western Murman and Eastern Murman. Murman environmental conditions are determined by westerly warm Murman Coastal Current (Denisov & Dzhenyuk, 1995). Interaction between Murman Coastal Current and Arctic water backflow causes remarkable cooling gradient from Western to Eastern Murman. Owing to gradient changes in environment, this whole area has transitional biogeographical status – from boreal to arctic zone (Zenkevich, 1963). After E.F. Gourjanova, I.G. Zaks and P.V. Ushakov (Gourjanova & Ushakov, 1929; Gourjanova *et al.*, 1929; Gourjanova *et al.*, 1930), we consider Western Murman, Kola Inlet and Eastern Murman as three distinct regions of Murman Coast.

Sampling

Samples were collected at 18 sites across the Murman Coast (all for *C. edule* and 12 of them for *M. balthica*) (Figure 1, table 1). The sampling took place in July–August, from 2002 to 2010. At each site, tidal zone was divided into three levels according to E.F. Gourjanova, I.G. Zaks and P.V. Ushakov (Gourjanova *et al.*, 1929), and five samples were taken from every level with soft bottom. All samples from one site were further analysed together disregarding the tidal level. At sites inhabited by *M. balthica* alone, 5 cm deep cores with a surface area 0.03 m² were taken; for sites where *M. balthica* and *C. edule* occurred together, 0.1 m² cores were used. All samples were sieved through 1 mm mesh.

All collected bivalves were sorted, identified and counted. Shell length of each individual was measured using calipers to the nearest 0.5 mm. Age of individuals was determined by counting annual growth marks on outer side of the shell. This method of age estimation was used basing on previous studies on *M. balthica* and *C. edule* in the region which confirmed that structural marks on shells of the molluscs in the subarctic seas reflect the annual growth patterns due to the strong seasonal variation in sea water temperature (Agarova, 1979; Maximovich *et al.*, 1992; Maximovich & Gerasimova, 2004; Nazarova *et al.*, 2010).

Statistical analysis

Similar statistical procedures were employed for both species. We started with describing the abundances from each site. We used both medians and means as measures of central tendency and created Tukey boxplots (Tukey, 1976) to illustrate abundance variability. Abundance data from individual samples from one site were used for plotting. Also, year-to-year variability of mean abundance and size-frequency distributions of the species were analysed. Kruskal-Wallis test (Hollander & Wolfe, 1973) was used to compare regional average abundances. The data from published records of mean *C. edule* and *M. balthica* abundance are presented together with our own observations in two maps (Figure 5). For comparisons, we used only mean densities of bigger than spat-sized individuals of both species. Full details on all papers used in assessment of global distribution patterns are listed in electronic supplementary (appendix A).

All calculations were done using R (R-core, 2014). In all cases, statistical significances were tested at 5% probability level.

RESULTS

The abundance of *Macoma balthica* at Western Murman and Kola Inlet sites varied from hundreds to thousands individuals per square meter. *M. balthica* mean density was significantly different in three regions of the Murman coast (Kruskal-Wallis test: $\chi^2 = 17.6$, $p < 0.0001$). Highest mean *M. balthica* density (3350 ± 520 ind. m^{-2}) was observed in 2005 at Cape Abram (Kola Inlet). *M. balthica* density at other Kola Inlet sites (Pala, Retinskoe and Nagornoe) was much alike Western Murman sites (Figures 2A and 3A). The Eastern Murman *M. balthica* mean density was lower than in Western Murman and Kola Inlet and rarely exceeded 100 ind. m^{-2} (Figure 3A). However, two major exceptions from general pattern were found in 2008 (1208 ± 73 ind. m^{-2} in Ivanovskaya bay and 387 ± 77 ind. m^{-2} in Yarnyshnaya bay respectively). In Dalne-Zelenetskaya bay (Eastern Murman) mean long-term (from 2002 to 2008) abundance was 47 ± 3 ind. m^{-2} (Figure 2A).

The most common type of *M. balthica* size structure was bimodal. Either larger or smaller molluscs can prevail though, and the dominant groups may differ at nearby sites even in the same or

subsequent years. For example, in 2008 Yarnyshnaya was dominated by 4 mm molluscs, while 17-18 mm-long *M. balthica* prevailed in nearby Dalne-Zelenetskaya and Gavrilovo (Figure 4A).

Alive individuals of *C. edule* were revealed at 11 sites out of 18. We have not found any alive cockles in Klimkovka, Dolgaya, Tyuva, Gavrilovo, Drozdovka, Ivanovskaya and Gremikha. Density of *C. edule* in Pechenga, Retinskoe, Bolshaya Volokovaya was so low that there was no possibility to estimate it using our sampling scheme, only solitary individuals were present at the tidal flats during our observations. Only empty shells of *C. edule* were found in Gavrilovo and Drozdovka, while Shelpino population became extinct after 2009.

In Ura bay only solitary cockles were present in 2005, but in 2010 abundance increased up to 282 ± 68 ind. m^{-2} . The other site with rather high abundance was Pala bay (45 ± 9 ind. m^{-2} in 2006). In subsequent 2007 cockle abundance at this site dropped, and only solitary cockles were present (Figure 2B). Cockle abundance was notably lower at all other sites (less than 10 ind. m^{-2} at the majority of sites studied) (Figure 3B). Mean abundance at the Western Murman was significantly higher than at other two regions (Kruskal-Wallis test: $\chi^2 = 21.4$, $p < 0.001$).

Unimodal age structures in *C. edule* with prevalence of single generation were typical for most sites (22-24 mm-long cockles in Ura bay, 2005; 26-28 mm in Shelpino, 2005 and 20-24 mm in Pala, 2007) (Figure 4B).

DISCUSSION

Our results generally match with the expectation that the amphi-boreal species *Macoma balthica* should show higher densities in the Barents Sea in comparison to the more temperate-zone *Cerastoderma edule*. This cockle is supposed to be the most sensitive to low air temperatures bivalve on European shores – severe winters were shown to cause mass mortality among adult *Cerastoderma edule* in the Wadden Sea (Strasser *et al.*, 2001). What effects of severe winters are more critical for cockle survival is still unclear, but probably the sudden rise of the air temperature in subsequent spring may be even more crucial than winter freezing of upper bottom layer during low tides (Strasser *et al.*, 2001). The long-term monitoring data from the Wadden Sea provided

evidences that long-term variability in average winter temperatures affects bivalve population dynamics in numerous ways (Beukema *et al.*, 2009; Beukema & Dekker, 2014). Community-level analyses revealed that mild winters in the Wadden Sea give advantage to recruitment in decapods *Crangon crangon* and *Carcinus maenas*, which in turn essentially feed on postlarvae of bivalves *Macoma balthica*, *Cerastoderma edule*, *Mya arenaria* and *Mytilus edulis* (Beukema & Dekker, 2014). The Barents Sea lacks *Carcinus maenas*, the distribution edge of the crab is in Northern Norway. *Crangon crangon* also cannot be regarded as an important predator in the Barents Sea intertidal because it does not seem to form permanent populations at the Murman Coast (Kuznetsov, 1960).

Regional variability in mean abundance found in both *M. balthica* and *C. edule* could be explained by a difference in general environmental conditions between Western and Eastern Murman. Western Murman is an area of numerous fjord-like bays, among which Kola Inlet is the longest (57 km) and Pechenga and Ura bay are more typical – 16 and 19 km long respectively, while Eastern Murman coast is relatively straight. Also, westerly Murman Coastal Current dominates the nearshore hydrodynamics in Eastern Murman, while longer bays of Western Murman are more isolated and thus provide more stable conditions (Denisov & Dzhenyuk, 1995).

According to our data, mean *Macoma* abundance was significantly lower in Eastern Murman sites in comparison to Western Murman and Kola Inlet sites. In 1973 mean density of *M. balthica* at Dalne-Zelenetskaya bay (23 ± 9 ind. m^{-2}) was much alike the present-day values (Agarova *et al.*, 1976). At Western Murman and in Kola Inlet clam densities were comparable with their densities in the White Sea, in the Baltic Sea and in the nearest waters – the northern part of the Norwegian Sea (e.g. Semenova, 1974; Aschan, 1988; Bostrom & Bonsdorf, 2000; Oug, 2001; Varfolomeeva & Naumov, 2013; Rousi *et al.*, 2013; Gerasimova & Maximovich, 2013). Values of mean *M. balthica* abundances similar to the Eastern Murman were typical to the southern part of the Norwegian Sea and the Wadden Sea (Snelli, 1968; Beukema, 1976; Jensen & Jensen, 1985; Reise *et al.*, 1994) (Figure 5A).

Populations of *Macoma balthica* in the South-Western Barents Sea (Murman coast) are certainly not marginal within the species' European part of the range (figures 1A and 5A) – *M. balthica* also occurs in the Kara Sea on the North-East and is quite typical for the White Sea intertidal communities, where it reaches high densities (Semenova, 1974; Denisenko *et al.*, 2003; Strelkov *et al.*, 2007; Gerasimova & Maximovich, 2013).

Densities of *M. balthica* in Ivanovskaya, the easternmost site for this species in our dataset, were dissimilar to other Eastern Murman localities, because at this site *M. balthica* appears to be rather subtidal than intertidal species – it occurs just below the low water level. Subtidal *Macoma balthica* population was previously reported in the Pechora Sea (southeastern part of the Barents Sea) with mean densities of 654 ind. m⁻², which is two times lower than in Ivanovskaya (Denisenko *et al.*, 2003).

Our data indicate that *Macoma balthica* abundance distribution within the species range does not follow the abundant centre pattern (Figure 5A) and is more like a ramped north (for names of distribution shapes see Sagarin & Gaines, 2002). Surprisingly, mean individual densities of bigger than spat-sized *M. balthica* at Western Murman sites were more similar to those in the White and Baltic Seas, and were higher than long-term mean values available for North Sea and Bay of Biscay (see appendix A).

Mean densities of adult *Cerastoderma edule* on the Murman Coast of the Barents Sea being at most sites around 10 ind. m⁻² were generally lower than in other parts of the species range, presumably in the North Sea and Irish Sea (Beukema, 1989; Ivell, 1989; André & Rosenberg, 1991; Bachelet *et al.*, 1992; Ramon, 2003; Dare *et al.*, 2004) (Figure 5B). Long-term mean density of cockles in the Dalne-Zelenetskaya bay was equal to previously reported values (Agarova *et al.*, 1976). Very similar densities to the Barents Sea cockle were previously reported from Tromsø, Northern Norway (Oug, 2001). These data support the hypothesis that low abundance is a typical feature of the northern part of the cockle range. Taking into account that at the southernmost distribution limit, Merja Zerga (Morocco), cockle density may reach 1200 ind. m⁻² with long-term

mean of 638 ind. m⁻² (Gam *et al.*, 2010), abundance pattern could be probably characterized as an “abundance center”, but with a higher abundance at the southern range edge, as compared to the Barents Sea (Figure 5B). Current easternmost *C. edule* local population is situated in Tryashina (~68° 59' N, ~36° 30' E), we did not find any alive cockles further to the east. However, in Drozdovka (~100 km east from Tryashina) we have found two non-fossilized cockle shells, which means that temporary settlement of this species occurred there several years previously to our visit. At five other sites (Pechenga, Retinskoe, Tyuva, Bolshaya Volokovaya, Gavrilovo) it was impossible to estimate cockle density using standard sampling methods and only shells were found, which clearly reflects the fact that Murman coast is a dynamic range edge for *C. edule*. No cockles were also found in the inner part of the Kola Inlet, Cape Abram and Nagornoe (Figure 3B). A possible explanation to the observation may be the known sensitivity of *C. edule* to pollution load (Savari *et al.*, 1989), which should be significant near the city and the port of Murmansk, the largest coastal industrial area in the Barents Sea region.

Cerastoderma edule population dynamics has a very typical feature of periodically occurring mass mortality events, when adult densities sharply decline from thousands of individuals to single cockles per square meter due to a variety of factors. Slow population recovery takes place after that (Strasser *et al.*, 2001; Malham *et al.*, 2012; Callaway *et al.*, 2013). Cockle density in Wadden Sea could reach 54000 ind. m⁻² including individuals of all age groups during periods called after Ducrotoy *et al.* (1991) as “recovery” (Malham *et al.*, 2012). According to our study, the cockle population in Ura bay on Western Murman unambiguously had features of a recovery phase in 2010 after a “crisis” phase observed in 2005 (Figure 2B). During our first visit, only solitary individuals were found across the tidal flat, while in 2010 mean cockle density was 282 ind. m⁻², which is the highest density value for the Barents Sea. Our results on cockle local population recovery in Ura bay support the dependence (at least partial) of Barents sea populations of invertebrates and pelagic fish on larval drift from west (Gjøsaeter, 1995; Mikkelsen & Pedersen, 2004).

ACKNOWLEDGEMENTS

We are grateful to all participants of Saint-Petersburg State University Barents Sea expeditions of 2002–2009 years for their help in the field work and to the administration of Kandalaksha State Natural Reserve for supporting our activity. We also greatly appreciate valuable comments of the anonymous Reviewer #3.

FINANCIAL SUPPORT

The study was partially supported by RFBR grants No.12-04-01507-a, 12-04-10128-k, 13-04-10131-k, and SPbSU Action 2 research project No.1.38.253.2014.

REFERENCES

- Agarova I.Ya.** (1979) Some aspects of the linear growth of bivalve mollusks in the population of *Cerastoderma edule* L. In *Molluscs: main results of studies: abstracts of the annual all-union conference*. Leningrad, pp. 80–82. (in Russian)
- Agarova I.Ya., Voronova M.N., Galtsova V.V., Ioffe B.I., Letova N.V., Streltsov V.E. and Streltsova S.I.** (1976) In *Distribution and ecology of benthic fauna on intertidal sandbank Dalniy Plaj. Ecological studies of soft-bottom littoral*. Apatity, pp. 95–186. (in Russian)
- André C. and Rosenberg R.** (1991) Adult-larval interactions in the suspension-feeding bivalves *Cerastoderma edule* and *Mya arenaria*. *Marine ecology progress series* 71(3), 227–234.
- Aschan M.** (1988) Soft bottom macrobenthos in a Baltic archipelago: Spatial variation and optimal sampling strategy. *Annales Zoologici Fennici* 25 (2), 153–164.
- Bachelet G., Guillou J. and Labourg P. J.** (1992) Adult-larval and juvenile interactions in the suspension-feeding bivalve, *Cerastoderma edule* (L.): field observations and experiments. In *Marine eutrophication and population dynamics*. Olsen and Olsen, Fredensborg, pp.175–182.
- Beukema J. J.** (1976) Biomass and species richness of the macro-benthic animals living on the tidal flats of the Dutch Wadden Sea. *Netherlands Journal of Sea Research* 10(2), 236–261.
- Beukema J. J.** (1989) Long-term changes in macrozoobenthic abundance on the tidal flats of the western part of the Dutch Wadden Sea. *Helgoländer Meeresuntersuchungen* 43(3–4), 405–

415.

- Beukema J. J. and Dekker R.** (2014) Variability in predator abundance links winter temperatures and bivalve recruitment: correlative evidence from long-term data in a tidal flat. *Marine Ecology Progress Series* 513, 1–15.
- Beukema J. J., Dekker R. and Jansen J. M.** (2009) Some like it cold: populations of the tellinid bivalve *Macoma balthica* (L.) suffer in various ways from a warming climate. *Marine Ecology Progress Series* 384, 135–145.
- Boström C. and Bonsdorff E.** (2000) Zoobenthic community establishment and habitat complexity-the importance of seagrass shoot-density, morphology and physical disturbance for faunal recruitment. *Marine Ecology Progress Series* 205, 123–138.
- Brown J. H., Mehlman D. W. and Stevens G. C.** (1995) Spatial variation in abundance. *Ecology* 76(7), 2028–2043.
- Callaway R., Burdon D., Deasey A., Mazik K. and Elliott M.** (2013) The riddle of the sands: how population dynamics explains causes of high bivalve mortality. *Journal of Applied Ecology* 50(4), 1050–1059.
- Dare P.J., Bell M.C., Walker P. and Bannister R.C.A.** (2004) *Historical and current status of cockle and mussel stocks in The Wash*. CEFAS Lowestoft. 85 pp.
- Denisenko S. G., Denisenko N. V., Lehtonen K. K., Andersin A. B. and Laine A. O.** (2003) Macrozoobenthos of the Pechora Sea (SE Barents Sea): community structure and spatial distribution in relation to environmental conditions. *Marine Ecology Progress Series* 258, 109–123.
- Denisov V. V. and Dzhenyuk S. L.** (1995) Chapter 2. Abiotic conditions. Abiotic conditions of Kola Peninsula. In *The Biological Resources of the Coastal zone of the Kola Peninsula. Modern State and Rational Usage*. Russian Academy of Sciences, Apatity, Murmansk, Russia. pp. 10–25. (In Russian)
- Ducrotoy J. P., Rybarczyk H., Souprayen J., Bachelet G., Beukema J., Desprez M. and Ibanez**

- F. (1991) A comparison of the population dynamics of the cockle (*Cerastoderma edule*, L.) in North-Western Europe. In M. Elliott and J.-P. Ducrotoy. (eds.) *Estuaries and coasts: spatial and temporal intercomparisons. International Symposium Series, ECSA-19 Symposium*, Olsen and Olsen, Fredensborg. pp. 173–184
- Gam M., de Montaudouin X. and Bazairi H.** (2010) Population dynamics and secondary production of the cockle *Cerastoderma edule*: A comparison between Merja Zerga (Moroccan Atlantic Coast) and Arcachon Bay (French Atlantic Coast). *Journal of Sea Research* 63(3), 191–201.
- Gaston K. J.** (2009) Geographic range limits of species. *Proceedings of the Royal Society B: Biological Sciences* 276(1661), 1391–1393.
- Genelt-Yanovskiy E., Poloskin A., Granovitch A., Nazarova S. and Strelkov P.** (2010) Population structure and growth rates at biogeographic extremes: A case study of the common cockle, *Cerastoderma edule* (L.) in the Barents Sea. *Marine pollution bulletin* 61(4), 247–253.
- Gerasimova A. V. and Maximovich N. V.** (2013) Age–size structure of common bivalve mollusc populations in the White Sea: the causes of instability. *Hydrobiologia* 706(1), 119–137.
- Gjørøseter H.** (1995) Pelagic fish and the ecological impact of the modern fishing industry in the Barents Sea. *Arctic* 48(3), 267–278.
- Gourjanova E.F., Zaks I.G. and Ushakov P.V.** (1930) A littoral of western Murman. *Studies of the Seas of USSR* 2, 47–52. (In Russian)
- Gourjanova E.F., Zaks I.G. and Ushakov P.V.** (1929) A littoral of Kola Inlet. *Proceedings of the Leningrad Society of Naturalists* 60(2), 17–107. (In Russian)
- Gourjanova E.F. and Ushakov P.V.** (1929) A littoral of eastern Murman. *Studies of the Seas of USSR* 10, 5–40. (In Russian)
- Hollander M. L. and Wolfe D.A.** (1973) *Nonparametric Statistical Methods*. New York: John Wiley and Sons Perry Press.
- Holt R. D. and Keitt T. H.** (2000) Alternative causes for range limits: a metapopulation

perspective. *Ecology Letters* 3(1), 41–47.

Ivell R. (1981) A quantitative study of a *Cerastoderma* – *Nephtys* community in the Limfjord, Denmark, with the special reference to production of *Cerastoderma edule*. *Journal of Molluscan Studies* 47, 147–170.

Jensen K. T. and Jensen J. N. (1985) The importance of some epibenthic predators on the density of juvenile benthic macrofauna in the Danish Wadden Sea. *Journal of Experimental Marine Biology and Ecology* 89(2), 157–174.

Johnstone J. F. and Chapin F. S. (2003). Non-equilibrium succession dynamics indicate continued northern migration of lodgepole pine. *Global Change Biology* 9(10), 1401–1409.

Krakau, M., Jacobsen, S., Jensen, K. T. and Reise, K. (2012). The cockle *Cerastoderma edule* at Northeast Atlantic shores: genetic signatures of glacial refugia. *Marine biology* 159(1), 221–230.

Kuznetsov V.V. (1960) *White Sea and biological features of its flora and fauna*. Moscow – Leningrad: Publishing House of the Academy of Sciences of the USSR. (In Russian)

Malham S. K., Hutchinson T. H. and Longshaw M. (2012) A review of the biology of European cockles (*Cerastoderma* spp.). *Journal of the Marine Biological Association of the United Kingdom* 92(07), 1563–1577.

Maximovich N.V., Gerasimova A. V. and Kunina T. A. (1992) Production characteristics of *Macoma balthica* population in Chupa bay (White Sea). I. Linear growth. *Vestnik Sankt-Peterburgskogo Universiteta. Series 3: Biology* 4 (24), 12–19. (In Russian)

Maximovich N. V. and Gerasimova A. V. (2004) Age determination of the White Sea bivalves by the shell morphology. In *Proceedings of the V Scientific Session of the Marine Biological Station of St. Petersburg State University, St. Petersburg, Russia*. St-Peterburg: St-Peterburg university press, pp. 29–30. (in Russian)

Mikkelsen N. and Pedersen T. (2004) How can the stock recruitment relationship of the Barents Sea capelin (*Mallotus villosus*) be improved by incorporating biotic and abiotic factors? *Polar*

Research 23(1), 19–26.

- Nazarova S. A., Genelt-Yanovsky E. A. and Maximovich N. V.** (2010) Linear growth of *Macoma balthica* in the Murmansk tidal zone (the Barents Sea) *Vestnik Sankt-Peterburgskogo Universiteta. Series 3: Biology* 4, 35–43. (In Russian)
- Nikula R., Strelkov P. and Väinölä R.** (2007) Diversity and trans-arctic invasion history of mitochondrial lineages in the north atlantic *Macoma balthica* complex (Bivalvia: Tellinidae). *Evolution* 61(4), 928–941.
- Oug E.** (2001) Polychaetes in intertidal rocky and sedimentary habitats in the region of Tromsø, northern Norway. *Sarsia* 86(1), 75–83.
- Pfenninger M., Nowak C. and Magnin F.** (2007) Intraspecific range dynamics and niche evolution in *Candidula* land snail species. *Biological Journal of the Linnean Society* 90(2), 303–317.
- Ramón M.** (2003) Population dynamics and secondary production of the cockle *Cerastoderma edule* (L.) in a backbarrier tidal flat in the Wadden Sea. *Scientia Marina* 67(4), 429–443.
- R Core Team** (2014) *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- Reise K., Herre E. and Sturm M.** (1994) Biomass and abundance of macrofauna in intertidal sediments of Königshafen in the northern Wadden Sea. *Helgoländer Meeresuntersuchungen* 48(2-3), 201–215.
- Rivadeneira M. M., Hernáez P., Antonio Baeza J., Boltana S., Cifuentes M., Correa C. and Thiel M.** (2010) Testing the abundant-centre hypothesis using intertidal porcelain crabs along the Chilean coast: linking abundance and life-history variation. *Journal of biogeography* 37(3), 486–498.
- Sagarin R. D., and Gaines S. D.** (2002) The ‘abundant centre’ distribution: to what extent is it a biogeographical rule? *Ecology letters* 5(1), 137–147.
- Sagarin R. D., Gaines S. D. and Gaylord B.** (2006) Moving beyond assumptions to understand

abundance distributions across the ranges of species. *Trends in Ecology and Evolution* 21(9), 524–530.

Savari A., Sylvestre C., Sheader M., Le Gal Y. and Lockwood A.P.M. (1989) Stress studies on the common cockle (*Cerastoderma edule* L.) in Southampton Water. Topics in Marine biology. *Scientia Marina* 53(2-3), 729-735.

Semenova N.L. (1974) The distribution of the bivalve *Macoma balthica* (L.) in some inlets of Kandalaksha Bay of the White Sea. *Proceedings of the White Sea Biological Station of the Moscow State University* 4, 87–102. (In Russian)

Sexton J. P., McIntyre P. J., Angert A. L. and Rice K. J. (2009) Evolution and ecology of species range limits. *Annual Review of Ecology, Evolution, and Systematics* 40, 415–436

Sneli J. A. (1968) The intertidal distribution of polychaetes and molluscs on a muddy shore in Nord-Møre, Norway. *Sarsia* 31(1), 63–68.

Strasser M., Reinwald T. and Reise K. (2001) Differential effects of the severe winter of 1995/96 on the intertidal bivalves *Mytilus edulis*, *Cerastoderma edule* and *Mya arenaria* in the Northern Wadden Sea. *Helgoland Marine Research* 55(3), 190–197.

Strelkov P., Nikula R. and Väinölä R. (2007) *Macoma balthica* in the White and Barents Seas: properties of a widespread marine hybrid swarm (Mollusca: Bivalvia). *Molecular ecology* 16(19), 4110–4127.

Svenning J. C., Normand S. and Kageyama, M. (2008) Glacial refugia of temperate trees in Europe: insights from species distribution modelling. *Journal of Ecology* 96(6), 1117–1127.

Tukey J. W. (1976) *Exploratory data analysis*. Massachusetts: Addison-Wesley.

Varfolomeeva M. and Naumov A. (2013) Long-term temporal and spatial variation of macrobenthos in the intertidal soft-bottom flats of two small bights (Chupa Inlet, Kandalaksha Bay, White Sea). *Hydrobiologia* 706(1), 175–189.

Zenkevich L.A. (1963) *Biology of the Seas of the U.S.S.R.* Moscow. Translated from Russian to English by S. Botsharskaya. New York: Interscience Publishers.

Table 1. Investigation sites and sampling details

Region	Site (code)	Latitude, longitude	Observation years
Western Murian	Pechenga (PG)	69.58, 31.27	2005
	Ura bay (UR)	68.99, 36.51	2010
Kola Inlet	Pala bay (PL)	69.19, 33.37	2006-2007
	Retinskoe (RT)	69.11, 33.38	2005
	Cape Abram (AB)	68.98, 33.03	2008
	Severnoe Nagornoe (NG)	68.90, 33.06	2005
	Tyuva (TU)	69.17, 33.63	2005
	Bolshaya Volokovaya (BV)	69.27, 33.62	2005
	Klimkovka (KL)	69.23, 34.64	2009
Eastern Murian	Dolgaya (DO)	69.17, 33.63	2009
	Gavrilovo (GV)	69.17, 35.86	2008
	Yarnyshnaya (YA)	69.09, 36.05	2004 - 2008
	Dalne-Zelenetskaya (DZ)	69.11, 36.10	2002 - 2009
	Shelpino (SH)	69.10, 36.21	2005, 2008
	Porchnikha (PR)	69.08, 36.25	2006 - 2007
	Tryashina (TR)	68.99, 36.51	2008
	Drozdovka (DR)	68.30, 38.44	2008
	Ivanovskaya subtidal (IV)	68.29, 38.71	2008
	Gremikha (GR)	68.03, 39.85	2009




Fig. 1. Distribution range map of *Macoma balthica* (European part) and *Cerastoderma edule* (A) and study area map (B). See table 1 for sample sites details.
175x85mm (300 x 300 DPI)




Fig. 2. Year-to-year variability of *Macoma balthica* (A) and *Cerastoderma edule* (B) mean abundance (lgN). Symbol colour corresponds to the region: Western Murman – solid grey, Kola Inlet – solid black and Eastern Murman – open symbols.
101x208mm (300 x 300 DPI)




Fig. 3. Mean abundance (N, ind.m⁻²) of *Macoma balthica* (A) and *Cerastoderma edule* (B) in the Barents sea. Samples are arranged from West to East. At the box-plots box marked Q1, median and Q3, whiskers marked 1.5 IQR. Mean abundance indicated at the top.
101x210mm (300 x 300 DPI)




Fig. 4. Size frequency distributions of *Macoma balthica* (A) into 1 mm classes and *Cerastoderma edule* (B) into 2 mm classes.
211x211mm (300 x 300 DPI)




Fig. 4. Size frequency distributions of *Macoma balthica* (A) into 1 mm classes and *Cerastoderma edule* (B) into 2 mm classes.
211x211mm (300 x 300 DPI)




Fig. 5. Geographic abundance distributions of *Macoma balthica* (A) and *Cerastoderma edule* (B). Area of circles is proportional to the mean abundance. For sources of data see Appendix A.
109x84mm (300 x 300 DPI)




Fig. 5. Geographic abundance distributions of *Macoma balthica* (A) and *Cerastoderma edule* (B). Area of circles is proportional to the mean abundance. For sources of data see Appendix A.
108x139mm (300 x 300 DPI)