

Journal of the
MARINE BIOLOGICAL ASSOCIATION
of the United Kingdom



CAMBRIDGE
UNIVERSITY PRESS

**Abundance distribution patterns of intertidal bivalves
Macoma balthica and *Cerastoderma edule* at the Murman
coast tidal flats (the Barents Sea)**

Journal:	<i>Journal of the Marine Biological Association of the United Kingdom</i>
Manuscript ID:	JMBA-11-14-SIEMBS-0452.R2
Manuscript Type:	Special Issue - EMBS European Marine Biology Symposium
Date Submitted by the Author:	n/a
Complete List of Authors:	Nazarova, Sophia; St-Petersburg State University, Ichthyology and Hydrobiology ; Russian State Pedagogical University, Zoology Shunkina, Ksenia; Zoological Institute RAS, Laboratory of Evolutional Morphology Genelt-Yanovskiy, Evgeny; Saint-Petersburg State University, Ichthyology and hydrobiology; Baltic Fund for Nature,
Keywords:	<i>Cerastoderma edule</i> , cockle, <i>Macoma balthica</i> , Barents sea, Geographic abundance distribution, latitudinal variation, species ranges
Abstract:	<p>Density distribution of the common infaunal bivalves, <i>Macoma balthica</i> and <i>Cerastoderma edule</i>, was studied along the Murman Coast of the Barents Sea during 2002–2010. In both species, abundance was generally higher in West Murman in contrast to East Murman. Highest density of <i>Macoma balthica</i> reaching 1535 ind. m⁻² was observed in the Kola Inlet. <i>Cerastoderma edule</i> was less abundant; its density rarely exceeded 10 ind. m⁻² in all but one site, where 282 ind. m⁻² was registered. Reconstruction of abundance distribution across <i>Macoma balthica</i> European geographic range revealed that it does not match “abundant-centre” pattern, having features of ramped north. On the other hand, distribution of <i>Cerastoderma edule</i> abundance across the range generally follows “abundant-centre” pattern but southern edge populations show relatively higher abundances as compared to those at the north edge (the Barents Sea).</p> <p>Nazarova_et_al_abstract.doc</p>

SCHOLARONE™
Manuscripts

For Review Only

1 INTRODUCTION

2 Patterns of species' abundance at the biogeographic extremes may provide insights into such
3 essential issues in ecology as identification of borders between adjacent biogeographic regions
4 and causes of species range limits. Large-scale distribution of species abundance is also a key
5 element of applied sciences such as planning of protected areas and designation of species into
6 regional Red Lists. A widespread paradigm, known as “abundant-centre hypothesis” (ACH) is
7 widely used to test whether the highest species abundance belongs to the range centre and
8 declines towards range edges (Sagarin et al., 2006). Intertidal fauna is an ideal test system for
9 studying latitudinal gradients. In Western Europe, many intertidal and high subtidal marine
10 invertebrates have extended ranges with distinct north (or north-eastern) and south (or south-
11 western) limits characterized by diametrically opposed combination of environmental conditions.
12 Recent studies on wide-spread marine coastal invertebrates have shown that abundance centre
13 pattern does not apply to most species equally, and even closely related species could show
14 different spatial patterns (Sagarin & Gaines, 2002; Rivadeneira et al., 2010).

15 Empirical approaches to quantify abundance in different parts of the range vary widely. One of
16 the main challenges for examining the abundant centre pattern is a logistical difficulty of
17 sampling throughout the whole species range. Particularly, it could be a problem to get adequate
18 samples from the range edge, where local populations are not stable in time and can
19 spontaneously become extinct and later reoccupy suitable habitats (Sagarin & Gaines, 2002;
20 Sexton et al., 2009).

21 Species ranges are not stable in time, often shifting, expanding and contracting (Gaston, 2009).
22 Fluctuations of environmental variables, such as climate oscillations, may alter population fitness
23 at the distribution limits and range shifts can lag behind environmental changes (Pfenninger et
24 al., 2007; Svenning et al., 2008). Marginal populations exhibit greater temporal variability in
25 abundance being near the species' limit of environmental tolerance, especially in extreme years
26 (Gaston, 2009). Since temporal variation can introduce errors in estimation of spatial abundance

27 distribution, it is highly important to apply monitoring data series to improve the understanding
28 of differences in abundance between central and edge populations (Holt & Keitt, 2000;
29 Johnstone & Chapin, 2003). The spatial distribution and patchy structure of population density
30 are also likely to determine range limits. Edge populations usually considered to be smaller and
31 more spatially fragmented than populations in the centre of the range (Brown et al., 1995). They
32 may exhibit reductions in diversity and in number of immigrants due to spatial arrangement only,
33 thus resulting in an isolation-by-distance effect (Sexton et al., 2009).

34 Here, we present the results of the study of abundance distribution of *Macoma balthica* and
35 *Cerastoderma edule*, two key bivalve species at the tidal flats of the Barents Sea. *Macoma*
36 *balthica* is currently regarded as a complex of Pacific *Macoma balthica balthica* and Atlantic
37 *Macoma balthica rubra* subspecies (Nikula et al., 2007). In Europe this complex spreads from
38 Bay of Biscay up north to the western Kara sea (Figure 1 A). Southern distribution range limit of
39 *Macoma* has shifted several hundreds of kilometres north during last decades (Beukema et al.,
40 2009). Pure *M. balthica balthica* northern distribution limit belongs to the Varanger Peninsula,
41 while populations in the White and Barents seas are proved to form a broad “hybrid swarm” -
42 their genetic composition is intermediate between Atlantic and Pacific forms (Strelkov et al.,
43 2007). *Cerastoderma edule* range is shaped into at least two genetically different groups of
44 populations along the North Atlantic coast, but separation between these groups does not have a
45 subspecies level (Krakau et al., 2012). Earlier we suggested that low density of individuals is
46 typical for *C. edule* in the Barents Sea, a North-Eastern distribution limit for the species (Genelt-
47 Yanovskiy et al., 2010)(Figure 1A). To study latitudinal clines in geographical abundance
48 distributions, we put together published records of mean *M. balthica* and *C. edule* density across
49 species' European ranges with our own observations from the Barents Sea.

50 MATERIALS AND METHODS

51 Study area

52 Murman Coast is a northern border of the Kola Peninsula with a coastal line of about 700 km,

including numerous bays and fjords. The longest among them, Kola Inlet, divide Murman Coast into two regions: Western Murman and Eastern Murman. Murman environmental conditions are determined by westerly warm Murman Coastal Current (Denisov & Dzhenyuk, 1995). Interaction between Murman Coastal Current and Arctic water backflow causes remarkable cooling gradient from West to East Murman. Owing to gradient changes in environment this whole area has transitional biogeographical status – from boreal to arctic zone (Zenkevich, 1963). After E.F. Gourjanova, I.G. Zaks and P.V. Ushakov (Gourjanova & Ushakov, 1929; Gourjanova et al., 1929; Gourjanova et al., 1930) we consider Western Murman, Kola Inlet and Eastern Murman as three distinct regions of Murman Coast.

Sampling

Samples were collected at 18 sites across the Murman Coast (all for *C. edule* and 12 of them for *M. balthica*) (Figure 1, table 1). The sampling took place in July–August, from 2002 to 2010. At sites inhabited by *Macoma* alone, 5 cm deep cores with a surface area 0.03 m² were taken; for sites where *Macoma* and *Cerastoderma* occurred together, 0.1 m² corers were used. All samples were sieved through 1 mm mesh. Each soft-bottom tidal level was characterized with five samples at all sites studied.

All collected bivalves were sorted, identified and counted. Shell length of each individual was measured using calipers to the nearest 0.5 mm. Age of individuals was determined by counting annual growth marks on outer side of the shell. This method of age estimation was used basing on previous studies on *Macoma* and *Cerastoderma* in the region, confirming that structural marks on shells of the molluscs in the subarctic seas reflect the annual growth patterns, due to the strong seasonal variation in sea water temperature (Agarova, 1979; Maximovich et al., 1992; Maximovich & Gerasimova, 2004, Nazarova et al, 2010).

Statistical analysis

Similar statistical procedures were employed for both species. We started with describing the abundances from each site. We used both medians and means as measures of central tendency

79 and created Tukey boxplots (Tukey, 1976) to illustrate abundance variability. Data on abundance
80 from individual samples from one site was used for plotting. Also year-to-year variability of
81 mean abundance and size-frequency distributions of the species were analysed. Kruskal-Wallis
82 test (Hollander, Wolfe, 1973) was used to compare regional average abundances. The data from
83 published records of mean *C. edule* and *M. balthica* abundance are presented together with our
84 own observations in two maps (Figure 5). For the comparisons we used only mean densities of
85 bigger than spat-sized individuals of *Macoma* and *Cerastoderma*. Full details on all papers used
86 in assessment of global distribution patterns are listed in electronic supplementary (appendix A).
87 All calculations were done with R (R-core, 2014). In all cases, statistical significances were
88 tested at 5% probability level.

89 RESULTS

90 The abundance of *Macoma balthica* at Western Murman and Kola Inlet sites varied from
91 hundreds to thousands individuals per square meter. *M. balthica* mean density was significantly
92 different at three regions of the Murman coast (Kruskal-Wallis test: $\chi^2 = 17.6$, $p < 0.0001$).
93 Highest mean *M. balthica* density (3350 ± 520 ind. m^{-2}) was observed in 2005 at Cape Abram
94 (Kola Inlet). *M. balthica* density at other Kola Inlet sites (Pala, Retinskoe and Nagornoe) was
95 much alike Western Murman sites (figures 2A and 3A). The Eastern Murman *M. balthica* mean
96 density was lower than in Western Murman and Kola Inlet and rarely exceeded 100 ind. m^{-2}
97 (Figure 3A). However, two major exceptions from general pattern were found in 2008
98 (1208 ± 73 ind. m^{-2} in Ivanovskaya bay and 387 ± 77 ind. m^{-2} in Yarnyshnaya bay respectively). In
99 Dalne-Zelenetskaya bay (Eastern Murman) mean long-term (from 2002 to 2008) abundance was
100 47 ± 3 ind. m^{-2} (Figure 2A).

101 The most common type of *M. balthica* size structure was bimodal, but both cases of prevalence
102 of larger and smaller molluscs could occur in same or subsequent year. So, in 2008 Yarnyshnaya
103 was dominated by 4 mm molluscs, while 17-18 mm-long *M. balthica* prevailed in nearby Dalne-
104 Zelenetskaya and Gavrilovo (Figure 4A).

105 Alive individuals of *C. edule* were revealed at 11 sites out of 18. We have not found any alive
106 cockles in Klimkovka, Dolgaya, Tyuva, Gavrilovo, Drozdovka, Ivanovskaya and Gremikha.
107 Density of *C. edule* in Pechenga, Retinskoe, Bolshaya Volokovaya was so low that there was no
108 possibility to estimate it using our sampling scheme, only solitary individuals were present at the
109 tidal flats during our observations. Only empty shells of *C. edule* were found in Gavrilovo and
110 Drozdovka, while Shelpino population became extinct after 2009.

111 In Ura bay only solitary cockles were present in 2005, but in 2010 abundance increased up to
112 282 ± 68 ind. m^{-2} . The other site with rather high abundance was Pala bay (45 ± 9 ind. m^{-2} in 2006).
113 In subsequent 2007 cockle abundance at this site had dropped, and only solitary cockles were
114 present (Figure 2B). Cockle abundance was notably lower at all other sites (less than 10 ind. m^{-2}
115 at the majority of sites studied) (Figure 3B). Mean abundance at the Western Murman was
116 significantly higher than at other two regions (Kruskal-Wallis test: $\chi^2 = 21.4$, $p < 0.001$).

117 Unimodal age structures in with prevalence of single *C. edule* generation were typical for most
118 sites (22-24 mm-long cockles in Ura bay, 2005; 26-28 mm in Shelpino, 2005 and 20-24 mm in
119 Pala, 2007) (Figure 4B).

120 DISCUSSION

121 Our results generally match with the expectation that the amphi-boreal species *Macoma*
122 *balthica* will show higher densities in the Barents Sea in comparison to the more temperate-zone
123 *Cerastoderma edule*. This cockle is supposed to be the most sensitive bivalve to low air
124 temperatures on European shores - severe winters were shown to cause mass mortalities among
125 adult *Cerastoderma edule* in the Wadden Sea (Strasser et al., 2001). What effects of severe
126 winters are more critical for cockle survival is still unclear, but probably the sudden rise of the
127 air temperature in subsequent spring may be even more crucial than winter freezing of upper
128 bottom layer during low tides (Strasser et al., 2001). The long-term monitoring data from the
129 Wadden Sea provided evidences that long-term variability in average winter temperatures affects
130 bivalve population dynamics in numerous ways (Beukema et al., 2009; Beukema & Dekker,

2014). Community-level analyses revealed that mild winters in the Wadden Sea give advantage to recruitment in decapods *Crangon crangon* and *Carcinus maenas*, which in turn essentially feed on postlarvae of bivalves *Macoma balthica*, *Cerastoderma edule*, *Mya arenaria* and *Mytilus edulis* (Beukema & Dekker, 2014). The Barents Sea is lacking *Carcinus maenas*. The distribution edge is in Northern Norway. *Crangon crangon* also cannot be regarded as an important predator in the Barents Sea intertidal because it does not seem to form permanent populations in the Murman Coast (Kuznetsov, 1960).

Regional variability in mean abundance, found in both *M. balthica* and *C. edule* could be interpreted by a difference in general environmental conditions between Western and Eastern Murman. West Murman is an area of numerous fjord-like bays, among which Kola Inlet is the longest (57 km) and Pechenga and Ura bay are more typical – 16 and 19 km long respectively, while Eastern Murman coast is relatively straight. Also, westerly Murman Coastal Current dominates the nearshore hydrodynamics in Eastern Murman, while longer bays of Western Murman are more isolated and thus provide more stable conditions (Denisov & Dzhenyuk, 1995).

According to our data, mean *Macoma* abundance was significantly lower in Eastern Murman sites in comparison to Western Murman and Kola Inlet sites. In 1973 mean density of *M. balthica* at Dalne-Zelenetskaya bay (23 ± 9 ind. m^{-2}) was much alike the present-day values (Agarova et al., 1976). At Western Murman and in Kola Inlet clam densities were comparable with their densities in the White Sea, in the Baltic Sea and in the nearest waters – the northern part of the Norwegian Sea (for example Semenova, 1974; Aschan, 1988; Bostrom & Bonsdorf, 2000; Oug, 2001; Varfolomeeva & Naumov, 2013; Rousi et al., 2013; Gerasimova & Maximovich, 2013). Similar to the Eastern Murman values of mean *Macoma* abundances were typical to the southern part of the Norwegian Sea and the Wadden Sea (Sneli, 1968; Beukema, 1976; Jensen & Jensen, 1985; Reise et al., 1994) (Figure 5A).

Populations of *Macoma balthica* in the South-Western Barents Sea (Murman coast) are

157 certainly not marginal within the species' European part of the range (figures 1A and 5A) –
158 *Macoma* also occurs in the Kara Sea on the North-East and quite typical for the White Sea
159 intertidal communities, where it reaches high densities (Semenova, 1974; Denisenko et al., 2003;
160 Strelkov et al., 2007; Gerasimova & Maximovich, 2013).

161 Densities of *M. balthica* in Ivanovskaya, the easternmost site for this species in our dataset,
162 were dissimilar with other Eastern Murman localities, because in this site *Macoma* appeared to
163 be more subtidal than intertidal species – it occurs just below the low water level. Subtidal
164 *Macoma balthica* population was previously reported in the Pechora Sea (southeastern part of
165 the Barents Sea) with mean densities of 654 ind. m⁻², which is two times lower than in
166 Ivanovskaya (Denisenko et al., 2003).

167 Our data suggest that *Macoma balthica* abundance distribution within the species range does
168 not follow the “abundant centre” (Figure 5A) and is more like a “ramped north” (for names of
169 distribution shapes see Sagarin & Gaines, 2002). Surprisingly, mean individual densities of
170 bigger than spat-sized *Macoma* in Western Murman sites were more similar to those in the White
171 and Baltic Seas, and were higher than long-term mean values available for North Sea and Bay of
172 Biscay (see appendix A).

173 Mean densities of adult *Cerastoderma edule* on the Murman Coast of the Barents Sea, varying
174 in most sites around 10 ind. m⁻² were generally lower than in other parts of the species range,
175 presumably in the North Sea and Irish Sea (Beukema, 1989; Ivell, 1989; André & Rosenberg,
176 1991; Bachelet et al., 1992; Ramon, 2003; Dare et al., 2004) (Figure 5B). Long-term mean
177 density of cockles in the Dalne-Zelenetskaya bay was equal to previously reported values
178 (Agarova et al., 1976). Very similar densities to the Barents Sea cockle were previously reported
179 from Tromsø, North Norway (Oug, 2001). These data support the hypothesis that low abundance
180 is a typical feature of the northern part of the cockle range. Taking into account that at the
181 southernmost distribution limit, Merja Zerga (Morocco), cockle density may reach 1200 ind. m⁻²
182 with long-term mean of 638 ind. m⁻² (Gam et al., 2010), abundance pattern could be probably

183 characterized as an “abundance center”, but with a higher abundance at the southern range edge,
184 as compared to the Barents Sea.(Figure 5B). Current easternmost *Cerastoderma* local population
185 is located in Tryashina (~68° 59’ N, ~36° 30’ E), we did not find any alive cockles further east.
186 However, in Drozdovka (~100 km east from Tryashina) we have found two non-fossilized cockle
187 shells, which mean that temporary settlement of this species occurred here some years ago.
188 Murman coast clearly is a dynamic range edge for *Cerastoderma*, and thus in five other sites
189 (Pechenga, Retinskoe, Tyuva, Bolshaya Volokovaya, Gavrilovo) it was impossible to estimate
190 cockle density using standard sampling methods and only shells were found. No cockles were
191 also found in the inner part of the Kola Inlet, Cape Abram and Nagornoe (Figure 3B). We
192 suppose this could be explained by the known sensitivity of *Cerastoderma* to pollution load
193 (Savari et al., 1989), which should be significant near the city and port of Murmansk, largest
194 coastal industrial area in the Barents Sea region.

195 *Cerastoderma edule* population dynamics has a very typical feature of periodically occurring
196 mass mortality events, when adult densities sharply decline from thousands of individuals to
197 single cockles per square meter due to a variety of factors. After that slow population recovery
198 takes place (Strasser et al., 2001; Malham et al., 2012; Callaway et al., 2013). In the Wadden Sea
199 during periods, called after Ducrotoy et al. (1991) as “recovery”, cockle density may reach
200 54000 ind. m⁻² including individuals of all age groups (Malham et al., 2012). According to our
201 study, the cockle population in Ura bay on Western Murman unambiguously had features of a
202 recovery phase in 2010 after “crisis” phase observed in 2005 (Figure 2B). During our first visit,
203 only solitary individuals were found across the tidal flat, while in 2010 mean cockle density was
204 282 ind. m⁻², which is the highest density value for the Barents Sea. Our results on cockle local
205 population recovery in Ura Bay support the dependence (at least partial) of Barents sea
206 populations of invertebrates and pelagic fish on larval drift from west (Gjøsaeter, 1995;
207 Mikkelsen & Pedersen, 2004).

208 ACKNOWLEDGEMENTS

We are grateful to all participants of Saint-Petersburg State University Barents Sea expeditions of 2002–2009 years for their help in the field work and to the administration of Kandalaksha State Natural Reserve for supporting our activity. We also greatly appreciate valuable comments of the anonymous Reviewer 3.

FINANCIAL SUPPORT

The study was partially supported by RFBR grants No.12-04-01507-a, 12-04-10128-k, 13-04-10131-k, and SPbSU Action 2 research project No.1.38.253.2014.

REFERENCES

- Agarova I.Ya.** (1979) Some aspects of the linear growth of bivalve mollusks in the population of *Cerastoderma edule* L. *Molluscs: main results of studies: abstracts of the annual all-union conference*. Leningrad 6, 80–82. (in Russian)
- Agarova I.Ya., Voronova M.N., Galtsova V.V., Ioffe B.I., Letova N.V., Streltsov V.E. and Streltsova S.I.** (1976) Distribution and ecology of benthic fauna on intertidal sandbank Dalniy Plaj. *Ecological studies of soft-bottom littoral*. Apatity, 95–186. (in Russian)
- André C. and Rosenberg R.** (1991) Adult-larval interactions in the suspension-feeding bivalves *Cerastoderma edule* and *Mya arenaria*. *Marine ecology progress series*. 71(3), 227–234.
- Aschan M.** (1988) Soft bottom macrobenthos in a Baltic archipelago: Spatial variation and optimal sampling strategy. In *Annales Zoologici Fennici* 25 (2), 153–164.
- Bachelet G., Guillou J., and Labourg P. J.** (1992) Adult-larval and juvenile interactions in the suspension-feeding bivalve, *Cerastoderma edule* (L.): field observations and experiments. In *Marine eutrophication and population dynamics*. Olsen and Olsen, Fredensborg, pp.175–182.
- Beukema J. J.** (1976) Biomass and species richness of the macro-benthic animals living on the tidal flats of the Dutch Wadden Sea. *Netherlands Journal of Sea Research*, 10(2), 236–261.
- Beukema J. J.** (1989) Long-term changes in macrozoobenthic abundance on the tidal flats of the western part of the Dutch Wadden Sea. *Helgoländer Meeresuntersuchungen* 43(3–4), 405–

- 235 415.
- 236 **Beukema J. J., and Dekker R.** (2014) Variability in predator abundance links winter
237 temperatures and bivalve recruitment: correlative evidence from long-term data in a tidal
238 flat. *Marine Ecology Progress Series* 513, 1–15.
- 239 **Beukema J. J., Dekker R., and Jansen J. M.** (2009) Some like it cold: populations of the
240 tellinid bivalve *Macoma balthica* (L.) suffer in various ways from a warming climate.
241 *Marine Ecology Progress Series* 384, 135–145.
- 242 **Boström C. and Bonsdorff E.** (2000) Zoobenthic community establishment and habitat
243 complexity-the importance of seagrass shoot-density, morphology and physical disturbance
244 for faunal recruitment. *Marine Ecology Progress Series* 205, 123–138.
- 245 **Brown J. H., Mehlman D. W., and Stevens G. C.** (1995) Spatial variation in
246 abundance. *Ecology* 76(7), 2028–2043.
- 247 **Callaway R., Burdon D., Deasey A., Mazik K. and Elliott M.** (2013) The riddle of the sands:
248 how population dynamics explains causes of high bivalve mortality. *Journal of Applied*
249 *Ecology* 50(4), 1050–1059.
- 250 **Dare P.J., Bell M.C., Walker P. and Bannister R.C.A.** (2004) *Historical and current status of*
251 *cockle and mussel stocks in The Wash. CEFAS Lowestoft.* 85 pp.
- 252 **Denisenko S. G., Denisenko N. V., Lehtonen K. K., Andersin A. B., and Laine A. O.** (2003)
253 Macrozoobenthos of the Pechora Sea (SE Barents Sea): community structure and spatial
254 distribution in relation to environmental conditions. *Marine Ecology Progress Series* 258,
255 109–123.
- 256 **Denisov V. V., Dzhenyuk S. L.** (1995) Chapter 2. Abiotic conditions. Abiotic conditions of Kola
257 Peninsula. In: *The Biological Resources of the Coastal zone of the Kola Peninsula. Modern*
258 *State and Rational Usage.* Russian Academy of Sciences, Apatity, Murmansk, Russia. pp.
259 10–25. (In Russian)
- 260 **Ducrotoy J. P., Rybarczyk H., Souprayen J., Bachelet G., Beukema J., Desprez M., ... and**

- 261 **Ibanez F.** (1991) A comparison of the population dynamics of the cockle (*Cerastoderma*
262 *edule*, L.) in North-Western Europe. In: M. Elliott and J.-P. Ducrotoy. (eds.) *Estuaries and*
263 *coasts: spatial and temporal intercomparisons. International Symposium Series, ECSA-19*
264 *Symposium*, Olsen and Olsen, Fredensborg. pp. 173–184
- 265 **Gam M., de Montaudouin X., and Bazairi H.** (2010) Population dynamics and secondary
266 production of the cockle *Cerastoderma edule*: A comparison between Merja Zerga
267 (Moroccan Atlantic Coast) and Arcachon Bay (French Atlantic Coast). *Journal of Sea*
268 *Research*, 63(3), 191–201.
- 269 **Gaston K. J.** (2009) Geographic range limits of species. *Proceedings of the Royal Society B:*
270 *Biological Sciences*, 276(1661), 1391–1393.
- 271 **Genelt-Yanovskiy E., Poloskin A., Granovitch A., Nazarova S., and Strelkov P.** (2010)
272 Population structure and growth rates at biogeographic extremes: A case study of the
273 common cockle, *Cerastoderma edule* (L.) in the Barents Sea. *Marine pollution bulletin*,
274 61(4), 247–253.
- 275 **Gerasimova A. V. and Maximovich N. V.** (2013) Age–size structure of common bivalve
276 mollusc populations in the White Sea: the causes of instability. *Hydrobiologia*, 706(1), 119–
277 137.
- 278 **Gjøsaeter H.** (1995) Pelagic fish and the ecological impact of the modern fishing industry in the
279 Barents Sea. *Arctic* 48(3), 267–278.
- 280 **Gourjanova E.F., Zaks I.G. and Ushakov P.V.** (1930) A littoral of western Murman. *Studies of*
281 *the Seas of USSR* 2, 47–52. (In Russian)
- 282 **Gourjanova E.F., Zaks I.G. and Ushakov P.V.** (1929) A littoral of Kola Inlet. *Proceedings of*
283 *the Leningrad Society of Naturalists* 60(2), 17–107. (In Russian)
- 284 **Gourjanova E.F. and Ushakov P.V.** (1929) A littoral of eastern Murman. *Studies of the Seas of*
285 *USSR* 10, 5–40. (In Russian)
- 286 **Hollander M. L. and Wolfe D.A.** (1973) *Nonparametric Statistical Methods*. New York: John

- 287 Wiley and Sons Perry Press.
- 288 **Holt R. D. and Keitt T. H.** (2000) Alternative causes for range limits: a metapopulation
289 perspective. *Ecology Letters* 3(1), 41–47.
- 290 **Ivell R.** (1981) A quantitative study of a *Cerastoderma* – *Nephtys* community in the Limfjord,
291 Denmark, with the special reference to production of *Cerastoderma edule*. *Journal of*
292 *Molluscan Studies* 47, 147–170.
- 293 **Jensen K. T. and Jensen J. N.** (1985) The importance of some epibenthic predators on the
294 density of juvenile benthic macrofauna in the Danish Wadden Sea. *Journal of Experimental*
295 *Marine Biology and Ecology* 89(2), 157–174.
- 296 **Johnstone J. F. and Chapin F. S.** (2003). Non-equilibrium succession dynamics indicate
297 continued northern migration of lodgepole pine. *Global Change Biology* 9(10), 1401–1409.
- 298 **Krakau, M., Jacobsen, S., Jensen, K. T., and Reise, K.** (2012). The cockle *Cerastoderma*
299 *edule* at Northeast Atlantic shores: genetic signatures of glacial refugia. *Marine biology*,
300 159(1), 221–230.
- 301 **Kuznetsov V.V.** (1960) *White Sea and biological features of its flora and fauna*. Moscow –
302 Leningrad: Publishing House of the Academy of Sciences of the USSR. (In Russian)
- 303 **Malham S. K., Hutchinson T. H. and Longshaw M.** (2012) A review of the biology of
304 European cockles (*Cerastoderma* spp.). *Journal of the Marine Biological Association of the*
305 *United Kingdom* 92(07), 1563–1577.
- 306 **Maximovich N.V., Gerasimova A. V. and Kunina T. A.** (1992) Production characteristics of
307 *Macoma balthica* population in Chupa bay (White Sea). I. Linear growth. *Vestnik Sankt-*
308 *Peterburgskogo Universiteta. Series 3: Biology* 4 (24), 12–19. (In Russian)
- 309 **Maximovich N. V. and Gerasimova A. V.** (2004) Age determination of the White Sea bivalves
310 by the shell morphology. In Proceedings of the V Scientific Session of the Marine Biological
311 Station of St. Petersburg State University, St. Petersburg, Russia: 29–30. (in Russian)
- 312 **Mikkelsen N. and Pedersen T.** (2004) How can the stock recruitment relationship of the

- 313 Barents Sea capelin (*Mallotus villosus*) be improved by incorporating biotic and abiotic
314 factors? *Polar Research* 23(1), 19–26.
- 315 **Nazarova S. A., Genelt-Yanovsky E. A. and Maximovich N. V.** (2010) Linear growth of
316 *Macoma balthica* in the Murmansk tidal zone (the Barents Sea) *Vestnik Sankt-*
317 *Peterburgskogo Universiteta. Series 3: Biology* 4, 35–43. (In Russian)
- 318 **Nikula R., Strelkov P. and Väinölä R.** (2007) Diversity and trans-arctic invasion history of
319 mitochondrial lineages in the north atlantic *Macoma balthica* complex (Bivalvia:
320 Tellinidae). *Evolution* 61(4), 928–941.
- 321 **Oug E.** (2001) Polychaetes in intertidal rocky and sedimentary habitats in the region of Tromsø,
322 northern Norway. *Sarsia* 86(1), 75–83.
- 323 **Pfenninger M., Nowak C., and Magnin F.** (2007) Intraspecific range dynamics and niche
324 evolution in *Candidula* land snail species. *Biological Journal of the Linnean Society* 90(2),
325 303–317.
- 326 **Ramón M.** (2003) Population dynamics and secondary production of the cockle *Cerastoderma*
327 *edule* (L.) in a backbarrier tidal flat in the Wadden Sea. *Scientia Marina* 67(4), 429–443.
- 328 **R Core Team** (2014) *R: A language and environment for statistical computing*. R Foundation for
329 Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- 330 **Reise K., Herre E. and Sturm M.** (1994) Biomass and abundance of macrofauna in intertidal
331 sediments of Königshafen in the northern Wadden Sea. *Helgoländer Meeresuntersuchungen*
332 48(2-3), 201–215.
- 333 **Rivadeneira M. M., Hernáez P., Antonio Baeza J., Boltana S., Cifuentes M., Correa C., ...**
334 **and Thiel M.** (2010) Testing the abundant-centre hypothesis using intertidal porcelain crabs
335 along the Chilean coast: linking abundance and life-history variation. *Journal of*
336 *biogeography*, 37(3), 486–498.
- 337 **Sagarin R. D., and Gaines S. D.** (2002) The ‘abundant centre’ distribution: to what extent is it a
338 biogeographical rule? *Ecology letters* 5(1), 137–147.

- 339 **Sagarin R. D., Gaines S. D. and Gaylord B.** (2006) Moving beyond assumptions to understand
340 abundance distributions across the ranges of species. *Trends in Ecology and Evolution* 21(9),
341 524–530.
- 342 **Savari A., Sylvestre C., Sheader M., Le Gal Y. and Lockwood A.P.M.** (1989) Stress studies on
343 the common cockle (*Cerastoderma edule* L.) in Southampton Water. *Topics in Marine*
344 *biology. Scientia Marina* 53(2-3), 729-735.
- 345 **Semenova N.L.** (1974) The distribution of the bivalve *Macoma balthica* (L.) in some inlets of
346 Kandalaksha Bay of the White Sea. *Proceedings of the White Sea Biological Station of the*
347 *Moscow State University* 4, 87–102. (In Russian)
- 348 **Sexton J. P., McIntyre P. J., Angert A. L. and Rice K. J.** (2009) Evolution and ecology of
349 species range limits. *Annual Review of Ecology, Evolution, and Systematics* 40, 415–436
- 350 **Sneli J. A.** (1968) The intertidal distribution of polychaetes and molluscs on a muddy shore in
351 Nord-Møre, Norway. *Sarsia* 31(1), 63–68.
- 352 **Strasser M., Reinwald T. and Reise K.** (2001) Differential effects of the severe winter of
353 1995/96 on the intertidal bivalves *Mytilus edulis*, *Cerastoderma edule* and *Mya arenaria* in
354 the Northern Wadden Sea. *Helgoland Marine Research* 55(3), 190–197.
- 355 **Strelkov P., Nikula R. and Väinölä R.** (2007) *Macoma balthica* in the White and Barents Seas:
356 properties of a widespread marine hybrid swarm (Mollusca: Bivalvia). *Molecular ecology*
357 16(19), 4110–4127.
- 358 **Svenning J. C., Normand S. and Kageyama, M.** (2008) Glacial refugia of temperate trees in
359 Europe: insights from species distribution modelling. *Journal of Ecology* 96(6), 1117–1127.
- 360 **Tukey J. W.** (1976) *Exploratory data analysis*. Massachusetts: Addison-Wesley.
- 361 **Varfolomeeva M. and Naumov A.** (2013) Long-term temporal and spatial variation of
362 macrobenthos in the intertidal soft-bottom flats of two small bights (Chupa Inlet,
363 Kandalaksha Bay, White Sea). *Hydrobiologia* 706(1), 175–189.
- 364 **Zenkevich L.A.** (1963) *Biology of the Seas of the U.S.S.R.* Moscow. Translated from Russian to

365 English by S. Botsharskaya. New York: Interscience Publishers.

For Review Only

Table 1. Investigation sites and sampling details

Region	Site (code)	Latitude, longitude	Observation years
Western Murman	Pechenga (PG)	69.58, 31.27	2005
	Ura bay (UR)	68.99, 36.51	2010
Kola bay	Pala bay (PL)	69.19, 33.37	2006-2007
	Retinskoe (RT)	69.11, 33.38	2005
	Cape Abram (AB)	68.98, 33.03	2008
	Severnoe Nagornoe (NG)	68.90, 33.06	2005
	Tyuva (TU)	69.17, 33.63	2005
	Bolshaya Volokovaya (BV)	69.27, 33.62	2005
	Klimkovka (KL)	69.23, 34.64	2009
Eastern Murman	Dolgaya (DO)	69.17, 33.63	2009
	Gavrilovo (GV)	69.17, 35.86	2008
	Yarnyshnaya (YA)	69.09, 36.05	2004 - 2008
	Dalne-Zelenetskaya (DZ)	69.11, 36.10	2002 - 2009
	Shelpino (SH)	69.10, 36.21	2005, 2008
	Porchnikha (PR)	69.08, 36.25	2006 - 2007
	Tryashina (TR)	68.99, 36.51	2008
	Drozdovka (DR)	68.30, 38.44	2008
	Ivanovskaya subtidal (IV)	68.29, 38.71	2008
	Gremikha (GR)	68.03, 39.85	2009




Fig. 1. Distribution range map of *Macoma balthica* (European part) and *Cerastoderma edule* (A) and study area map (B). See table 1 for sample sites details.
175x85mm (300 x 300 DPI)




Fig. 2. Year-to-year variability of *Macoma balthica* (A) and *Cerastoderma edule* (B) mean abundance (lgN). Symbol colour corresponds to the region: Western Murman – solid grey, Kola Inlet – solid black and Eastern Murman – open symbols.

101x208mm (300 x 300 DPI)




Fig. 3. Mean abundance (N , ind.m⁻²) of *Macoma balthica* (A) and *Cerastoderma edule* (B) in the Barents sea. Samples are arranged from West to East. At the box-plots box marked Q1, median and Q3, whiskers marked 1.5 IQR. Mean abundance indicated at the top.
101x210mm (300 x 300 DPI)




Fig. 4. Size frequency distributions of *Macoma balthica* (A) into 1 mm classes and *Cerastoderma edule* (B) into 2 mm classes.
211x211mm (300 x 300 DPI)




Fig. 4. Size frequency distributions of *Macoma balthica* (A) into 1 mm classes and *Cerastoderma edule* (B) into 2 mm classes.
211x211mm (300 x 300 DPI)




Fig. 5. Geographic abundance distributions of *Macoma balthica* (A) and *Cerastoderma edule* (B). Circles squares are proportional to mean abundance. For sources of data see Appendix A.
109x84mm (300 x 300 DPI)




Fig. 5. Geographic abundance distributions of *Macoma balthica* (A) and *Cerastoderma edule* (B). Circles squares are proportional to mean abundance. For sources of data see Appendix A.
108x139mm (300 x 300 DPI)