

# Algebras in Tensor Triangular Categories

Seperability, Descent and Finite Étale Extensions

---

David Rubinstein

18th October, 2021

## Derived Category of Quasi-Coherent Sheaves

Let  $j : U \hookrightarrow V$  be a Zariski open Subscheme of a scheme  $V$ . It is a well known (well, to some!) theorem that the derived pullback functor  $j^* : D^{\text{qcoh}}(V) \rightarrow D^{\text{qcoh}}(U)$  is a smashing localization.

## Derived Category of Quasi-Coherent Sheaves

Let  $j : U \hookrightarrow V$  be a Zariski open Subscheme of a scheme  $V$ . It is a well known (well, to some!) theorem that the derived pullback functor  $j^* : D^{\text{qcoh}}(V) \rightarrow D^{\text{qcoh}}(U)$  is a smashing localization.

To spell that out: We can recover  $D^{\text{qcoh}}(U)$  as a localization of  $D^{\text{qcoh}}(V)$ ;

$$D^{\text{qcoh}}(V) / D_Z^{\text{qcoh}}(V) \cong D^{\text{qcoh}}(U)$$

where

## Derived Category of Quasi-Coherent Sheaves

Let  $j : U \hookrightarrow V$  be a Zariski open Subscheme of a scheme  $V$ . It is a well known (well, to some!) theorem that the derived pullback functor  $j^* : D^{\text{qcoh}}(V) \rightarrow D^{\text{qcoh}}(U)$  is a smashing localization.

To spell that out: We can recover  $D^{\text{qcoh}}(U)$  as a localization of  $D^{\text{qcoh}}(V)$ ;

$$D^{\text{qcoh}}(V) / D_Z^{\text{qcoh}}(V) \cong D^{\text{qcoh}}(U)$$

where

- $Z = V - U$  and  $D_Z^{\text{qcoh}}(V) = \ker(j^*)$  are the objects supported on  $Z$ ;
- the localization functor  $D^{\text{qcoh}}(V) \rightarrow D^{\text{qcoh}}(V) / D_Z^{\text{qcoh}}(V) \cong D^{\text{qcoh}}(U)$  has a right adjoint, which is automatically fully faithful.

## Derived Category of Quasi-Coherent Sheaves

Let  $j : U \hookrightarrow V$  be a Zariski open Subscheme of a scheme  $V$ . It is a well known (well, to some!) theorem that the derived pullback functor  $j^* : D^{\text{qcoh}}(V) \rightarrow D^{\text{qcoh}}(U)$  is a smashing localization.

To spell that out: We can recover  $D^{\text{qcoh}}(U)$  as a localization of  $D^{\text{qcoh}}(V)$ ;

$$D^{\text{qcoh}}(V) / D_Z^{\text{qcoh}}(V) \cong D^{\text{qcoh}}(U)$$

where

- $Z = V - U$  and  $D_Z^{\text{qcoh}}(V) = \ker(j^*)$  are the objects supported on  $Z$ ;
- the localization functor  $D^{\text{qcoh}}(V) \rightarrow D^{\text{qcoh}}(V) / D_Z^{\text{qcoh}}(V) \cong D^{\text{qcoh}}(U)$  has a right adjoint, which is automatically fully faithful.
- Since the right adjoint is fully faithful we can really view  $D^{\text{qcoh}}(U)$  as being a "piece" of  $D^{\text{qcoh}}(V)$

# Some Leading Questions

## How to Extend This?

The above construction gives some hope that it might be possible to do something analogous in other contexts. For example,

# Some Leading Questions

## How to Extend This?

The above construction gives some hope that it might be possible to do something analogous in other contexts. For example,

- can we do the same for any morphism of schemes: If  $f : V \rightarrow X$  is a morphism of schemes, is the derived pullback of  $f$  a localization? If not always, are there conditions we can place on it to make it so?

# Some Leading Questions

## How to Extend This?

The above construction gives some hope that it might be possible to do something analogous in other contexts. For example,

- can we do the same for any morphism of schemes: If  $f : V \rightarrow X$  is a morphism of schemes, is the derived pullback of  $f$  a localization? If not always, are there conditions we can place on it to make it so?
- In many other cases we have "inclusion" maps that induce maps of tensor triangular categories. Are these induced maps also localizations?



# Some Leading Questions

## How to Extend This?

The above construction gives some hope that it might be possible to do something analogous in other contexts. For example,

- can we do the same for any morphism of schemes: If  $f : V \rightarrow X$  is a morphism of schemes, is the derived pullback of  $f$  a localization? If not always, are there conditions we can place on it to make it so?
- In many other cases we have "inclusion" maps that induce maps of tensor triangular categories. Are these induced maps also localizations?
- For example: If  $H \hookrightarrow G$  is subgroup of a (finite) group  $G$ , is the restriction of scalars functor  $\text{Stab}(kG) \rightarrow \text{Stab}(kH)$  (or  $D(kG) \rightarrow D(kH)$ ) a localization?

## How do you multiply things???

Before we jump right into the world of tensor triangulated categories, let us start with a recap on plain old "tensor" categories; otherwise known as Monoidal Categories. These are morally the categories in which we can "multiply" objects in a coherent way.

## How do you multiply things???

Before we jump right into the world of tensor triangulated categories, let us start with a recap on plain old "tensor" categories; otherwise known as Monoidal Categories. These are morally the categories in which we can "multiply" objects in a coherent way.

- **Def (simplified):** A Monoidal Category  $(\mathcal{C}, \otimes, \mathbb{1})$  is a category  $\mathcal{C}$  equipped with a bifunctor  $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$  and a "unit" object  $\mathbb{1}$  satisfying a bunch of coherence axioms:

## How do you multiply things???

Before we jump right into the world of tensor triangulated categories, let us start with a recap on plain old "tensor" categories; otherwise known as Monoidal Categories. These are morally the categories in which we can "multiply" objects in a coherent way.

- **Def (simplified):** A Monoidal Category  $(\mathcal{C}, \otimes, \mathbb{1})$  is a category  $\mathcal{C}$  equipped with a bifunctor (which we will refer to as multiplication)  $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$  and a "unit" object  $\mathbb{1}$  satisfying a bunch of coherence axioms:
- Multiplication should be associative:  $(a \otimes b) \otimes c \cong a \otimes (b \otimes c)$

## How do you multiply things???

Before we jump right into the world of tensor triangulated categories, let us start with a recap on plain old "tensor" categories; otherwise known as Monoidal Categories. These are morally the categories in which we can "multiply" objects in a coherent way.

- **Def (simplified):** A Monoidal Category  $(\mathcal{C}, \otimes, \mathbb{1})$  is a category  $\mathcal{C}$  equipped with a bifunctor (which we will refer to as multiplication)  $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$  and a "unit" object  $\mathbb{1}$  satisfying a bunch of coherence axioms:
- Multiplication should be associative:  $(a \otimes b) \otimes c \cong a \otimes (b \otimes c)$
- Multiplication by the unit does nothing:  $\mathbb{1} \otimes a \cong a \cong a \otimes \mathbb{1}$

## Some Warnings

**Warnings:** This definition given above is extremely imprecise. For a more thorough definition of monoidal categories you can view the resources being shared. Let us just quickly comment a few things

## Some Warnings

**Warnings:** This definition given above is extremely imprecise. For a more thorough definition of monoidal categories you can view the resources being shared. Let us just quickly comment a few things:

- The associative isomorphisms above are really a given choice of natural isomorphisms
- There are two distinct maps in the unital isomorphisms (one for tensoring on the left and one for on the right)
- We have not made the claim yet that  $a \otimes b \cong b \otimes a$  yet.

# Symmetric Monoidal Categories

## Definition and Some Examples

**Def:** A **Symmetric Monoidal Category** is a monoidal category with a choice of braiding  $\tau : a \otimes b \cong b \otimes a$  that plays nice with the associativity isomorphisms and the two choices of unital isomorphisms.



## Definition and Some Examples

**Def:** A **Symmetric Monoidal Category** is a monoidal category with a choice of braiding  $\tau : a \otimes b \cong b \otimes a$  that plays nice with the associativity isomorphisms and the two choices of unital isomorphisms.

Let us look at some examples:

- $(\mathbf{Ab}, \otimes_{\mathbb{Z}}, \mathbb{Z})$
- More generally for  $R$  a commutative ring we have  $(R\text{-Mod}, \otimes_R, R)$

## Definition and Some Examples

**Def:** A **Symmetric Monoidal Category** is a monoidal category with a choice of braiding  $\tau : a \otimes b \cong b \otimes a$  that plays nice with the associativity isomorphisms and the two choices of unital isomorphisms.

Let us look at some examples:

- $(\mathbf{Ab}, \otimes_{\mathbb{Z}}, \mathbb{Z})$
- More generally for  $R$  a commutative ring we have  $(R\text{-Mod}, \otimes_R, R)$
- Let  $G$  be a finite group. Then  $(kG\text{-Mod}, \otimes_k, k)$
- More generally  $H$  be a Hopf Algebra over a field  $k$ . Then  $(H\text{-Mod}, \otimes_k, k)$

# Symmetric Monoidal Functors and their Adjoints

## (Symmetric) Monoidal Functors

A (lax) monoidal functor  $F : (\mathcal{C}, \otimes_{\mathcal{C}}, \mathbb{1}_{\mathcal{C}}) \rightarrow (\mathcal{D}, \otimes_{\mathcal{D}}, \mathbb{1}_{\mathcal{D}})$  is a functor equipped with a morphism

$$\varphi_0 : \mathbb{1}_{\mathcal{D}} \rightarrow F(\mathbb{1}_{\mathcal{C}})$$

and a natural transformation

$$\varphi_{a,b} : F(a) \otimes_{\mathcal{D}} F(b) \rightarrow F(a \otimes_{\mathcal{C}} b)$$

# Symmetric Monoidal Functors and their Adjoints

## (Symmetric) Monoidal Functors

A (lax) monoidal functor  $F : (\mathcal{C}, \otimes_{\mathcal{C}}, \mathbb{1}_{\mathcal{C}}) \rightarrow (\mathcal{D}, \otimes_{\mathcal{D}}, \mathbb{1}_{\mathcal{D}})$  is a functor equipped with a morphism

$$\varphi_0 : \mathbb{1}_{\mathcal{D}} \rightarrow F(\mathbb{1}_{\mathcal{C}})$$

and a natural transformation

$$\varphi_{a,b} : F(a) \otimes_{\mathcal{D}} F(b) \rightarrow F(a \otimes_{\mathcal{C}} b)$$

A strong monoidal functor is a lax monoidal functor where the two maps defined above are isomorphisms.

# Symmetric Monoidal Functors and their Adjoints

## (Symmetric) Monoidal Functors

A (lax) monoidal functor  $F : (\mathcal{C}, \otimes_{\mathcal{C}}, \mathbb{1}_{\mathcal{C}}) \rightarrow (\mathcal{D}, \otimes_{\mathcal{D}}, \mathbb{1}_{\mathcal{D}})$  is a functor equipped with a morphism

$$\varphi_0 : \mathbb{1}_{\mathcal{D}} \rightarrow F(\mathbb{1}_{\mathcal{C}})$$

and a natural transformation

$$\varphi_{a,b} : F(a) \otimes_{\mathcal{D}} F(b) \rightarrow F(a \otimes_{\mathcal{C}} b)$$

A strong monoidal functor is a lax monoidal functor where the two maps defined above are isomorphisms.

A (strong or lax) symmetric monoidal functor is a (strong or lax) monoidal functor such the following diagram commutes for all  $a, b$ :

# Symmetric Monoidal Functors and their Adjoints

## (Symmetric) Monoidal Functors

A (lax) monoidal functor  $F : (\mathcal{C}, \otimes_{\mathcal{C}}, \mathbb{1}_{\mathcal{C}}) \rightarrow (\mathcal{D}, \otimes_{\mathcal{D}}, \mathbb{1}_{\mathcal{D}})$  is a functor equipped with a morphism

$$\varphi_0 : \mathbb{1}_{\mathcal{D}} \rightarrow F(\mathbb{1}_{\mathcal{C}})$$

and a natural transformation

$$\varphi_{a,b} : F(a) \otimes_{\mathcal{D}} F(b) \rightarrow F(a \otimes_{\mathcal{C}} b)$$

A strong monoidal functor is a lax monoidal functor where the two maps defined above are isomorphisms.

A (strong or lax) symmetric monoidal functor is a (strong or lax) monoidal functor such the following diagram commutes for all  $a, b$ :

$$\begin{array}{ccc} F(a) \otimes F(b) & \xrightarrow{\varphi_{a,b}} & F(a \otimes b) \\ \tau_{F(a), F(b)} \downarrow & & \downarrow F(\tau_{a,b}) \\ F(b) \otimes F(a) & \xrightarrow{\varphi_{b,a}} & F(b \otimes a) \end{array}$$

# Symmetric Monoidal Functors and their Adjoints

## (Symmetric) Monoidal Adjoints

Let  $F : \mathcal{C} \rightarrow \mathcal{D}$  be a strong monoidal functor and suppose  $F$  has a right adjoint  $G$ .

**FACT:**  $G$  is a lax monoidal functor. Note that we therefore have the following two maps:

# Symmetric Monoidal Functors and their Adjoints

## (Symmetric) Monoidal Adjoints

Let  $F : \mathcal{C} \rightarrow \mathcal{D}$  be a strong monoidal functor and suppose  $F$  has a right adjoint  $G$ .

**FACT:**  $G$  is a lax monoidal functor. Note that we therefore have the following two maps:

$$\mathbb{1}_{\mathcal{C}} \rightarrow G(\mathbb{1}_{\mathcal{D}})$$

and

$$G(\mathbb{1}_{\mathcal{D}}) \otimes_{\mathcal{C}} G(\mathbb{1}_{\mathcal{D}}) \rightarrow G(\mathbb{1}_{\mathcal{D}} \otimes_{\mathcal{D}} \mathbb{1}_{\mathcal{D}}) \cong G(\mathbb{1}_{\mathcal{D}})$$



# Symmetric Monoidal Functors and their Adjoints

## (Symmetric) Monoidal Adjoints

Let  $F : \mathcal{C} \rightarrow \mathcal{D}$  be a strong monoidal functor and suppose  $F$  has a right adjoint  $G$ .

**FACT:**  $G$  is a lax monoidal functor. Note that we therefore have the following two maps:

$$\mathbb{1}_{\mathcal{C}} \rightarrow G(\mathbb{1}_{\mathcal{D}})$$

and

$$G(\mathbb{1}_{\mathcal{D}}) \otimes_{\mathcal{C}} G(\mathbb{1}_{\mathcal{D}}) \rightarrow G(\mathbb{1}_{\mathcal{D}} \otimes_{\mathcal{D}} \mathbb{1}_{\mathcal{D}}) \cong G(\mathbb{1}_{\mathcal{D}})$$

That is, there is a sort of "multiplication map" for  $G(\mathbb{1}_{\mathcal{D}})$ . Let us formalize that.

## Definition and Examples

Given a Symmetric Monoidal Category  $(\mathcal{C}, \otimes, \mathbb{1})$  we can talk about those objects  $A$  that admit a multiplication structure.

**Def:** A ring object in a Symmetric Monoidal Category is

## Definition

Given a Symmetric Monoidal Category  $(\mathcal{C}, \otimes, \mathbb{1})$  we can talk about those objects  $A$  that admit a multiplication structure.

**Def:** A ring object in a Symmetric Monoidal Category is an object  $A$  with maps:

- $\mu : A \otimes A \rightarrow A$  (called the multiplication map)
- $\eta : \mathbb{1} \rightarrow A$  (called the unit map)

such that the following diagrams commutes

# Ring Objects

## Definition

Given a Symmetric Monoidal Category  $(\mathcal{C}, \otimes, \mathbb{1})$  we can talk about those objects  $A$  that admit a multiplication structure.

**Def:** A ring object in a Symmetric Monoidal Category is an object  $A$  with maps:

- $\mu : A \otimes A \rightarrow A$  (called the multiplication map)
- $\eta : \mathbb{1} \rightarrow A$  (called the unit map)

such that the following diagrams commutes

$$\begin{array}{ccc} A \otimes A \otimes A & \xrightarrow{\mu \otimes 1} & A \otimes A \\ \downarrow 1 \otimes \mu & & \downarrow \mu \\ A \otimes A & \xrightarrow{\mu} & A \end{array}$$

$$\begin{array}{ccccc} A \otimes \mathbb{1} & \xrightarrow{1 \otimes \eta} & A \otimes A & \xleftarrow{\eta \otimes 1} & \mathbb{1} \otimes A \\ & \searrow & \downarrow \mu & \swarrow & \\ & & A & & \end{array}$$

# Ring Objects

## Definition

Given a Symmetric Monoidal Category  $(\mathcal{C}, \otimes, \mathbb{1})$  we can talk about those objects  $A$  that admit a multiplication structure.

**Def:** A ring object in a Symmetric Monoidal Category is an object  $A$  with maps:

- $\mu : A \otimes A \rightarrow A$  (called the multiplication map)
- $\eta : \mathbb{1} \rightarrow A$  (called the unit map)

such that the following diagrams commutes

$$\begin{array}{ccc} A \otimes A \otimes A & \xrightarrow{\mu \otimes 1} & A \otimes A \\ \downarrow 1 \otimes \mu & & \downarrow \mu \\ A \otimes A & \xrightarrow{\mu} & A \end{array}$$

$$\begin{array}{ccccc} A \otimes \mathbb{1} & \xrightarrow{1 \otimes \eta} & A \otimes A & \xleftarrow{\eta \otimes 1} & \mathbb{1} \otimes A \\ & \searrow & \downarrow \mu & \swarrow & \\ & & A & & \end{array}$$

We say the ring  $A$  is **commutative** if the multiplication map commutes with the braiding: that is if  $\mu \circ \tau = \mu$

# Examples of Ring Objects

## In the Categories we Previously Mentioned

This is nothing more than the categorified version of a ring. To see that, consider:

- Ring objects in  $(\mathbf{Ab}, \otimes_{\mathbb{Z}}, \mathbb{Z})$  are nothing more than...

# Examples of Ring Objects

## In the Categories we Previously Mentioned

This is nothing more than the categorified version of a ring. To see that, consider:

- Ring objects in  $(\mathbf{Ab}, \otimes_{\mathbb{Z}}, \mathbb{Z})$  are nothing more than Rings!
- More generally ring objects in  $(\mathbf{R} - \mathbf{Mod}, \otimes_{\mathbf{R}}, \mathbf{R})$  are....

# Examples of Ring Objects

## In the Categories we Previously Mentioned

This is nothing more than the categorified version of a ring. To see that, consider:

- Ring objects in  $(\mathbf{Ab}, \otimes_{\mathbb{Z}}, \mathbb{Z})$  are nothing more than Rings!
- More generally ring objects in  $(\mathbf{R-Mod}, \otimes_{\mathbf{R}}, \mathbf{R})$  are  $\mathbf{R}$ -algebras!
- For  $G$  finite Group, ring objects in  $(\mathbf{kG-Mod}, \otimes_{\mathbf{k}}, \mathbf{k})$  are  $\mathbf{k}$ -algebras with actions of  $G$  as algebra automorphisms.



# Examples of Ring Objects

## In the Categories we Previously Mentioned

This is nothing more than the categorified version of a ring. To see that, consider:

- Ring objects in  $(\mathbf{Ab}, \otimes_{\mathbb{Z}}, \mathbb{Z})$  are nothing more than Rings!
- More generally ring objects in  $(\mathbf{R} - \mathbf{Mod}, \otimes_{\mathbf{R}}, \mathbf{R})$  are  $\mathbf{R}$ -algebras!
- For  $G$  finite Group, ring objects in  $(\mathbf{k}G - \mathbf{Mod}, \otimes_{\mathbf{k}}, \mathbf{k})$  are  $\mathbf{k}$ -algebras with actions of  $G$  as algebra automorphisms.
- Recall a right adjoint  $G$  of any strong monoidal functor  $F$  is a lax monoidal functor. Then we saw that in fact  $G(\mathbb{1})$  is a ring object.

# Modules over Ring objects

Given me a ring and I'll give you a Module

Given a ring object we can talk about modules over the ring.

**Def:** Let  $A$  be a (commutative) ring object in a symmetric monoidal category  $(\mathcal{C}, \otimes, \mathbb{1})$ . A left  $A$ -module  $M$  is an object of  $\mathcal{C}$  equipped with a map  $\rho : A \otimes M \rightarrow M$  such that the following two diagrams commute:

# Modules over Ring objects

Given me a ring and I'll give you a Module

Given a ring object we can talk about modules over the ring.

**Def:** Let  $A$  be a (commutative) ring object in a symmetric monoidal category  $(\mathcal{C}, \otimes, \mathbb{1})$ . A left  $A$ -module  $M$  is an object of  $\mathcal{C}$  equipped with a map  $\rho : A \otimes M \rightarrow M$  such that the following two diagrams commute:

$$\begin{array}{ccc} A \otimes A \otimes M & \xrightarrow{1 \otimes \rho} & A \otimes M \\ \downarrow \mu \otimes 1 & & \downarrow \rho \\ A \otimes M & \xrightarrow{\rho} & M \end{array}$$

$$\begin{array}{ccc} \mathbb{1} \otimes M & \xrightarrow{\eta \otimes 1} & A \otimes M \\ & \searrow \iota_M & \downarrow \rho \\ & & M \end{array}$$

# Modules over Ring objects

Given me a ring and I'll give you a Module

Given a ring object we can talk about modules over the ring.

**Def:** Let  $A$  be a (commutative) ring object in a symmetric monoidal category  $(\mathcal{C}, \otimes, \mathbb{1})$ . A left  $A$ -module  $M$  is an object of  $\mathcal{C}$  equipped with a map  $\rho : A \otimes M \rightarrow M$  such that the following two diagrams commute:

$$\begin{array}{ccc} A \otimes A \otimes M & \xrightarrow{1 \otimes \rho} & A \otimes M \\ \mu \otimes 1 \downarrow & & \downarrow \rho \\ A \otimes M & \xrightarrow{\rho} & M \end{array}$$

$$\begin{array}{ccc} \mathbb{1} \otimes M & \xrightarrow{\eta \otimes 1} & A \otimes M \\ & \searrow \wr_M & \downarrow \rho \\ & & M \end{array}$$

Remark: These axioms are just souped up versions of the usual two axioms that  $a.(b.m) = (ab).m$  and  $1.m = m$  we are familiar with for Modules.

# Modules over Ring objects

## Category of Modules

Note that every object  $X \in \mathcal{C}$  gives rise to a "free A-module"  $A \otimes X$  whose structure map is given by the multiplication on  $A$ .

# Modules over Ring objects

## Category of Modules

Note that every object  $X \in \mathcal{C}$  gives rise to a "free  $A$ -module"  $A \otimes X$  whose structure map is given by the multiplication on  $A$ .

**Def:** The category of  $A$  modules is denoted  $A - \text{Mod}_{\mathcal{C}}$  and consists of

(Objects)  $A$ -modules

(Morphisms)  $A$ -linear maps. That is a map  $f : M \rightarrow N$  that commutes with the action of  $A$  on  $M$  and  $N$ .

# Modules over Ring objects

## Category of Modules

Note that every object  $X \in \mathcal{C}$  gives rise to a "free  $A$ -module"  $A \otimes X$  whose structure map is given by the multiplication on  $A$ .

**Def:** The category of  $A$  modules is denoted  $A - \text{Mod}_{\mathcal{C}}$  and consists of

(Objects)  $A$ -modules

(Morphisms)  $A$ -linear maps. That is a map  $f : M \rightarrow N$  that commutes with the action of  $A$  on  $M$  and  $N$ .

Note we then have an "extension of scalars functor"

$$F_A := A \otimes - : \mathcal{C} \rightarrow A - \text{Mod}_{\mathcal{C}}$$

which has a right adjoint

$$U_A : A - \text{Mod}_{\mathcal{C}} \rightarrow \mathcal{C}$$

that forgets the action of  $A$ .

# Modules over Ring objects

## Category of Modules

Note that every object  $X \in \mathcal{C}$  gives rise to a "free  $A$ -module"  $A \otimes X$  whose structure map is given by the multiplication on  $A$ .

**Def:** The category of  $A$  modules is denoted  $A - \text{Mod}_{\mathcal{C}}$  and consists of

(Objects)  $A$ -modules

(Morphisms)  $A$ -linear maps. That is a map  $f : M \rightarrow N$  that commutes with the action of  $A$  on  $M$  and  $N$ .

Note we then have an "extension of scalars functor"

$$F_A := A \otimes - : \mathcal{C} \rightarrow A - \text{Mod}_{\mathcal{C}}$$

which has a right adjoint

$$U_A : A - \text{Mod}_{\mathcal{C}} \rightarrow \mathcal{C}$$

that forgets the action of  $A$ .

**Remark:** We typically call the essential image of  $F$  the category of "Free Modules" and denote it by  $A - \text{Free}_{\mathcal{C}}$ . The adjunction above then of course restricts to:

$$\begin{array}{ccc} & \mathcal{C} & \\ F_A \nearrow & & \nwarrow U_A \\ A - \text{Free}_{\mathcal{C}} & \xrightarrow{\quad} & A - \text{Mod}_{\mathcal{C}} \end{array}$$



## Modules from an Adjunction

Let

$$\begin{array}{c} \mathcal{C} \\ \downarrow F \quad \uparrow G \\ \mathcal{D} \end{array}$$

be an adjoint pair between (Symmetric) Monoidal Categories. Recall that if  $F$  is strong Monoidal, then  $G$  is lax monoidal, turning  $A := G(\mathbb{1})$  into a ring object; so we can consider the category of  $A$ -Modules in  $\mathcal{C}$ .

# Realization of Ring Objects

## Modules from an Adjunction

Let

$$\begin{array}{c} \mathcal{C} \\ \uparrow F \\ \downarrow G \\ \mathcal{D} \end{array}$$

be an adjoint pair between (Symmetric) Monoidal Categories. Recall that if  $F$  is strong Monoidal, then  $G$  is lax monoidal, turning  $A := G(\mathbb{1})$  into a ring object; so we can consider the category of  $A$ -Modules in  $\mathcal{C}$ .

## Theorem:

There exist unique functors  $L$  and  $K$  making the following diagram commute:

$$\begin{array}{ccccc} & & \mathcal{C} & & \\ & \nearrow F_A & \uparrow F & \nwarrow U_A & \\ & \searrow U_A & \downarrow G & \nearrow F_A & \\ A - \text{Free}_{\mathcal{C}} & \xrightarrow{\quad L \quad} & \mathcal{D} & \xrightarrow{\quad K \quad} & A - \text{Mod}_{\mathcal{C}} \end{array}$$

## Tensor Triangulated Categories

**Def:** A **Tensor Triangulated Category** is a triangulated category  $\mathcal{T}$  equipped with a triangulated bi-functor  $- \otimes - : \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{T}$  and unit object that turns  $\mathcal{T}$  into a symmetric monoidal category. We will often call a tensor triangulated category a tt. category and will still denote it by  $(\mathcal{T}, \otimes, \mathbb{1})$

## Tensor Triangulated Categories

**Def:** A **Tensor Triangulated Category** is a triangulated category  $\mathcal{T}$  equipped with a triangulated bi-functor  $- \otimes - : \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{T}$  and unit object that turns  $\mathcal{T}$  into a symmetric monoidal category. We will often call a tensor triangulated category a tt. category and will still denote it by  $(\mathcal{T}, \otimes, \mathbb{1})$

Let us consider some examples (compare these to the examples of (symmetric) monoidal categories):

1.  $(D(R), \otimes_R^L, R)$ . The derived category of a commutative ring  $R$  with derived tensor product.

## Tensor Triangulated Categories

**Def:** A **Tensor Triangulated Category** is a triangulated category  $\mathcal{T}$  equipped with a triangulated bi-functor  $- \otimes - : \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{T}$  and unit object that turns  $\mathcal{T}$  into a symmetric monoidal category. We will often call a tensor triangulated category a tt. category and will still denote it by  $(\mathcal{T}, \otimes, \mathbb{1})$

Let us consider some examples (compare these to the examples of (symmetric) monoidal categories):

1.  $(D(R), \otimes_R^L, R)$ . The derived category of a commutative ring  $R$  with derived tensor product.
2.  $(\text{Stab}(kG), \otimes_k, k)$ . The "stable module" category for the group ring  $kG$ .

## Tensor Triangulated Categories

**Def:** A **Tensor Triangulated Category** is a triangulated category  $\mathcal{T}$  equipped with a triangulated bi-functor  $- \otimes - : \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{T}$  and unit object that turns  $\mathcal{T}$  into a symmetric monoidal category. We will often call a tensor triangulated category a tt. category and will still denote it by  $(\mathcal{T}, \otimes, \mathbb{1})$

Let us consider some examples (compare these to the examples of (symmetric) monoidal categories):

1.  $(D(R), \otimes_R^L, R)$ . The derived category of a commutative ring  $R$  with derived tensor product.
2.  $(\text{Stab}(kG), \otimes_k, k)$ . The "stable module" category for the group ring  $kG$ .
3.  $(SH, \wedge, S)$ . The "stable homotopy" category with smash product
4.  $(SH(G), \wedge, S)$ . The  $G$ -equivariant stable homotopy category for a finite group  $G$ .

## Tensor Triangulated Categories

**Def:** A **Tensor Triangulated Category** is a triangulated category  $\mathcal{T}$  equipped with a triangulated bi-functor  $- \otimes - : \mathcal{T} \times \mathcal{T} \rightarrow \mathcal{T}$  and unit object that turns  $\mathcal{T}$  into a symmetric monoidal category. We will often call a tensor triangulated category a tt. category and will still denote it by  $(\mathcal{T}, \otimes, \mathbb{1})$

Let us consider some examples (compare these to the examples of (symmetric) monoidal categories):

1.  $(D(R), \otimes_R^L, R)$ . The derived category of a commutative ring  $R$  with derived tensor product.
2.  $(\text{Stab}(kG), \otimes_k, k)$ . The "stable module" category for the group ring  $kG$ .
3.  $(SH, \wedge, S)$ . The "stable homotopy" category with smash product
4.  $(SH(G), \wedge, S)$ . The  $G$ -equivariant stable homotopy category for a finite group  $G$ .
5.  $(DM^{(\text{ét})}(S, R), \otimes, R)$ . The derived category of (étale) motives over base scheme  $S$  with coefficients in a commutative ring  $R$ .

## Some leading questions/examples

**Question:** Let  $R$  be a commutative ring and let  $A$  be an  $R$ -algebra: Is  $A[0]$  a ring object in  $D(R)$ ?



## Some leading questions/examples

**Question:** Let  $R$  be a commutative ring and let  $A$  be an  $R$ -algebra: Is  $A[0]$  a ring object in  $D(R)$ ?

**Answer:** Sadly, not always. This is because  $(A \otimes^L A)[0] \neq A[0] \otimes^L A[0]$  unless  $A$  is flat. So a flat  $R$ -Algebra  $A$  remains a ring object in  $D(R)$ .

## Some leading questions/examples

**Question 1:** Let  $R$  be a commutative ring and let  $A$  be an  $R$ -algebra: Is  $A[0]$  a ring object in  $D(R)$ ?

**Answer 1:** Sadly, not always. This is because  $(A \otimes^L A)[0] \neq A[0] \otimes^L A[0]$  unless  $A$  is flat. So a flat  $R$ -Algebra  $A$  remains a ring object in  $D(R)$ .

**Question 2:** Let  $A$  be a ring object in a tt category  $\mathcal{T}$ , and consider again the category of  $A$ -modules  $A - \text{Mod}_{\mathcal{T}}$ . Is  $A - \text{Mod}_{\mathcal{T}}$  triangulated?

## Some leading questions/examples

**Question 1:** Let  $R$  be a commutative ring and let  $A$  be an  $R$ -algebra: Is  $A[0]$  a ring object in  $D(R)$ ?

**Answer 1:** Sadly, not always. This is because  $(A \otimes^L A)[0] \neq A[0] \otimes^L A[0]$  unless  $A$  is flat. So a flat  $R$ -Algebra  $A$  remains a ring object in  $D(R)$ .

**Question 2:** Let  $A$  be a ring object in a tt category  $\mathcal{T}$ , and consider again the category of  $A$ -modules  $A - \text{Mod}_{\mathcal{T}}$ . Is  $A - \text{Mod}_{\mathcal{T}}$  triangulated?

**Answer 2:** Sadly... not always.

## Separability

**Def:** Let  $A$  be a ring object in a tt category  $\mathcal{T}$ . We say that  $A$  is **separable** if the multiplication map  $\mu : A \otimes A \rightarrow A$  admits a section as a two sided  $A$ -module.

## Separability

**Def:** Let  $A$  be a ring object in a tt category  $\mathcal{T}$ . We say that  $A$  is **separable** if the multiplication map  $\mu : A \otimes A \rightarrow A$  admits a section as a two sided  $A$ -module. That is, a morphism  $\sigma : A \rightarrow A \otimes A$  such that  $\mu\sigma = \text{id}_A$  and that the following diagram commutes:

## Separability

**Def:** Let  $A$  be a ring object in a tt category  $\mathcal{T}$ . We say that  $A$  is **separable** if the multiplication map  $\mu : A \otimes A \rightarrow A$  admits a section as a two sided  $A$ -module. That is, a morphism  $\sigma : A \rightarrow A \otimes A$  such that  $\mu\sigma = \text{id}_A$  and that the following diagram commutes:

$$\begin{array}{ccccc} & & A \otimes A & & \\ & \swarrow \sigma \otimes 1 & \downarrow \mu & \searrow 1 \otimes \sigma & \\ A \otimes A \otimes A & & A & & A \otimes A \otimes A \\ & \searrow 1 \otimes \mu & \downarrow \sigma & \swarrow \mu \otimes 1 & \\ & & A \otimes A & & \end{array}$$

# Separable Rings

## Separability

**Def:** Let  $A$  be a ring object in a tt category  $\mathcal{T}$ . We say that  $A$  is **separable** if the multiplication map  $\mu : A \otimes A \rightarrow A$  admits a section as a two sided  $A$ -module. That is, a morphism  $\sigma : A \rightarrow A \otimes A$  such that  $\mu\sigma = \text{id}_A$  and that the following diagram commutes:

$$\begin{array}{ccccc} & & A \otimes A & & \\ \sigma \otimes 1 \swarrow & & \downarrow \mu & \searrow 1 \otimes \sigma & \\ A \otimes A \otimes A & & A & & A \otimes A \otimes A \\ \downarrow 1 \otimes \mu & & \downarrow \sigma & & \downarrow \mu \otimes 1 \\ & & A \otimes A & & \end{array}$$

## Theorem:

Let  $A$  be a separable ring in a tt category  $\mathcal{T}$ . Then the category  $A - \text{Mod}_{\mathcal{T}}$  is canonically triangulated such that the extension of scalars functor

$$F_A : \mathcal{T} \rightarrow A - \text{Mod}_{\mathcal{T}}$$

is a tt functor.

## Smashing Localizations

Before diving into some concrete examples, let us note a particular case of this theory. Let  $\mathcal{T}$  be a tt category and consider a smashing localization  $L : \mathcal{T} \rightarrow \mathcal{T}$



## Smashing Localizations

Before diving into some concrete examples, let us note a particular case of this theory. Let  $\mathcal{T}$  be a tt category and consider a smashing localization  $L : \mathcal{T} \rightarrow \mathcal{T}$ . Recall: A smashing localization is a localization  $L$  such that

1.  $L$  admits a fully faithful right adjoint (allowing us to think of the localization as a subcategory: the essential image of "L-local objects")

## Smashing Localizations

Before diving into some concrete examples, let us note a particular case of this theory. Let  $\mathcal{T}$  be a tt category and consider a smashing localization  $L : \mathcal{T} \rightarrow \mathcal{T}$ . Recall: A smashing localization is a localization  $L$  such that

1.  $L$  admits a fully faithful right adjoint (allowing us to think of the localization as a subcategory: the essential image of "L-local objects")
2. For any object  $t \in \mathcal{T}$ , we have a canonical isomorphism  $L(t) \cong L(1) \otimes t$

## Smashing Localizations

Before diving into some concrete examples, let us note a particular case of this theory. Let  $\mathcal{T}$  be a tt category and consider a smashing localization  $L : \mathcal{T} \rightarrow \mathcal{T}$ . Recall: A smashing localization is a localization  $L$  such that

1.  $L$  admits a fully faithful right adjoint (allowing us to think of the localization as a subcategory: the essential image of "L-local objects")
2. For any object  $t \in \mathcal{T}$ , we have a canonical isomorphism  $L(t) \cong L(\mathbb{1}) \otimes t$

Then let us let  $A := L(\mathbb{1})$ . One then shows that  $A \otimes A \cong A$ ; this allows us to view  $A$  as a separable algebra (by taking  $\sigma$  to be the inverse of the isomorphism.)

## Smashing Localizations

Before diving into some concrete examples, let us note a particular case of this theory. Let  $\mathcal{T}$  be a tt category and consider a smashing localization  $L : \mathcal{T} \rightarrow \mathcal{T}$ . Recall: A smashing localization is a localization  $L$  such that

1.  $L$  admits a fully faithful right adjoint (allowing us to think of the localization as a subcategory: the essential image of "L-local objects")
2. For any object  $t \in \mathcal{T}$ , we have a canonical isomorphism  $L(t) \cong L(\mathbb{1}) \otimes t$

Then let us let  $A := L(\mathbb{1})$ . One then shows that  $A \otimes A \cong A$ ; this allows us to view  $A$  as a separable algebra (by taking  $\sigma$  to be the inverse of the isomorphism.)

In this case then, the category  $A - \text{Mod}_{\mathcal{T}}$  is just the category of L-local objects.

# Examples: Take 1

## Smashing Localizations

Before diving into some concrete examples, let us note a particular case of this theory. Let  $\mathcal{T}$  be a tt category and consider a smashing localization  $L : \mathcal{T} \rightarrow \mathcal{T}$ . Recall: A smashing localization is a localization  $L$  such that

1.  $L$  admits a fully faithful right adjoint (allowing us to think of the localization as a subcategory: the essential image of "L-local objects")
2. For any object  $t \in \mathcal{T}$ , we have a canonical isomorphism  $L(t) \cong L(\mathbb{1}) \otimes t$

Then let us let  $A := L(\mathbb{1})$ . One then shows that  $A \otimes A \cong A$ ; this allows us to view  $A$  as a separable algebra (by taking  $\sigma$  to be the inverse of the isomorphism).

In this case then, the category  $A - \text{Mod}_{\mathcal{T}}$  is just the category of L-local objects.

## Back to OG example

**Remark:** The case of the open immersion of schemes  $U \hookrightarrow V$  can thus be stated as follows: Letting  $A = j_*(\mathcal{O}_U)$  we have that  $A\text{-Mod} \cong j_*j^*\text{-Local objects} \cong D^{\text{qcoh}}(U)$ .

That is, we can view  $D^{\text{qcoh}}(U)$  as being a sort of Module category **inside**  $D^{\text{qcoh}}(V)$

## The Main Construction

Let  $\mathcal{C} \begin{smallmatrix} \xleftarrow{G} \\ \xrightarrow{F} \end{smallmatrix} \mathcal{D}$  be an adjunction of tt categories.

## The Main Construction

Let  $\mathcal{C} \xrightleftharpoons[F]{G} \mathcal{D}$  be an adjunction of tt categories. Then recall that  $A := G(\mathbb{1}_{\mathcal{D}})$  is a ring object and that we had the following picture

$$\begin{array}{ccccc}
 & & \mathcal{C} & & \\
 & \nearrow F_A & \uparrow F & \nwarrow U_A & \\
 A - \text{Free}_{\mathcal{C}} & \xrightarrow{\quad L \quad} & \mathcal{D} & \xrightarrow{\quad K \quad} & A - \text{Mod}_{\mathcal{C}} \\
 & \nwarrow U_A & \downarrow G & \nearrow F_A & \\
 & & & & 
 \end{array}$$

# Finite Étale Extensions

## The Main Construction

Let  $\mathcal{C} \xrightleftharpoons[F]{G} \mathcal{D}$  be an adjunction of tt categories. Then recall that  $A := G(\mathbb{1}_{\mathcal{D}})$  is a ring object and that we had the following picture

$$\begin{array}{ccccc}
 & & \mathcal{C} & & \\
 & \swarrow F_A & \uparrow F & \nwarrow U_A & \\
 A - \text{Free}_{\mathcal{C}} & \xrightarrow{\quad L \quad} & \mathcal{D} & \xrightarrow{\quad K \quad} & A - \text{Mod}_{\mathcal{C}} \\
 & \nwarrow U_A & \downarrow G & \swarrow F_A & \\
 & & \mathcal{D} & & 
 \end{array}$$

## Main Definition:

Let  $F$  and  $A$  be as above. We say  $F$  is a **finite étale extension** if  $A$  is a (compact) separable ring object such that

- the functor  $\mathcal{D} \xrightarrow{K} A - \text{Mod}_{\mathcal{C}}$  is an tt equivalence of tt categories
- under which the functor  $F$  becomes isomorphic to the extension of scalars functor  $F_A$  and
- $G$  becomes isomorphic to the forgetful functor  $U_A$



## The Reason for the Name

Recall that if  $A$  is a flat  $R$ -Algebra then  $A[0]$  remains a ring object in  $D(R)$ .

**Def:** An **étale  $R$ -Algebra  $S$**  is a separable, flat  $R$ -algebra of finite presentation.

## The Reason for the Name

Recall that if  $A$  is a flat  $R$ -Algebra then  $A[0]$  remains a ring object in  $D(R)$ .

**Def:** An **étale  $R$ -Algebra  $S$**  is a separable, flat  $R$ -algebra of finite presentation.

**Thrm 1:** Let  $S$  be an étale  $R$ -algebra and consider the extension of scalars functor:

$$D(R) \xrightarrow{F := S[0] \otimes_R^L -} D(S)$$

Then  $F$  is a finite étale extension. That is the category  $D(S)$  is canonically equivalent to the category of  $S$ -Modules inside  $D(R)$ .

## The Reason for the Name

Recall that if  $A$  is a flat  $R$ -Algebra then  $A[0]$  remains a ring object in  $D(R)$ .

**Def:** An **étale  $R$ -Algebra**  $S$  is a separable, flat  $R$ -algebra of finite presentation.

**Thrm 1:** Let  $S$  be an étale  $R$ -algebra and consider the extension of scalars functor:

$$D(R) \xrightarrow{F := S[0] \otimes_R^L -} D(S)$$

Then  $F$  is a finite étale extension. That is the category  $D(S)$  is canonically equivalent to the category of  $S$ -Modules inside  $D(R)$ .

## Thrm 2

**Thrm 2:** Let  $f : V \rightarrow X$  be a separated étale morphism of quasi-compact, quasi-separated schemes. Then the functor

$$f^* : D^{\text{qcoh}}(X) \rightarrow D^{\text{qcoh}}(V)$$

is a finite étale extension. That is, we have an equivalence of categories

$$D^{\text{qcoh}}(V) \cong \text{Rf}_*(\mathbb{1}) - \text{Mod}_{D^{\text{qcoh}}(X)}$$

## Modular Representation Theory

Let  $G$  be a finite group and consider the tt category  $\mathrm{Stab}(kG)$ . Let  $H \leq G$  be a subgroup and recall that we get the following adjunction:

## Modular Representation Theory

Let  $G$  be a finite group and consider the tt category  $\mathrm{Stab}(kG)$ . Let  $H \leq G$  be a subgroup and recall that we get the following adjunction:

$$\begin{array}{ccc} & \mathrm{Stab}(kG) & \\ \mathrm{Res}_H^G \downarrow & \updownarrow & \mathrm{Ind}_H^G := kG \otimes - \\ & \mathrm{Stab}(kH) & \end{array}$$

## Modular Representation Theory

Let  $G$  be a finite group and consider the tt category  $\text{Stab}(kG)$ . Let  $H \leq G$  be a subgroup and recall that we get the following adjunction:

$$\begin{array}{ccc} & \text{Stab}(kG) & \\ \text{Res}_H^G \updownarrow & & \updownarrow \text{Ind}_H^G := kG \otimes - \\ & \text{Stab}(kH) & \end{array}$$

**Thrm:** Let  $A_H^G := \text{Ind}_H^G(\mathbb{1}) \cong k(G/H)$ . The Restriction to a subgroup functor is a finite étale extension. That is, the category  $\text{Stab}(kH)$  is canonically isomorphic to the category of  $A$ -Modules in  $\text{Stab}(kG)$  under which the restriction functor is isomorphic to the extension of scalars functor  $F_A$ .

## Modular Representation Theory

Let  $G$  be a finite group and consider the  $\mathbf{tt}$  category  $\mathrm{Stab}(kG)$ . Let  $H \leq G$  be a subgroup and recall that we get the following adjunction:

$$\begin{array}{ccc} \mathrm{Stab}(kG) & & \\ \mathrm{Res}_H^G \updownarrow & \mathrm{Ind}_H^G := kG \otimes - & \\ \mathrm{Stab}(kH) & & \end{array}$$

**Thrm:** Let  $A_H^G := \mathrm{Ind}_H^G(\mathbb{1}) \cong k(G/H)$ . The Restriction to a subgroup functor is a finite étale extension. That is, the category  $\mathrm{Stab}(kH)$  is canonically isomorphic to the category of  $A$ -Modules in  $\mathrm{Stab}(kG)$  under which the restriction functor is isomorphic to the extension of scalars functor  $F_A$ .

**Remark:** One can then phrase questions about extending representations of  $H$  to  $G$  in terms of "descent" of the ring  $A_H^G$ . I will not mention much more about this, but will leave some references for you to look at. The big takeaway is that this ring  $A_H^G$  satisfies descent iff  $[G : H]$  is invertible in  $k$ .

## Equivariant Homotopy Theory

Let  $G$  be a compact Lie Group (ex; a finite group) and consider the tt category  $\mathrm{SH}(G)$ . Let  $H \leq G$  be a closed subgroup- we get the following adjunction:



## Equivariant Homotopy Theory

Let  $G$  be a compact Lie Group (ex; a finite group) and consider the tt category  $\mathrm{SH}(G)$ . Let  $H \leq G$  be a closed subgroup- we get the following adjunction:

$$\begin{array}{ccccc}
 & & \mathrm{SH}(G) & & \\
 & \uparrow & | & \uparrow & \\
 \mathrm{Ind}_H^G := G_+ \wedge_H - & \mathrm{Res}_H^G & & \mathrm{CoInd}_H^G := F_H(G_+, -) & \\
 & \downarrow & | & \downarrow & \\
 & & \mathrm{SH}(H) & & 
 \end{array}$$

## Equivariant Homotopy Theory

Let  $G$  be a compact Lie Group (ex; a finite group) and consider the tt category  $SH(G)$ . Let  $H \leq G$  be a closed subgroup- we get the following adjunction:

$$\begin{array}{ccccc}
 & & SH(G) & & \\
 & \uparrow & | & \uparrow & \\
 Ind_H^G := G + \wedge_H - & \text{Res}_H^G & & CoInd_H^G := F_H(G +, -) & \\
 & \downarrow & | & \downarrow & \\
 & & SH(H) & & 
 \end{array}$$

Note that when  $[G : H] < \infty$  there is an isomorphism of functors between  $Ind_H^G \cong CoInd_H^G$ .

# Étale Extensions in Equivariant contexts

## Equivariant Homotopy Theory

Let  $G$  be a compact Lie Group (ex; a finite group) and consider the tt category  $SH(G)$ . Let  $H \leq G$  be a closed subgroup- we get the following adjunction:

$$\begin{array}{ccccc}
 & & SH(G) & & \\
 & \uparrow & | & \uparrow & \\
 Ind_H^G := G_+ \wedge_H - & \text{Res}_H^G & & CoInd_H^G := F_H(G_+, -) & \\
 & \downarrow & | & \downarrow & \\
 & & SH(H) & & 
 \end{array}$$

Note that when  $[G : H] < \infty$  there is an isomorphism of functors between  $Ind_H^G \cong CoInd_H^G$ .

## Theorem:

Let  $H \leq G$  be a closed subgroup of finite index. Let  $A := F_H(G_+, \mathbb{1}_{SH(H)}) \cong G_+ \wedge_H \mathbb{1}_{SH(H)} \cong \sum^{\infty} (G/H)_+$ . Then restriction to  $H$  is a finite étale extension; that is the category of  $A$ -Modules in  $SH(G)$  is equivalent to  $SH(H)$ .

## Some Topics to Read if Interested

There are many directions one can take with this:

- Read about what the extension of scalars functor does on Spectra
- Classify all separable algebras in a given tt category
- Read about descent for separable algebras
- See how far you can push the analogy of a ring: going up theorem, "residue fields", Galois extensions, etc
- Reading about the behavior of finite étale morphisms on the "big" categories