

(Tensor) Triangulated Categories

What are they good for- part 2?

David Rubinstein

An analogue of Ring Theory

1. Last talk we discussed the axioms and some elementary examples and properties of Triangulated Categories. We saw that while they were not Abelian categories in general, they behaved rather similarly.

An analogue of Ring Theory

1. Last talk we discussed the axioms and some elementary examples and properties of Triangulated Categories. We saw that while they were not Abelian categories in general, they behaved rather similarly.
2. Moreover, we saw that we could get pretty far treating them as some sort of strange Group.

An analogue of Ring Theory

1. Last talk we discussed the axioms and some elementary examples and properties of Triangulated Categories. We saw that while they were not Abelian categories in general, they behaved rather similarly.
2. Moreover, we saw that we could get pretty far treating them as some sort of strange Group.
 - 2.1 We treated thick subcategories as normal subgroups, Triangulated Functors as Group Homomorphisms, and Verdier Localization as the natural projection map.

An analogue of Ring Theory

1. Last talk we discussed the axioms and some elementary examples and properties of Triangulated Categories. We saw that while they were not Abelian categories in general, they behaved rather similarly.
2. Moreover, we saw that we could get pretty far treating them as some sort of strange Group.
 - 2.1 We treated thick subcategories as normal subgroups, Triangulated Functors as Group Homomorphisms, and Verdier Localization as the natural projection map.
3. Today we will continue this analogy, and will now talk about what the Triangulated Analogue of a Ring should be.
 - 3.1 We call such "rings" Tensor-Triangulated Categories

An analogue of Ring Theory

1. Last talk we discussed the axioms and some elementary examples and properties of Triangulated Categories. We saw that while they were not Abelian categories in general, they behaved rather similarly.
2. Moreover, we saw that we could get pretty far treating them as some sort of strange Group.
 - 2.1 We treated thick subcategories as normal subgroups, Triangulated Functors as Group Homomorphisms, and Verdier Localization as the natural projection map.
3. Today we will continue this analogy, and will now talk about what the Triangulated Analogue of a Ring should be.
 - 3.1 We call such "rings" Tensor-Triangulated Categories
 - 3.2 With this product structure in hand (to be defined soon), we can make progress on some promised Unification we hinted at last week. Let us recall those leading questions now.

Some Leading Questions

Unification

1. Since so many examples of these Triangulated Categories exist ranging from algebra/geometry (examples 2,3,4) to topology (1,5,6) to analysis (7) the vague hope is that by studying "Triangulated Categories" writ large, one can learn things about all these topics in one fell swoop.
2. For example- Can we relate the ideas of line bundles in Algebraic-Geometry and Endotrivial Modules in Representation Theory? Moreover, can we relate $\text{Spec}(R)$ for a commutative ring with Support Varieties $\mathcal{V}_G := \text{Proj}(H^\bullet(G, k))$
3. There is a famous nilpotence theorem of Hopkins and Smith in Algebraic Topology- are there analogous nilpotent theorems in other contexts? In Alg Topology this nilpotence theorem provides a stratification for our category, can we expect the same for other nilpotence theorems?

The Axioms, Take 2

Bring on the Tensor

Throughout the next slides, we will let \mathcal{K} be an essentially small triangulated category. (That is, there is only a Set of Isomorphism Classes of Objects).

The Axioms, Take 2

Bring on the Tensor

Throughout the next slides, we will let \mathcal{K} be an essentially small triangulated category. (That is, there is only a Set of Isomorphism Classes of Objects).

1. We say \mathcal{K} is a Tensor Triangulated Category (tt category), written as $(\mathcal{K}, \otimes, 1)$ if there is a functor $- \otimes - : \mathcal{K} \times \mathcal{K} \rightarrow \mathcal{K}$ satisfying a bunch of axioms (see below) and such that $- \otimes -$ is a triangulated functor in each variable.

The Axioms, Take 2

Bring on the Tensor

Throughout the next slides, we will let \mathcal{K} be an essentially small triangulated category. (That is, there is only a Set of Isomorphism Classes of Objects).

1. We say \mathcal{K} is a Tensor Triangulated Category (tt category), written as $(\mathcal{K}, \otimes, 1)$ if there is a functor $- \otimes - : \mathcal{K} \times \mathcal{K} \rightarrow \mathcal{K}$ satisfying a bunch of axioms (see below) and such that $- \otimes -$ is a triangulated functor in each variable.

1.1 We have the following natural isomorphisms:

$$l_a : 1 \otimes a \cong a$$

$$r_a : a \otimes 1 \cong a$$

$$\tau_{a,b} : a \otimes b \cong b \otimes a$$

along with associativity axioms in such a way that everything behaves nice with one another.

The Axioms, Take 2

Bring on the Tensor

Throughout the next slides, we will let \mathcal{K} be an essentially small triangulated category. (That is, there is only a Set of Isomorphism Classes of Objects).

1. We say \mathcal{K} is a Tensor Triangulated Category (tt category), written as $(\mathcal{K}, \otimes, 1)$ if there is a functor $- \otimes - : \mathcal{K} \times \mathcal{K} \rightarrow \mathcal{K}$ satisfying a bunch of axioms (see below) and such that $- \otimes -$ is a triangulated functor in each variable.

1.1 We have the following natural isomorphisms:

$$l_a : 1 \otimes a \cong a$$

$$r_a : a \otimes 1 \cong a$$

$$\tau_{a,b} : a \otimes b \cong b \otimes a$$

along with associativity axioms in such a way that everything behaves nice with one another.

2. We think of \otimes as a commutative product and 1 as the unit. Indeed, in the literature, you will often see the above referred to as a "Symmetric Monoidal Category" (although this can refer to any tensor category, not necessarily triangulated), or an "Axiomatic Stable Homotopy Theory"

All assumptions on the table

Duelizable Objects

Before we can investigate what the tensor gives us, we need to give one more definition/assumption.

Dualizable Objects

Before we can investigate what the tensor gives us, we need to give one more definition/assumption.

1. Let $(\mathcal{K}, \otimes, 1)$ be a tt category. Then we say \mathcal{K} is a closed symmetric monoidal category if, for each $k \in \mathcal{K}$ the functor $k \otimes -$ has a (triangulated) right adjoint, denoted $\text{hom}(k, -)$. These can be assembled into a (triangulated) bifunctor $\text{hom}(-, -)$

Dualizable Objects

Before we can investigate what the tensor gives us, we need to give one more definition/assumption.

1. Let $(\mathcal{K}, \otimes, 1)$ be a tt category. Then we say \mathcal{K} is a closed symmetric monoidal category if, for each $k \in \mathcal{K}$ the functor $k \otimes -$ has a (triangulated) right adjoint, denoted $\text{hom}(k, -)$. These can be assembled into a (triangulated) bifunctor $\text{hom}(-, -)$
2. Then for each $k \in \mathcal{K}$ we denote the dual of k as $k^\vee := \text{hom}(k, 1)$. There is a natural map $k^\vee \otimes l \rightarrow \text{hom}(k, l)$ given from the counit of the adjunction.

All assumptions on the table

Duelizable Objects

Before we can investigate what the tensor gives us, we need to give one more definition/assumption.

1. Let $(\mathcal{K}, \otimes, 1)$ be a tt category. Then we say \mathcal{K} is a closed symmetric monoidal category if, for each $k \in \mathcal{K}$ the functor $k \otimes -$ has a (triangulated) right adjoint, denoted $\text{hom}(k, -)$. These can be assembled into a (triangulated) bifunctor $\text{hom}(-, -)$
2. Then for each $k \in \mathcal{K}$ we denote the dual of k as $k^\vee := \text{hom}(k, 1)$. There is a natural map $k^\vee \otimes l \rightarrow \text{hom}(k, l)$ given from the counit of the adjunction.
3. We say that \mathcal{K} is rigid, if this natural map $k^\vee \otimes l \xrightarrow{\sim} \text{hom}(k, l)$ is an isomorphism for all k, l . (sometimes this condition is called "Strongly Duelizable")

All assumptions on the table

Dualizable Objects

Before we can investigate what the tensor gives us, we need to give one more definition/assumption.

1. Let $(\mathcal{K}, \otimes, 1)$ be a tt category. Then we say \mathcal{K} is a closed symmetric monoidal category if, for each $k \in \mathcal{K}$ the functor $k \otimes -$ has a (triangulated) right adjoint, denoted $\text{hom}(k, -)$. These can be assembled into a (triangulated) bifunctor $\text{hom}(-, -)$
2. Then for each $k \in \mathcal{K}$ we denote the dual of k as $k^\vee := \text{hom}(k, 1)$. There is a natural map $k^\vee \otimes l \rightarrow \text{hom}(k, l)$ given from the counit of the adjunction.
3. We say that \mathcal{K} is rigid, if this natural map $k^\vee \otimes l \xrightarrow{\sim} \text{hom}(k, l)$ is an isomorphism for all k, l . (sometimes this condition is called "Strongly Dualizable")
4. Rigidity is an assumption that just simplifies our life a lot- for example, the tensor-hom adjunctions imply the k is a direct summand of $k \otimes k \otimes k^\vee$ and that k^\vee is a direct summand of $k \otimes k^\vee \otimes k^\vee$. We will note in a few slides why this is useful.

Examples of tt Categories

Bring on the Examples

The following are examples of tt categories. We shall write them all as $(\mathcal{K}, \otimes, 1)$

1. Let k be a field of char p dividing the order of the group G . Then $(\text{stab}(kG), \otimes_k, k)$ is a tt category, where the tensor is the usual tensor with diagonal action of g
 $g(M \otimes_k N) = gM \otimes_k gN$

Examples of tt Categories

Bring on the Examples

The following are examples of tt categories. We shall write them all as $(\mathcal{K}, \otimes, 1)$

1. Let k be a field of char p dividing the order of the group G . Then $(\text{stab}(kG), \otimes_k, k)$ is a tt category, where the tensor is the usual tensor with diagonal action of g
 $g(M \otimes_k N) = gM \otimes_k gN$
2. Let R be a commutative ring. Then $(D^b(R), \otimes^L, R)$ is a tt category where \otimes^L is the Left derived tensor product.

Examples of tt Categories

Bring on the Examples

The following are examples of tt categories. We shall write them all as $(\mathcal{K}, \otimes, 1)$

1. Let k be a field of char p dividing the order of the group G . Then $(\text{stab}(kG), \otimes_k, k)$ is a tt category, where the tensor is the usual tensor with diagonal action of g
 $g(M \otimes_k N) = gM \otimes_k gN$
2. Let R be a commutative ring. Then $(D^b(R), \otimes^L, R)$ is a tt category where \otimes^L is the Left derived tensor product.
3. Again, let R be a commutative ring. Then $(D^{\text{perf}}(R), \otimes^L, R)$ is a tt category (where $D^{\text{perf}}(R)$ is the derived category of perfect complexes)

Examples of tt Categories

Bring on the Examples

The following are examples of tt categories. We shall write them all as $(\mathcal{K}, \otimes, 1)$

1. Let k be a field of char p dividing the order of the group G . Then $(\text{stab}(kG), \otimes_k, k)$ is a tt category, where the tensor is the usual tensor with diagonal action of g
 $g(M \otimes_k N) = gM \otimes_k gN$
2. Let R be a commutative ring. Then $(D^b(R), \otimes^L, R)$ is a tt category where \otimes^L is the Left derived tensor product.
3. Again, let R be a commutative ring. Then $(D^{\text{perf}}(R), \otimes^L, R)$ is a tt category (where $D^{\text{perf}}(R)$ is the derived category of perfect complexes)
4. $(SH^{\text{finite}}, \wedge, \mathbb{S}^0)$ or more generally the G -Equivariant Stable Homotopy Category for G a Group $(SH^{\text{finite}}(G), \wedge, \mathbb{S}^0)$

Examples of tt Categories

Bring on the Examples

The following are examples of tt categories. We shall write them all as $(\mathcal{K}, \otimes, 1)$

1. Let k be a field of char p dividing the order of the group G . Then $(\text{stab}(kG), \otimes_k, k)$ is a tt category, where the tensor is the usual tensor with diagonal action of g
 $g(M \otimes_k N) = gM \otimes_k gN$
2. Let R be a commutative ring. Then $(D^b(R), \otimes^L, R)$ is a tt category where \otimes^L is the Left derived tensor product.
3. Again, let R be a commutative ring. Then $(D^{\text{perf}}(R), \otimes^L, R)$ is a tt category (where $D^{\text{perf}}(R)$ is the derived category of perfect complexes)
4. $(SH^{\text{finite}}, \wedge, \mathbb{S}^0)$ or more generally the G -Equivariant Stable Homotopy Category for G a Group $(SH^{\text{finite}}(G), \wedge, \mathbb{S}^0)$
5. Variations of the theme for points 2 and 3 for X a quasi compact, quasi separated scheme.

What's the Big Ideal?

Until further notice, let $(\mathcal{K}, \otimes, 1)$ be a closed, rigid tt category. Then, continuing our analogy of treating \mathcal{K} as a ring, we introduce the following definitions:

What's the Big Ideal?

Until further notice, let $(\mathcal{K}, \otimes, 1)$ be a closed, rigid tt category. Then, continuing our analogy of treating \mathcal{K} as a ring, we introduce the following definitions:

1. We say $\mathcal{I} \subseteq \mathcal{K}$ is a tensor-ideal, if \mathcal{I} is a thick subcategory of \mathcal{K} such that $\mathcal{K} \otimes \mathcal{I} \subseteq \mathcal{I}$

What's the Big Ideal?

Until further notice, let $(\mathcal{K}, \otimes, 1)$ be a closed, rigid tt category. Then, continuing our analogy of treating \mathcal{K} as a ring, we introduce the following definitions:

1. We say $\mathcal{I} \subseteq \mathcal{K}$ is a tensor-ideal, if \mathcal{I} is a thick subcategory of \mathcal{K} such that $\mathcal{K} \otimes \mathcal{I} \subseteq \mathcal{I}$
2. We say $\mathcal{P} \subseteq \mathcal{K}$ is a prime tensor ideal if, whenever $a \otimes b \in \mathcal{P}$ then either a or b is in \mathcal{P}

What's the Big Ideal?

Until further notice, let $(\mathcal{K}, \otimes, 1)$ be a closed, rigid tt category. Then, continuing our analogy of treating \mathcal{K} as a ring, we introduce the following definitions:

1. We say $\mathcal{I} \subseteq \mathcal{K}$ is a tensor-ideal, if \mathcal{I} is a thick subcategory of \mathcal{K} such that $\mathcal{K} \otimes \mathcal{I} \subseteq \mathcal{I}$
2. We say $\mathcal{P} \subseteq \mathcal{K}$ is a prime tensor ideal if, whenever $a \otimes b \in \mathcal{P}$ then either a or b is in \mathcal{P}
3. Let $(\mathcal{T}, \otimes, 1)$ be another closed, rigid, tt category. Then a tt functor $F : \mathcal{K} \rightarrow \mathcal{T}$ is a triangulated functor that is also "strong monoidal" (that is $F(a \otimes b) \cong F(a) \otimes F(b)$) with a whole bunch of compatibility axioms as well)

What's the Big Ideal?

Until further notice, let $(\mathcal{K}, \otimes, 1)$ be a closed, rigid tt category. Then, continuing our analogy of treating \mathcal{K} as a ring, we introduce the following definitions:

1. We say $\mathcal{I} \subseteq \mathcal{K}$ is a tensor-ideal, if \mathcal{I} is a thick subcategory of \mathcal{K} such that $\mathcal{K} \otimes \mathcal{I} \subseteq \mathcal{I}$
2. We say $\mathcal{P} \subseteq \mathcal{K}$ is a prime tensor ideal if, whenever $a \otimes b \in \mathcal{P}$ then either a or b is in \mathcal{P}
3. Let $(\mathcal{T}, \otimes, 1)$ be another closed, rigid, tt category. Then a tt functor $F : \mathcal{K} \rightarrow \mathcal{T}$ is a triangulated functor that is also "strong monoidal" (that is $F(a \otimes b) \cong F(a) \otimes F(b)$) with a whole bunch of compatibility axioms as well)

Rmk: The assumption that \mathcal{K} is rigid, and the remark about k, k^\vee being summands of respective tensors implies that a tensor ideal \mathcal{I} is automatically closed under taking duals, and is radical ($k^{\otimes n} \in \mathcal{I} \implies k \in \mathcal{I}$). This second part will be very important in a few slides.

Example of tt functor

We saw last talk that given a thick subcategory $\mathcal{C} \subseteq \mathcal{K}$ that we could form the quotient category \mathcal{K}/\mathcal{C} in such a way that it was triangulated, and where the universal quotient functor $q : \mathcal{K} \rightarrow \mathcal{K}/\mathcal{C}$ is a triangulated functor. We want to extend this construction to the tt world.

Example of tt functor

We saw last talk that given a thick subcategory $\mathcal{C} \subseteq \mathcal{K}$ that we could form the quotient category \mathcal{K}/\mathcal{C} in such a way that it was triangulated, and where the universal quotient functor $q : \mathcal{K} \rightarrow \mathcal{K}/\mathcal{C}$ is a triangulated functor. We want to extend this construction to the tt world.

1. Now let $\mathcal{I} \subseteq \mathcal{K}$ be a tt ideal. Then we can form the triangulated category \mathcal{K}/\mathcal{I} with universal functor $q : \mathcal{K} \rightarrow \mathcal{K}/\mathcal{I}$ just because \mathcal{I} is thick.

Example of tt functor

We saw last talk that given a thick subcategory $\mathcal{C} \subseteq \mathcal{K}$ that we could form the quotient category \mathcal{K}/\mathcal{C} in such a way that it was triangulated, and where the universal quotient functor $q : \mathcal{K} \rightarrow \mathcal{K}/\mathcal{C}$ is a triangulated functor. We want to extend this construction to the tt world.

1. Now let $\mathcal{I} \subseteq \mathcal{K}$ be a tt ideal. Then we can form the triangulated category \mathcal{K}/\mathcal{I} with universal functor $q : \mathcal{K} \rightarrow \mathcal{K}/\mathcal{I}$ just because \mathcal{I} is thick.
2. The point is, that now we have the following, two hopefully unsurprising, facts:
 - 2.1 \mathcal{K}/\mathcal{I} is a tt category.
 - 2.2 The universal functor $q : \mathcal{K} \rightarrow \mathcal{K}/\mathcal{I}$ is a tensor functor.

A Naive Goal

Given a tt category \mathcal{K} , the ultimate goal would be to classify all objects up to isomorphism. Unfortunately, such a desire is absolutely impossible in general. So we must resign ourselves to a slightly weaker version of classification.

A Naive Goal

Given a tt category \mathcal{K} , the ultimate goal would be to classify all objects up to isomorphism. Unfortunately, such a desire is absolutely impossible in general. So we must resign ourselves to a slightly weaker version of classification.

1. The main point is that we have more structure to play around with! We can take tensors, cones, direct sums, direct summands, suspensions, ect. So rather than demanding we classify an object X up to isomorphism, we ask to classify X "up to the tt structure"

A Naive Goal

Given a tt category \mathcal{K} , the ultimate goal would be to classify all objects up to isomorphism. Unfortunately, such a desire is absolutely impossible in general. So we must resign ourselves to a slightly weaker version of classification.

1. The main point is that we have more structure to play around with! We can take tensors, cones, direct sums, direct summands, suspensions, ect. So rather than demanding we classify an object X up to isomorphism, we ask to classify X "up to the tt structure"
2. In other words we want to know when we can reach an object Y from X using all the structure at hand. That is we want to know if Y is in the tensor ideal generated by X !

A Naive Goal

Given a tt category \mathcal{K} , the ultimate goal would be to classify all objects up to isomorphism. Unfortunately, such a desire is absolutely impossible in general. So we must resign ourselves to a slightly weaker version of classification.

1. The main point is that we have more structure to play around with! We can take tensors, cones, direct sums, direct summands, suspensions, ect. So rather than demanding we classify an object X up to isomorphism, we ask to classify X "up to the tt structure"
2. In other words we want to know when we can reach an object Y from X using all the structure at hand. That is we want to know if Y is in the tensor ideal generated by X !
3. Therefore, our main task is to classify thick tensor ideals of \mathcal{K} ! How to do this.....?

The Balmer Spectrum

The definitions given thus far have a strong bias towards commutative algebra. It is therefore not surprising that the Prime Ideals play a special role.

The Balmer Spectrum

The definitions given thus far have a strong bias towards commutative algebra. It is therefore not surprising that the Prime Ideals play a special role.

1. The Spectrum of a tt category is the set $\mathrm{Spc}(\mathcal{K}) = \{\mathcal{P} \subseteq \mathcal{K} : \mathcal{P} \text{ is a prime ideal}\}$

The Balmer Spectrum

The definitions given thus far have a strong bias towards commutative algebra. It is therefore not surprising that the Prime Ideals play a special role.

1. The Spectrum of a tt category is the set $\mathrm{Spc}(\mathcal{K}) = \{\mathcal{P} \subseteq \mathcal{K} : \mathcal{P} \text{ is a prime ideal}\}$
2. The support of an object k is the collection of all prime ideals k is NOT in, $\mathrm{supp}(k) = \{\mathcal{P} \in \mathrm{Spc}(\mathcal{K}) : k \notin \mathcal{P}\}$
 - 2.1 Rmk: Recall the kernel of the universal functor $q : \mathcal{K} \rightarrow \mathcal{K}/\mathcal{P}$ is precisely \mathcal{P} . So saying $k \notin \mathcal{P}$ amounts to saying $k \neq 0$ in the tt category \mathcal{K}/\mathcal{P}

The Balmer Spectrum

The definitions given thus far have a strong bias towards commutative algebra. It is therefore not surprising that the Prime Ideals play a special role.

1. The Spectrum of a tt category is the set $\mathrm{Spc}(\mathcal{K}) = \{\mathcal{P} \subseteq \mathcal{K} : \mathcal{P} \text{ is a prime ideal}\}$
2. The support of an object k is the collection of all prime ideals k is NOT in,
 $\mathrm{supp}(k) = \{\mathcal{P} \in \mathrm{Spc}(\mathcal{K}) : k \notin \mathcal{P}\}$
 - 2.1 Rmk: Recall the kernel of the universal functor $q : \mathcal{K} \rightarrow \mathcal{K}/\mathcal{P}$ is precisely \mathcal{P} . So saying $k \notin \mathcal{P}$ amounts to saying $k \neq 0$ in the tt category \mathcal{K}/\mathcal{P}
3. We make $\mathrm{Spc}(\mathcal{K})$ into a topological space by giving letting $\{\mathrm{supp}(k)\}_{k \in \mathcal{K}}$ be a basis of closed subsets.

The Balmer Spectrum

The definitions given thus far have a strong bias towards commutative algebra. It is therefore not surprising that the Prime Ideals play a special role.

1. The Spectrum of a tt category is the set $\mathrm{Spc}(\mathcal{K}) = \{\mathcal{P} \subseteq \mathcal{K} : \mathcal{P} \text{ is a prime ideal}\}$
2. The support of an object k is the collection of all prime ideals k is NOT in, $\mathrm{supp}(k) = \{\mathcal{P} \in \mathrm{Spc}(\mathcal{K}) : k \notin \mathcal{P}\}$
 - 2.1 Rmk: Recall the kernel of the universal functor $q : \mathcal{K} \rightarrow \mathcal{K}/\mathcal{P}$ is precisely \mathcal{P} . So saying $k \notin \mathcal{P}$ amounts to saying $k \neq 0$ in the tt category \mathcal{K}/\mathcal{P}
3. We make $\mathrm{Spc}(\mathcal{K})$ into a topological space by giving letting $\{\mathrm{supp}(k)\}_{k \in \mathcal{K}}$ be a basis of closed subsets.
4. We shall see that classifying the tt ideals of \mathcal{K} more or less amounts to classifying nice subsets of $\mathrm{Spc}(\mathcal{K})$. Let us first see some basic properties of the Spectrum.

More on Balmer Spectrum

It should hopefully come as no surprise that the Balmer spectrum satisfies a universal property. To state what the universal property is, we need a few results first: Let $\mathcal{K}, \mathrm{Spc}(\mathcal{K}), \mathrm{supp}(\mathbf{k})$ be as above. Then,

More on Balmer Spectrum

It should hopefully come as no surprise that the Balmer spectrum satisfies a universal property. To state what the universal property is, we need a few results first: Let $\mathcal{K}, \mathrm{Spc}(\mathcal{K}), \mathrm{supp}(k)$ be as above. Then,

1. $\mathrm{supp}(1) = \mathrm{Spc}(\mathcal{K})$ and $\mathrm{supp}(0) = \emptyset$

More on Balmer Spectrum

It should hopefully come as no surprise that the Balmer spectrum satisfies a universal property. To state what the universal property is, we need a few results first: Let $\mathcal{K}, \mathrm{Spc}(\mathcal{K}), \mathrm{supp}(k)$ be as above. Then,

1. $\mathrm{supp}(1) = \mathrm{Spc}(\mathcal{K})$ and $\mathrm{supp}(0) = \emptyset$
2. $\mathrm{supp}(a \oplus b) = \mathrm{supp}(a) \cup \mathrm{supp}(b)$

More on Balmer Spectrum

It should hopefully come as no surprise that the Balmer spectrum satisfies a universal property. To state what the universal property is, we need a few results first: Let $\mathcal{K}, \mathrm{Spc}(\mathcal{K}), \mathrm{supp}(k)$ be as above. Then,

1. $\mathrm{supp}(1) = \mathrm{Spc}(\mathcal{K})$ and $\mathrm{supp}(0) = \emptyset$
2. $\mathrm{supp}(a \oplus b) = \mathrm{supp}(a) \cup \mathrm{supp}(b)$
3. $\mathrm{supp}(\Sigma x) = \mathrm{supp}(x)$ for all x

More on Balmer Spectrum

It should hopefully come as no surprise that the Balmer spectrum satisfies a universal property. To state what the universal property is, we need a few results first: Let $\mathcal{K}, \mathrm{Spc}(\mathcal{K}), \mathrm{supp}(k)$ be as above. Then,

1. $\mathrm{supp}(1) = \mathrm{Spc}(\mathcal{K})$ and $\mathrm{supp}(0) = \emptyset$
2. $\mathrm{supp}(a \oplus b) = \mathrm{supp}(a) \cup \mathrm{supp}(b)$
3. $\mathrm{supp}(\Sigma x) = \mathrm{supp}(x)$ for all x
4. $\mathrm{supp}(z) \subseteq \mathrm{supp}(x) \cup \mathrm{supp}(y)$ for any distinguished triangle $x \rightarrow y \rightarrow z \rightarrow \Sigma x$

More on Balmer Spectrum

It should hopefully come as no surprise that the Balmer spectrum satisfies a universal property. To state what the universal property is, we need a few results first: Let $\mathcal{K}, \mathrm{Spc}(\mathcal{K}), \mathrm{supp}(k)$ be as above. Then,

1. $\mathrm{supp}(1) = \mathrm{Spc}(\mathcal{K})$ and $\mathrm{supp}(0) = \emptyset$
2. $\mathrm{supp}(a \oplus b) = \mathrm{supp}(a) \cup \mathrm{supp}(b)$
3. $\mathrm{supp}(\Sigma x) = \mathrm{supp}(x)$ for all x
4. $\mathrm{supp}(z) \subseteq \mathrm{supp}(x) \cup \mathrm{supp}(y)$ for any distinguished triangle $x \rightarrow y \rightarrow z \rightarrow \Sigma x$
5. $\mathrm{supp}(x \otimes y) = \mathrm{supp}(x) \cap \mathrm{supp}(y)$

More on Balmer Spectrum

It should hopefully come as no surprise that the Balmer spectrum satisfies a universal property. To state what the universal property is, we need a few results first: Let $\mathcal{K}, \mathrm{Spc}(\mathcal{K}), \mathrm{supp}(k)$ be as above. Then,

1. $\mathrm{supp}(1) = \mathrm{Spc}(\mathcal{K})$ and $\mathrm{supp}(0) = \emptyset$
2. $\mathrm{supp}(a \oplus b) = \mathrm{supp}(a) \cup \mathrm{supp}(b)$
3. $\mathrm{supp}(\Sigma x) = \mathrm{supp}(x)$ for all x
4. $\mathrm{supp}(z) \subseteq \mathrm{supp}(x) \cup \mathrm{supp}(y)$ for any distinguished triangle $x \rightarrow y \rightarrow z \rightarrow \Sigma x$
5. $\mathrm{supp}(x \otimes y) = \mathrm{supp}(x) \cap \mathrm{supp}(y)$

We say a "Support Data" on a tt category is a pair (X, σ) where X is a topological space and $\sigma(k)$ is a closed subset in X for each $k \in \mathcal{K}$ satisfying conditions 1-5 above.

A morphism of support data $(X, \sigma) \xrightarrow{f} (Y, \beta)$ is a continuous map $X \rightarrow Y$ such that $f^{-1}(\beta(k)) = \sigma(k)$.

More on Balmer Spectrum

It should hopefully come as no surprise that the Balmer spectrum satisfies a universal property. To state what the universal property is, we need a few results first: Let $\mathcal{K}, \mathrm{Spc}(\mathcal{K}), \mathrm{supp}(k)$ be as above. Then,

1. $\mathrm{supp}(1) = \mathrm{Spc}(\mathcal{K})$ and $\mathrm{supp}(0) = \emptyset$
2. $\mathrm{supp}(a \oplus b) = \mathrm{supp}(a) \cup \mathrm{supp}(b)$
3. $\mathrm{supp}(\Sigma x) = \mathrm{supp}(x)$ for all x
4. $\mathrm{supp}(z) \subseteq \mathrm{supp}(x) \cup \mathrm{supp}(y)$ for any distinguished triangle $x \rightarrow y \rightarrow z \rightarrow \Sigma x$
5. $\mathrm{supp}(x \otimes y) = \mathrm{supp}(x) \cap \mathrm{supp}(y)$

We say a "Support Data" on a tt category is a pair (X, σ) where X is a topological space and $\sigma(k)$ is a closed subset in X for each $k \in \mathcal{K}$ satisfying conditions 1-5 above.

A morphism of support data $(X, \sigma) \xrightarrow{f} (Y, \beta)$ is a continuous map $X \rightarrow Y$ such that $f^{-1}(\beta(k)) = \sigma(k)$.

Thrm : $(\mathrm{Spc}(\mathcal{K}), \mathrm{supp})$ is the terminal support data on \mathcal{K}

Some basic properties

With the Balmer spectrum in hand, let us now see some basic applications.

Some basic properties

With the Balmer spectrum in hand, let us now see some basic applications.

1. We have $\mathrm{supp}(k) = \emptyset \iff k \cong 0$ (this uses that \mathcal{K} is rigid)

Some basic properties

With the Balmer spectrum in hand, let us now see some basic applications.

1. We have $\text{supp}(\mathbf{k}) = \emptyset \iff \mathbf{k} \cong 0$ (this uses that \mathcal{K} is rigid)
2. Let $\mathcal{P} \in \text{Spc}(\mathcal{K})$. Then $\overline{\{\mathcal{P}\}} = \{\mathcal{Q} \in \text{Spc}(\mathcal{K}) : \mathcal{Q} \subseteq \mathcal{P}\}$. In particular $\overline{\{\mathcal{P}_1\}} = \overline{\{\mathcal{P}_2\}} \iff \mathcal{P}_1 = \mathcal{P}_2$. We say such a space is T_0

Some basic properties

With the Balmer spectrum in hand, let us now see some basic applications.

1. We have $\mathrm{supp}(\mathbf{k}) = \emptyset \iff \mathbf{k} \cong 0$ (this uses that \mathcal{K} is rigid)
2. Let $\mathcal{P} \in \mathrm{Spc}(\mathcal{K})$. Then $\overline{\{\mathcal{P}\}} = \{\mathcal{Q} \in \mathrm{Spc}(\mathcal{K}) : \mathcal{Q} \subseteq \mathcal{P}\}$. In particular $\overline{\{\mathcal{P}_1\}} = \overline{\{\mathcal{P}_2\}} \iff \mathcal{P}_1 = \mathcal{P}_2$. We say such a space is T_0
3. Moreover, $\mathrm{Spc}(\mathcal{K})$ is a "Spectral" topological space: that is, it is T_0 ; it is quasi-compact; the quasi-compact open subsets of $\mathrm{Spc}(\mathcal{K})$ are closed under finite intersections and form an open basis of $\mathrm{Spc}(\mathcal{K})$; and every non-empty irreducible closed subset of $\mathrm{Spc}(\mathcal{K})$ has a generic point.

Some basic properties

With the Balmer spectrum in hand, let us now see some basic applications.

1. We have $\text{supp}(\mathbf{k}) = \emptyset \iff \mathbf{k} \cong 0$ (this uses that \mathcal{K} is rigid)
2. Let $\mathcal{P} \in \text{Spc}(\mathcal{K})$. Then $\overline{\{\mathcal{P}\}} = \{\mathcal{Q} \in \text{Spc}(\mathcal{K}) : \mathcal{Q} \subseteq \mathcal{P}\}$. In particular $\overline{\{\mathcal{P}_1\}} = \overline{\{\mathcal{P}_2\}} \iff \mathcal{P}_1 = \mathcal{P}_2$. We say such a space is T_0
3. Moreover, $\text{Spc}(\mathcal{K})$ is a "Spectral" topological space: that is, it is T_0 ; it is quasi-compact; the quasi-compact open subsets of $\text{Spc}(\mathcal{K})$ are closed under finite intersections and form an open basis of $\text{Spc}(\mathcal{K})$; and every non-empty irreducible closed subset of $\text{Spc}(\mathcal{K})$ has a generic point.
4. The assignment $\mathcal{K} \rightarrow \text{Spc}(\mathcal{K})$ is a contravariant functor. Given a tt functor $\mathcal{K} \xrightarrow{\mathbb{F}} \mathcal{T}$ we get a continuous (spectral) map $\text{Spc}(\mathcal{T}) \xrightarrow{\text{Spc}(\mathbb{F})} \text{Spc}(\mathcal{K})$

Thomason Subsets and the Classification Thrm

Remember, our goal is to classify tt ideals of \mathcal{K} . We need one final topological notion before we can state the classification.

Thomason Subsets and the Classification Thrm

Remember, our goal is to classify tt ideals of \mathcal{K} . We need one final topological notion before we can state the classification.

1. A Thomason subset $V \subseteq X$ of a spectral topological space X is a subset of the form $V = \bigcup_{\alpha} V_{\alpha}$ where each V_{α} is closed with quasi-compact open complements.

Thomason Subsets and the Classification Thrm

Remember, our goal is to classify tt ideals of \mathcal{K} . We need one final topological notion before we can state the classification.

1. A Thomason subset $V \subseteq X$ of a spectral topological space X is a subset of the form $V = \bigcup_{\alpha} V_{\alpha}$ where each V_{α} is closed with quasi-compact open complements.
2. Now recall, $\mathrm{Spc}(\mathcal{K})$ is spectral. So let $V \subseteq \mathrm{Spc}(\mathcal{K})$ be a Thomason subset, and let $\mathcal{K}_V = \{x \in \mathcal{K} : \mathrm{supp}(x) \subseteq V\}$. It turns out this is a tt ideal.

Thomason Subsets and the Classification Thrm

Remember, our goal is to classify tt ideals of \mathcal{K} . We need one final topological notion before we can state the classification.

1. A Thomason subset $V \subseteq X$ of a spectral topological space X is a subset of the form $V = \bigcup_{\alpha} V_{\alpha}$ where each V_{α} is closed with quasi-compact open complements.
2. Now recall, $\mathrm{Spc}(\mathcal{K})$ is spectral. So let $V \subseteq \mathrm{Spc}(\mathcal{K})$ be a Thomason subset, and let $\mathcal{K}_V = \{x \in \mathcal{K} : \mathrm{supp}(x) \subseteq V\}$. It turns out this is a tt ideal.
3. Now let $\mathcal{P} \in \mathrm{Spc}(\mathcal{K})$, and let $V = \bigcup_{x \in \mathcal{P}} \mathrm{supp}(x)$. Then V is a Thomason subset of $\mathrm{Spc}(\mathcal{K})$. Moreover,

Thomason Subsets and the Classification Thrm

Remember, our goal is to classify tt ideals of \mathcal{K} . We need one final topological notion before we can state the classification.

1. A Thomason subset $V \subseteq X$ of a spectral topological space X is a subset of the form $V = \bigcup_{\alpha} V_{\alpha}$ where each V_{α} is closed with quasi-compact open complements.
2. Now recall, $\mathrm{Spc}(\mathcal{K})$ is spectral. So let $V \subseteq \mathrm{Spc}(\mathcal{K})$ be a Thomason subset, and let $\mathcal{K}_V = \{x \in \mathcal{K} : \mathrm{supp}(x) \subseteq V\}$. It turns out this is a tt ideal.
3. Now let $\mathcal{P} \in \mathrm{Spc}(\mathcal{K})$, and let $V = \bigcup_{x \in \mathcal{P}} \mathrm{supp}(x)$. Then V is a Thomason subset of $\mathrm{Spc}(\mathcal{K})$. Moreover,
4. **The two assignments above give an order preserving bijection between**

$$\mathrm{Thom}(\mathrm{Spc}(\mathcal{K})) \leftrightarrow \mathrm{Thick}^{\otimes}(\mathcal{K})$$

the Thomason subsets of $\mathrm{Spc}(\mathcal{K})$ and the set of tt ideals in \mathcal{K} .

Applications

Before we provide some of the more classical examples of classifications, let us give some consequences of this notion of Spc.

1. Let $\langle x \rangle$ denote the tt ideal generated by an object $x \in \mathcal{K}$. Then we have that $y \in \langle x \rangle \iff \text{supp}(y) \subseteq \text{supp}(x)!!$

Applications

Before we provide some of the more classical examples of classifications, let us give some consequences of this notion of Spc.

1. Let $\langle x \rangle$ denote the tt ideal generated by an object $x \in \mathcal{K}$. Then we have that $y \in \langle x \rangle \iff \text{supp}(y) \subseteq \text{supp}(x)!!$
2. Suppose $\text{supp}(x) = y_1 \sqcup y_2$. Then x decomposes as $x = x_1 \oplus x_2$ with $\text{supp}(x_i) = y_i$ (has applications to the Pickard Group of \otimes -invertible objects)

Applications

Before we provide some of the more classical examples of classifications, let us give some consequences of this notion of Spc.

1. Let $\langle x \rangle$ denote the tt ideal generated by an object $x \in \mathcal{K}$. Then we have that $y \in \langle x \rangle \iff \text{supp}(y) \subseteq \text{supp}(x)!!$
2. Suppose $\text{supp}(x) = y_1 \sqcup y_2$. Then x decomposes as $x = x_1 \oplus x_2$ with $\text{supp}(x_i) = y_i$ (has applications to the Pickard Group of \otimes -invertible objects).
3. Suppose $\text{supp}(x) \cap \text{supp}(y) = \emptyset$. Then we have $\text{Hom}_{\mathcal{K}}(x, y) = 0$

Some Consequences of Spc

Applications

Before we provide some of the more classical examples of classifications, let us give some consequences of this notion of Spc.

1. Let $\langle x \rangle$ denote the tt ideal generated by an object $x \in \mathcal{K}$. Then we have that $y \in \langle x \rangle \iff \text{supp}(y) \subseteq \text{supp}(x)!!$
2. Suppose $\text{supp}(x) = y_1 \sqcup y_2$. Then x decomposes as $x = x_1 \oplus x_2$ with $\text{supp}(x_i) = y_i$ (has applications to the Pickard Group of \otimes -invertible objects).
3. Suppose $\text{supp}(x) \cap \text{supp}(y) = \emptyset$. Then we have $\text{Hom}_{\mathcal{K}}(x, y) = 0$

Now, recall that we noticed that $\text{Spc}(\mathcal{K})$ is a "spectral space." It is a theorem that every spectral space is isomorphic to $\text{Spec}(R)$ for some commutative ring R . In any tt-category, one can take the endomorphism ring of the unit, and get a commutative ring. So we might ask,

1. Is $\text{Spc}(\mathcal{K}) \cong \text{Spec}(\text{End}_{\mathcal{K}}(1))$?
2. Or more generally, is there a map $\text{Spc}(\mathcal{K}) \rightarrow \text{Spec}(\text{End}_{\mathcal{K}}(1))$.

Some Consequences of Spc

Applications

Before we provide some of the more classical examples of classifications, let us give some consequences of this notion of Spc.

1. Let $\langle x \rangle$ denote the tt ideal generated by an object $x \in \mathcal{K}$. Then we have that $y \in \langle x \rangle \iff \text{supp}(y) \subseteq \text{supp}(x)!!$
2. Suppose $\text{supp}(x) = y_1 \sqcup y_2$. Then x decomposes as $x = x_1 \oplus x_2$ with $\text{supp}(x_i) = y_i$
3. Suppose $\text{supp}(x) \cap \text{supp}(y) = \emptyset$. Then we have $\text{Hom}_{\mathcal{K}}(x, y) = 0$

Now, recall that we noticed that $\text{Spc}(\mathcal{K})$ is a "spectral space." It is a theorem that every spectral space is isomorphic to $\text{Spec}(R)$ for some commutative ring R . In any tt-category, one can take the endomorphism ring of the unit, and get a commutative ring. So we might ask,

1. Is $\text{Spc}(\mathcal{K}) \cong \text{Spec}(\text{End}_{\mathcal{K}}(1))$?
2. Or more generally, is there a map $\text{Spc}(\mathcal{K}) \rightarrow \text{Spec}(\text{End}_{\mathcal{K}}(1))$.
3. Thrm: There exist two natural continuous maps
 - 3.1 $\rho_{\mathcal{K}} : \text{Spc}(\mathcal{K}) \rightarrow \text{Spec}(\text{End}_{\mathcal{K}}(1))$.
 - 3.2 $\rho_{\mathcal{K}}^{\bullet} : \text{Spc}(\mathcal{K}) \rightarrow \text{Spec}^h(\text{End}_{\mathcal{K}}(1))$.

Some Consequences of Spc

Applications

Before we provide some of the more classical examples of classifications, let us give some consequences of this notion of Spc.

1. Let $\langle x \rangle$ denote the tt ideal generated by an object $x \in \mathcal{K}$. Then we have that $y \in \langle x \rangle \iff \text{supp}(y) \subseteq \text{supp}(x)!!$
2. Suppose $\text{supp}(x) = y_1 \sqcup y_2$. Then x decomposes as $x = x_1 \oplus x_2$ with $\text{supp}(x_i) = y_i$
3. Suppose $\text{supp}(x) \cap \text{supp}(y) = \emptyset$. Then we have $\text{Hom}_{\mathcal{K}}(x, y) = 0$

Now, recall that we noticed that $\text{Spc}(\mathcal{K})$ is a "spectral space." It is a theorem that every spectral space is isomorphic to $\text{Spec}(R)$ for some commutative ring R . In any tt-category, one can take the endomorphism ring of the unit, and get a commutative ring. So we might ask,

1. Is $\text{Spc}(\mathcal{K}) \cong \text{Spec}(\text{End}_{\mathcal{K}}(1))$?
2. Or more generally, is there a map $\text{Spc}(\mathcal{K}) \rightarrow \text{Spec}(\text{End}_{\mathcal{K}}(1))$.
3. Thrm: There exist two natural continuous maps
 - 3.1 $\rho_{\mathcal{K}} : \text{Spc}(\mathcal{K}) \rightarrow \text{Spec}(\text{End}_{\mathcal{K}}(1))$.
 - 3.2 $\rho_{\mathcal{K}}^{\bullet} : \text{Spc}(\mathcal{K}) \rightarrow \text{Spec}^h(\text{End}_{\mathcal{K}}^{\bullet}(1))$.

Moreover, these maps are very often surjective. We call these maps the comparison maps, and we will make use of them shortly.

Derived Category of a (nice enough) Scheme

Reproducing a Scheme

1. Thrm (Thomason, Neeman): Let X be a quasi-compact, quasi separated scheme, and let $\mathcal{K} = D^{\text{perf}}(X)$ be the derived category of perfect complexes. Then the Spectrum of \mathcal{K} is isomorphic to the underlying scheme itself $|X|$ via a homeomorphism $|X| \xrightarrow{\sim} \text{Spc}(D^{\text{perf}}(X))$ given by:

Derived Category of a (nice enough) Scheme

Reproducing a Scheme

1. Thrm (Thomason, Neeman): Let X be a quasi-compact, quasi separated scheme, and let $\mathcal{K} = D^{\text{perf}}(X)$ be the derived category of perfect complexes. Then the Spectrum of \mathcal{K} is isomorphic to the underlying scheme itself $|X|$ via a homeomorphism $|X| \xrightarrow{\sim} \text{Spc}(D^{\text{perf}}(X))$ given by:

$$x \rightarrow \mathcal{P}(x) = \{Y \in \mathcal{K} : Y_x \cong 0\}$$

where $\mathcal{P}(x)$ is the kernel of the residue field functor $D^{\text{perf}}(X) \rightarrow D^b(k(x))$ for each point x .

Derived Category of a (nice enough) Scheme

Reproducing a Scheme

1. Thrm (Thomason, Neeman): Let X be a quasi-compact, quasi separated scheme, and let $\mathcal{K} = D^{\text{perf}}(X)$ be the derived category of perfect complexes. Then the Spectrum of \mathcal{K} is isomorphic to the underlying scheme itself $|X|$ via a homeomorphism $|X| \xrightarrow{\sim} \text{Spc}(D^{\text{perf}}(X))$ given by:

$$x \rightarrow \mathcal{P}(x) = \{Y \in \mathcal{K} : Y_x \cong 0\}$$

where $\mathcal{P}(x)$ is the kernel of the residue field functor $D^{\text{perf}}(X) \rightarrow D^b(k(x))$ for each point x .

2. Affine Case: In particular, suppose A is a commutative ring, and let $X = \text{Spec}(A)$ be an affine scheme. Then we get $\text{Spc}(D^{\text{perf}}(A)) \cong \text{Spec}(A)$ (such a result also holds if A is commutative-graded, by replacing $\text{Spec}(A)$ with $\text{Spec}^h(A)$ the homogeneous spectrum)

Derived Category of a (nice enough) Scheme

Reproducing a Scheme

1. Thrm (Thomason, Neeman): Let X be a quasi-compact, quasi separated scheme, and let $\mathcal{K} = D^{\text{perf}}(X)$ be the derived category of perfect complexes. Then the Spectrum of \mathcal{K} is isomorphic to the underlying scheme itself $|X|$ via a homeomorphism $|X| \xrightarrow{\sim} \text{Spc}(D^{\text{perf}}(X))$ given by:

$$x \rightarrow \mathcal{P}(x) = \{Y \in \mathcal{K} : Y_x \cong 0\}$$

where $\mathcal{P}(x)$ is the kernel of the residue field functor $D^{\text{perf}}(X) \rightarrow D^b(k(x))$ for each point x .

2. Affine Case: In particular, suppose A is a commutative ring, and let $X = \text{Spec}(A)$ be an affine scheme. Then we get $\text{Spc}(D^{\text{perf}}(A)) \cong \text{Spec}(A)$ (such a result also holds if A is commutative- graded, by replacing $\text{Spec}(A)$ with $\text{Spec}^h(A)$ the homogeneous spectrum)
3. Moreover, one can always equip $\text{Spc}(\mathcal{K})$ with a sheaf of commutative rings, and in this case one recovers the structure sheaf of X , \mathcal{O}_X

Derived Category of a (nice enough) Scheme

Reproducing a Scheme

1. Thrm (Thomason, Neeman): Let X be a quasi-compact, quasi separated scheme, and let $\mathcal{K} = D^{\text{perf}}(X)$ be the derived category of perfect complexes. Then the Spectrum of \mathcal{K} is isomorphic to the underlying scheme itself $|X|$ via a homeomorphism $|X| \xrightarrow{\sim} \text{Spc}(D^{\text{perf}}(X))$ given by:

$$x \rightarrow \mathcal{P}(x) = \{Y \in \mathcal{K} : Y_x \cong 0\}$$

where $\mathcal{P}(x)$ is the kernel of the residue field functor $D^{\text{perf}}(X) \rightarrow D^b(k(x))$ for each point x .

2. Affine Case: In particular, suppose A is a commutative ring, and let $X = \text{Spec}(A)$ be an affine scheme. Then we get $\text{Spc}(D^{\text{perf}}(A)) \cong \text{Spec}(A)$. (such a result also holds if A is commutative- graded, by replacing $\text{Spec}(A)$ with $\text{Spec}^h(A)$ the homogeneous spectrum)
3. Moreover, one can always equip $\text{Spc}(\mathcal{K})$ with a sheaf of commutative rings, and in this case one recovers the structure sheaf of X , \mathcal{O}_X
4. There are generalizations of these to (nice enough) stacks and singularity categories a la Stevenson. They are beyond my current knowledge however.

Support Varieties

Let G be a finite group and k be a field of characteristic p dividing the order of G . Then consider $\mathcal{K} = \text{stab}(kG)$ the category of finite dimensional kG -modules, and recall that we can identify $\text{stab}(kG)$ as the Verdier quotient $\text{stab}(kG) \cong D^b(kG)/D^{\text{perf}}(kG)$ (see my last talk)

Support Varieties

Let G be a finite group and k be a field of characteristic p dividing the order of G . Then consider $\mathcal{K} = \text{stab}(kG)$ the category of finite dimensional kG -modules, and recall that we can identify $\text{stab}(kG)$ as the Verdier quotient $\text{stab}(kG) \cong D^b(kG)/D^{\text{perf}}(kG)$ (see my last talk)

1. There is a homeomorphism between the spectrum of \mathcal{K} and the projective support variety of G , $\text{Spc}(\text{stab}(kG)) \cong \text{Proj}(H^\bullet(G, k))$.

Support Varieties

Let G be a finite group and k be a field of characteristic p dividing the order of G . Then consider $\mathcal{K} = \text{stab}(kG)$ the category of finite dimensional kG -modules, and recall that we can identify $\text{stab}(kG)$ as the Verdier quotient $\text{stab}(kG) \cong D^b(kG)/D^{\text{perf}}(kG)$ (see my last talk)

1. There is a homeomorphism between the spectrum of \mathcal{K} and the projective support variety of G , $\text{Spc}(\text{stab}(kG)) \cong \text{Proj}(H^\bullet(G, k))$.
2. Furthermore, this homeomorphism can be extended to a homeomorphism $\text{Spc}(D^b(kG)) \cong \text{Spec}^h(H^\bullet(G, k))$ by adding exactly one closed point.

Support Varieties

Let G be a finite group and k be a field of characteristic p dividing the order of G . Then consider $\mathcal{K} = \text{stab}(kG)$ the category of finite dimensional kG -modules, and recall that we can identify $\text{stab}(kG)$ as the Verdier quotient $\text{stab}(kG) \cong D^b(kG)/D^{\text{per}f}(kG)$ (see my last talk)

1. There is a homeomorphism between the spectrum of \mathcal{K} and the projective support variety of G , $\text{Spc}(\text{stab}(kG)) \cong \text{Proj}(H^\bullet(G, k))$.
2. Furthermore, this homeomorphism can be extended to a homeomorphism $\text{Spc}(D^b(kG)) \cong \text{Spec}^h(H^\bullet(G, k))$ by adding exactly one closed point.
3. Rmk: The two above homeomorphisms hold for G a finite group scheme as well.

The Original classification

Now, the initial tt Category was SH^c and the classification of tt ideals in that case, in some sense motivated this entire discussion. The classification in this case is a little more complicated to state however, so let us first provide some context and notation to be used.

The Original classification

Now, the initial tt Category was SH^c and the classification of tt ideals in that case, in some sense motivated this entire discussion. The classification in this case is a little more complicated to state however, so let us first provide some context and notation to be used.

1. We will consider the "p-local Stable Homotopy category". This consists of objects in SH^c such that $\pi_*(X) \otimes \mathbb{Z}_{(p)} \cong \pi_*(X)$. Denote this subcategory by SH_p^c . This is the Bousfield Localization $q : \mathrm{SH}^c \rightarrow \mathrm{SH}_p^c$ with respect to the Homology theory $\pi_\bullet(-) \otimes \mathbb{Z}_{(p)}$

The Original classification

Now, the initial tt Category was SH^c and the classification of tt ideals in that case, in some sense motivated this entire discussion. The classification in this case is a little more complicated to state however, so let us first provide some context and notation to be used.

1. We will consider the " p -local Stable Homotopy category". This consists of objects in SH^c such that $\pi_\bullet(X) \otimes \mathbb{Z}_{(p)} \cong \pi_\bullet(X)$. Denote this subcategory by SH_p^c . This is the Bousfield Localization $q : \mathrm{SH}^c \rightarrow \mathrm{SH}_p^c$ with respect to the Homology theory $\pi_\bullet(-) \otimes \mathbb{Z}_{(p)}$
2. For each integer $n \geq 1$ and prime p , there is a Homology theory, called Morava k -theory, denoted as $K_{p,n} : \mathrm{SH}_p^c \rightarrow \mathbb{F}_p[v_n, v_n^{-1}] - \mathrm{Mod}$.
Let us denote $\mathcal{C}_{p,n} := q^{-1}(\ker(K_{p,n}))$ and $\mathcal{C}_{0,1}$ to be the kernel of the rationalization functor $\pi_\bullet(-) \otimes \mathbb{Q} \cong H_\bullet(-, \mathbb{Q}) : \mathrm{SH}^c \rightarrow \mathrm{SH}_{\mathbb{Q}}^c \cong D^b(\mathbb{Q})$

The Original classification

Now, the initial tt Category was SH^c and the classification of tt ideals in that case, in some sense motivated this entire discussion. The classification in this case is a little more complicated to state however, so let us first provide some context and notation to be used.


1. We will consider the "p-local Stable Homotopy category". This consists of objects in SH^c such that $\pi_*(X) \otimes \mathbb{Z}_{(p)} \cong \pi_*(X)$. Denote this subcategory by SH_p^c . It turns out this can be realized as a Bousfield Localization $q : \mathrm{SH}^c \rightarrow \mathrm{SH}_p^c$
2. For each integer $n \geq 1$ and prime p , there is a Homology theory, called Morava k -theory, denoted as $K_{p,n} : \mathrm{SH}_p^c \rightarrow \mathbb{F}_p[v_n, v_n^{-1}] - \mathrm{Mod}$.
Let us denote $\mathcal{C}_{p,n} := q^{-1}(\ker(K_{p,n}))$ and $\mathcal{C}_{0,1}$ to be the kernel of the rationalization functor $\pi_*(-) \otimes \mathbb{Q} \cong H_*(-, \mathbb{Q}) : \mathrm{SH}^c \rightarrow \mathrm{SH}_{\mathbb{Q}}^c \cong D^b(\mathbb{Q})$
3. Recall the "comparison map" defined some slides ago: $\rho : \mathrm{Spc}(\mathcal{K}) \rightarrow \mathrm{Spec}(\mathrm{End}_{\mathcal{K}}(1))$. In this case, the unit is $1 = \mathbb{S}^0$ and $\mathrm{End}_{\mathcal{K}}(\mathbb{S}^0) = \mathbb{Z}$

The Original classification

Now, the initial tt Category was SH^c and the classification of tt ideals in that case, in some sense motivated this entire discussion. The classification in this case is a little more complicated to state however, so let us first provide some context and notation to be used.

1. We will consider the "p-local Stable Homotopy category". This consists of objects in SH^c such that $\pi_{\bullet}(X) \otimes \mathbb{Z}_{(p)} \cong \pi_{\bullet}(X)$. Denote this subcategory by SH_p^c . It turns out this can be realized as a Bousfield Localization $q : \mathrm{SH}^c \rightarrow \mathrm{SH}_p^c$
2. For each integer $n \geq 1$ and prime p , there is a Homology theory, called Morava k -theory, denoted as $K_{p,n} : \mathrm{SH}_p^c \rightarrow \mathbb{F}_p[v_n, v_n^{-1}] - \mathrm{Mod}$.
Let us denote $\mathcal{C}_{p,n} := q^{-1}(\ker(K_{p,n}))$ and $\mathcal{C}_{0,1}$ to be the kernel of the rationalization functor $\pi_{\bullet}(-) \otimes \mathbb{Q} \cong H_{\bullet}(-, \mathbb{Q}) : \mathrm{SH}^c \rightarrow \mathrm{SH}_{\mathbb{Q}}^c \cong D^b(\mathbb{Q})$
3. Recall the "comparison map" defined some slides ago: $\rho : \mathrm{Spc}(\mathcal{K}) \rightarrow \mathrm{Spec}(\mathrm{End}_{\mathcal{K}}(1))$. In this case, the unit is $1 = \mathbb{S}^0$ and $\mathrm{End}_{\mathcal{K}}(\mathbb{S}^0) = \mathbb{Z}$
4. Then it turns out the Spectrum of SH^c is given by pulling back this comparison map $\rho_{\mathcal{K}}$. The picture is as follows:

The Original classification



The Original classification

In the above picture, a line indicates that the higher prime is in the closure of the lower one. We have more precisely:

1. The pre-image of the dense point $(0) \in \operatorname{Spec}(\mathbb{Z})$ is the single point $\mathcal{C}_{0,1}$

The Original classification

In the above picture, a line indicates that the higher prime is in the closure of the lower one. We have more precisely:

1. The pre-image of the dense point $(0) \in \operatorname{Spec}(\mathbb{Z})$ is the single point $\mathcal{C}_{0,1}$
2. For each prime p , the pre-image of $p\mathbb{Z}$ consists of the column $\mathcal{C}_{p,n}$ and where $\mathcal{C}_{p,\infty} = \ker(q_p : \operatorname{SH}^c \rightarrow \operatorname{SH}_p^c)$.

The Original classification

In the above picture, a line indicates that the higher prime is in the closure of the lower one. We have more precisely:

1. The pre-image of the dense point $(0) \in \operatorname{Spec}(\mathbb{Z})$ is the single point $\mathcal{C}_{0,1}$
2. For each prime p , the pre-image of $p\mathbb{Z}$ consists of the column $\mathcal{C}_{p,n}$ and where $\mathcal{C}_{p,\infty} = \ker(q_p : \operatorname{SH}^c \rightarrow \operatorname{SH}_p^c)$.
3. $\mathcal{C}_{0,1}$ is the unique dense point in $\operatorname{Spc}(\operatorname{SH}^c)$. For each prime p and integer $1 \leq n \neq \infty$ we have the closure $\overline{\{\mathcal{C}_{p,n}\}} = \{\mathcal{C}_{p,m} : n \leq m \leq \infty\}$. The closed points of $\operatorname{Spc}(\operatorname{SH}^c)$ are precisely the $\mathcal{C}_{p,\infty}$ for all p .

The Original classification

In the above picture, a line indicates that the higher prime is in the closure of the lower one. We have more precisely:

1. The pre-image of the dense point $(0) \in \operatorname{Spec}(\mathbb{Z})$ is the single point $\mathcal{C}_{0,1}$
2. For each prime p , the pre-image of $p\mathbb{Z}$ consists of the column $\mathcal{C}_{p,n}$ and where $\mathcal{C}_{p,\infty} = \ker(q_p : \operatorname{SH}^c \rightarrow \operatorname{SH}_p^c)$.
3. $\mathcal{C}_{0,1}$ is the unique dense point in $\operatorname{Spc}(\operatorname{SH}^c)$. For each prime p and integer $1 \leq n \neq \infty$ we have the closure $\overline{\{\mathcal{C}_{p,n}\}} = \{\mathcal{C}_{p,m} : n \leq m \leq \infty\}$. The closed points of $\operatorname{Spc}(\operatorname{SH}^c)$ are precisely the $\mathcal{C}_{p,\infty}$ for all p .
4. The support of an object x is:
 - 4.1 $\operatorname{supp}(x) = \emptyset$ when $x \cong 0$
 - 4.2 $\operatorname{supp}(x) = \operatorname{Spc}(\operatorname{SH}^c)$ when $x \notin \mathcal{C}_{0,1}$
 - 4.3 $\operatorname{supp}(x) =$ a finite union of "columns" when $x \in \mathcal{C}_{0,1}$. More concretely, $\operatorname{supp}(x) =$ finite unions of $\overline{\mathcal{C}_{p,m_p}}$ where $\mathcal{C}_{p,m_p} := \{\mathcal{C}_{p,n} : m_p \leq n \leq \infty\}$ and where m_p is the "type" of p .

The Original classification

In the above picture, a line indicates that the higher prime is in the closure of the lower one. We have more precisely:

1. The pre-image of the dense point $(0) \in \text{Spec}(\mathbb{Z})$ is the single point $\mathcal{C}_{0,1}$
2. For each prime p , the pre-image of $p\mathbb{Z}$ consists of the column $\mathcal{C}_{p,n}$ and where $\mathcal{C}_{p,\infty} = \ker(q_p : \text{SH}^c \rightarrow \text{SH}_p^c)$.
3. $\mathcal{C}_{0,1}$ is the unique dense point in $\text{Spc}(\text{SH}^c)$. For each prime p and integer $1 \leq n \neq \infty$ we have the closure $\overline{\{\mathcal{C}_{p,n}\}} = \{\mathcal{C}_{p,m} : n \leq m \leq \infty\}$. The closed points of $\text{Spc}(\text{SH}^c)$ are precisely the $\mathcal{C}_{p,\infty}$ for all p .
4. The support of an object x is:
 - 4.1 $\text{supp}(x) = \emptyset$ when $x \cong 0$
 - 4.2 $\text{supp}(x) = \text{Spc}(\text{SH}^c)$ when $x \notin \mathcal{C}_{0,1}$
 - 4.3 $\text{supp}(x) =$ a finite union of "columns" when $x \in \mathcal{C}_{0,1}$. More concretely, $\text{supp}(x) =$ finite unions of $\overline{\mathcal{C}_{p,m_p}}$ where $\mathcal{C}_{p,m_p} := \{\mathcal{C}_{p,n} : m_p \leq n \leq \infty\}$ and where m_p is the "type" of p .
5. The Thomason Subsets of $\text{Spc}(\text{SH}^c)$ are
 - 5.1 the empty set and the whole space itself
 - 5.2 Arbitrary unions of columns $\overline{\mathcal{C}_{p,m_p}}$
6. A great way to think about each column is that it expresses a "chromatic refinement" between the representing spectra $\text{H}\mathbb{Q}$ and $\text{H}\mathbb{F}_p$ (ie, between the primes (0) and $p\mathbb{Z}$).

Recent work by Balmer-Sanders

1. The examples above, while providing a great conceptual framework that unifies seemingly disjoint work, are more repackaging of old theorems rather than brave new work.
2. In many of the cases in fact, computing $\mathrm{Spc}(\mathcal{K})$ is done by ALREADY knowing the tt ideals of \mathcal{K} . But we would like to do the reverse: Given a tt category \mathcal{K} compute $\mathrm{Spc}(\mathcal{K})$ from first principles, and then from that, DEDUCE the tt-ideals of \mathcal{K} .

Recent work by Balmer-Sanders

1. The examples above, while providing a great conceptual framework that unifies seemingly disjoint work, are more repackaging of old theorems rather than brave new work.
2. In many of the cases in fact, computing $\mathrm{Spc}(\mathcal{K})$ is done by ALREADY knowing the tt ideals of \mathcal{K} . But we would like to do the reverse: Given a tt category \mathcal{K} compute $\mathrm{Spc}(\mathcal{K})$ from first principles, and then from that, DEDUCE the tt-ideals of \mathcal{K} .
3. Recent work by Balmer-Sanders (2017) does just that, for the case of $\mathrm{SH}^c(G)$ for G a finite group. They describe $\mathrm{Spc}(\mathrm{SH}^c(G))$ as a set for all finite groups G , and get close to completely describing the topology. Let us say a little about $\mathrm{SH}^c(G)$, and then we can give the statement of the theorem, and present an absolutely beautiful picture after the fact.

Recent work by Balmer-Sanders

1. The examples above, while providing a great conceptual framework that unifies seemingly disjoint work, are more repackaging of old theorems rather than brave new work.
2. In many of the cases in fact, computing $\mathrm{Spc}(\mathcal{K})$ is done by ALREADY knowing the tt ideals of \mathcal{K} . But we would like to do the reverse: Given a tt category \mathcal{K} compute $\mathrm{Spc}(\mathcal{K})$ from first principles, and then from that, DEDUCE the tt-ideals of \mathcal{K} .
3. Recent work by Balmer-Sanders (2017) does just that, for the case of $\mathrm{SH}^c(G)$ for G a finite group. They describe $\mathrm{Spc}(\mathrm{SH}^c(G))$ as a set for all finite groups G , and get close to completely describing the topology. Let us say a little about $\mathrm{SH}^c(G)$, and then we can give the statement of the theorem, and present an absolutely beautiful picture after the fact.
4. Recall the comparison map we have used a few times now $\rho_{\mathcal{K}} : \mathrm{Spc}(\mathcal{K}) \rightarrow \mathrm{Spc}(\mathrm{End}_{\mathcal{K}}(1))$. In this case, the endomorphism ring is the Burnside Ring of G , $\mathrm{End}_{\mathcal{K}}(1) \cong A(G)$.

Recent work by Balmer-Sanders

1. The examples above, while providing a great conceptual framework that unifies seemingly disjoint work, are more repackaging of old theorems rather than brave new work.
2. In many of the cases in fact, computing $\mathrm{Spc}(\mathcal{K})$ is done by ALREADY knowing the tt ideals of \mathcal{K} . But we would like to do the reverse: Given a tt category \mathcal{K} compute $\mathrm{Spc}(\mathcal{K})$ from first principles, and then from that, DEDUCE the tt-ideals of \mathcal{K} .
3. Recent work by Balmer-Sanders (2017) does just that, for the case of $\mathrm{SH}^c(G)$ for G a finite group. They describe $\mathrm{Spc}(\mathrm{SH}^c(G))$ as a set for all finite groups G , and get close to completely describing the topology. Let us say a little about $\mathrm{SH}^c(G)$, and then we can give the statement of the theorem, and present an absolutely beautiful picture after the fact.
4. Recall the comparison map we have used a few times now $\rho_{\mathcal{K}} : \mathrm{Spc}(\mathcal{K}) \rightarrow \mathrm{Spc}(\mathrm{End}_{\mathcal{K}}(1))$. In this case, the endomorphism ring is the Burnside Ring of G , $\mathrm{End}_{\mathcal{K}}(1) \cong A(G)$.
5. Remember we remarked that the computation of $\mathrm{Spc}(\mathrm{SH}^c)$ provided a "refinement" between the primes at (0) and $p\mathbb{Z}$ - we get a similar refinement of the spectrum of $A(G)$ in this case as well.

$\mathrm{Spc}(\mathrm{SH}^c)$ as a set.

1. Thrm: All G -equivariant primes are obtained by pulling back non-equivariant primes via geometric fixed point functors with respect to the various subgroups $H \leq G$. Moreover, there is no redundancy, in the sense that the primes $\mathcal{P}(H, p, n) = \mathcal{P}(H', p', n')$ iff H is conjugate to H' and the chromatic primes $\mathcal{C}_{p,n} = \mathcal{C}_{p',n'}$ coincide in SH^c (where $\mathcal{P}(H, p, n) := (\Phi^H)^{-1}(\mathcal{C}_{p,n})$ are the pulled back primes in SH^c by the "geometric fixed point functor").
If $K \trianglelefteq H$ has nonzero index then $\mathcal{P}(K, p, n+1) \subseteq \mathcal{P}(H, p, n)$ for every $n \geq 1$. There is no inclusion $\mathcal{P}(K, q, n) \subseteq \mathcal{P}(H, p, m)$ unless the Chromatic primes are included, $\mathcal{C}_{q,n} \subseteq \mathcal{C}_{p,m}$ and K is conjugate to a q -subnormal subgroup of H .

$\mathrm{Spc}(\mathrm{SH}^c)$ as a set.

1. Thrm: All G -equivariant primes are obtained by pulling back non-equivariant primes via geometric fixed point functors with respect to the various subgroups $H \leq G$. Moreover, there is no redundancy, in the sense that the primes $\mathcal{P}(H, p, n) = \mathcal{P}(H', p', n')$ iff H is conjugate to H' and the chromatic primes $\mathcal{C}_{p,n} = \mathcal{C}_{p',n'}$ coincide in SH^c (where $\mathcal{P}(H, p, n) := (\Phi^H)^{-1}(\mathcal{C}_{p,n})$ are the pulled back primes in SH^c by the "geometric fixed point functor").
If $K \trianglelefteq H$ has nonzero index then $\mathcal{P}(K, p, n+1) \subseteq \mathcal{P}(H, p, n)$ for every $n \geq 1$. There is no inclusion $\mathcal{P}(K, q, n) \subseteq \mathcal{P}(H, p, m)$ unless the Chromatic primes are included, $\mathcal{C}_{q,n} \subseteq \mathcal{C}_{p,m}$ and K is conjugate to a q -subnormal subgroup of H .
2. This completely describes the topology for groups of square free order. For example, the following picture is the spectrum of $\mathrm{SH}(C_p)$.

SH but on Steroids

