

(Tensor) Triangulated Categories

What are they good for?

David Rubinstein

Why Care about these categories?

The axioms suck (not really, but still)

The definition/axioms of triangulated categories are notoriously viewed as being bad. In fact, many people are not convinced that the axioms as they currently exist will remain in their current form.

So, rather jumping right into the axioms, let us first see that they at least provide a wide range of examples. (The range of applications provides a good argument for the side of the axioms it should be noted)

Why Care about these categories?

They show up everywhere!

1. The Stable Homotopy Category of (finite) Spectra

Why Care about these categories?

They show up everywhere!

1. The Stable Homotopy Category of (finite) Spectra
2. The Homotopy Category of a commutative ring $K(R\text{-mod})$ and its variants

Why Care about these categories?

They show up everywhere!

1. The Stable Homotopy Category of (finite) Spectra
2. The Homotopy Category of a commutative ring $K(R\text{-mod})$ and its variants
3. The Derived Category of (perfect complexes of) a Ring R (or Scheme X) $D(R)$ ($D(X)$)

Why Care about these categories?

They show up everywhere!

1. The Stable Homotopy Category of (finite) Spectra
2. The Homotopy Category of a commutative ring $K(R\text{-mod})$ and its variants
3. The Derived Category of (perfect complexes of) a Ring R (or Scheme X) $D(R)$ ($D(X)$)
4. The stable Module Category $\text{StMod}(kG)$ for G a finite group (scheme), and k a field of char p that divides the order of G

Why Care about these categories?

They show up everywhere!

1. The Stable Homotopy Category of (finite) Spectra
2. The Homotopy Category of a commutative ring $K(R\text{-mod})$ and its variants
3. The Derived Category of (perfect complexes of) a Ring R (or Scheme X) $D(R)$ ($D(X)$)
4. The stable Module Category $\text{StMod}(kG)$ for G a finite group (scheme), and k a field of char p that divides the order of G
5. The Equivariant Stable Homotopy Category of G -Spectra for G a compact Lie Group

Why Care about these categories?

They show up everywhere!

1. The Stable Homotopy Category of (finite) Spectra
2. The Homotopy Category of a commutative ring $K(R\text{-mod})$ and its variants
3. The Derived Category of (perfect complexes of) a Ring R (or Scheme X) $D(R)$ ($D(X)$)
4. The stable Module Category $\text{StMod}(kG)$ for G a finite group (scheme), and k a field of char p that divides the order of G
5. The Equivariant Stable Homotopy Category of G -Spectra for G a compact Lie Group
6. The stable \mathbb{A}^1 -homotopy category and derived categories of motives (Voevodsky–Morel, Voevodsky)

Why Care about these categories?

They show up everywhere!

1. The Stable Homotopy Category of (finite) Spectra
2. The Homotopy Category of a commutative ring $K(R\text{-mod})$ and its variants
3. The Derived Category of (perfect complexes of) a Ring R (or Scheme X) $D(R)$ ($D(X)$)
4. The stable Module Category $\text{StMod}(kG)$ for G a finite group (scheme), and k a field of char p that divides the order of G
5. The Equivariant Stable Homotopy Category of G -Spectra for G a compact Lie Group
6. The stable \mathbb{A}^1 -homotopy category and derived categories of motives (Voevodsky–Morel, Voevodsky)
7. C^* - Algebras in KK -Theory




Figure: Picture stolen from Paul Balmer- depicts the broad scope of TT-Categories

Groups VS Commutative Rings

1. When one is first learning about Algebra one first learns about Group Theory and then after learns about Ring Theory.

Groups VS Commutative Rings

1. When one is first learning about Algebra one first learns about Group Theory and then after learns about Ring Theory.
2. In doing so, we find that many of the groups we studied had a richer structure that we weren't able to exploit working with just the group structure. An analogous statement is true here.

Groups VS Commutative Rings

1. When one is first learning about Algebra one first learns about Group Theory and then after learns about Ring Theory.
2. In doing so, we find that many of the groups we studied had a richer structure that we weren't able to exploit working with just the group structure. An analogous statement is true here.

$$\begin{array}{l} \text{Groups} \iff \text{Triangulated Categories} \\ \text{Commutative Rings} \iff \text{Tensor Triangulated Categories} \end{array}$$

Groups VS Commutative Rings

1. When one is first learning about Algebra one first learns about Group Theory and then after learns about Ring Theory.
2. In doing so, we find that many of the groups we studied had a richer structure that we weren't able to exploit working with just the group structure. An analogous statement is true here.

$$\begin{array}{ccc} \text{Groups} & \Longleftrightarrow & \text{Triangulated Categories} \\ \text{Commutative Rings} & \Longleftrightarrow & \text{Tensor Triangulated Categories} \end{array}$$

3. This talk should be viewed as learning about Group Theory then.

Groups VS Commutative Rings

1. When one is first learning about Algebra one first learns about Group Theory and then after learns about Ring Theory.
2. In doing so, we find that many of the groups we studied had a richer structure that we weren't able to exploit working with just the group structure. An analogous statement is true here.

$$\begin{array}{l} \text{Groups} \iff \text{Triangulated Categories} \\ \text{Commutative Rings} \iff \text{Tensor Triangulated Categories} \end{array}$$

3. This talk should be viewed as learning about Group Theory then.
4. Next talk will be analogues to Ring Theory- via the work of Balmer et, al one can push this analogy quite far through the use of the Triangulated Spectrum.

Unification

1. Since so many examples of these Triangulated Categories exist ranging from algebra/geometry (examples 2,3,4) to topology (1,5,6) to analysis (7) the vague hope is that by studying "Triangulated Categories" writ large, one can learn things about all these topics in one fell swoop.

Unification

1. Since so many examples of these Triangulated Categories exist ranging from algebra/geometry (examples 2,3,4) to topology (1,5,6) to analysis (7) the vague hope is that by studying "Triangulated Categories" writ large, one can learn things about all these topics in one fell swoop.
2. Moreover, one can ask if we can transfer ideas from one branch to another using this framework of Triangulated Categories. Beren likes to describe the image above as being a conveyor belts of sorts- you feed in data from one example, and use the conveyor belt of TT-Categories to spit out that data in another example.

Unification

1. Since so many examples of these Triangulated Categories exist ranging from algebra/geometry (examples 2,3,4) to topology (1,5,6) to analysis (7) the vague hope is that by studying "Triangulated Categories" writ large, one can learn things about all these topics in one fell swoop.
2. Moreover, one can ask if we can transfer ideas from one branch to another using this framework of Triangulated Categories. Beren likes to describe the image above as being a conveyor belts of sorts- you feed in data from one example, and use the conveyor belt of TT-Categories to spit out that data in another example.
3. For example- Can we relate the ideas of line bundles in Algebraic-Geometry and Endotrivial Modules in Representation Theory? Moreover, can we relate $\text{Spec}(R)$ for a commutative ring with Support Varieties $\mathcal{V}_G := \text{Proj}(H^*(G, k))$

Unification

1. Since so many examples of these Triangulated Categories exist ranging from algebra/geometry (examples 2,3,4) to topology (1,5,6) to analysis (7) the vague hope is that by studying "Triangulated Categories" writ large, one can learn things about all these topics in one fell swoop.
2. Moreover, one can ask if we can transfer ideas from one branch to another using this framework of Triangulated Categories. Beren likes to describe the image above as being a conveyor belts of sorts- you feed in data from one example, and use the conveyor belt of TT-Categories to spit out that data in another example.
3. For example- Can we relate the ideas of line bundles in Algebraic-Geometry and Endotrivial Modules in Representation Theory? Moreover, can we relate $\text{Spec}(R)$ for a commutative ring with Support Varieties $\mathcal{V}_G := \text{Proj}(H^*(G, k))$
4. There is a famous nilpotence theorem of Hopkins and Smith in Algebraic Topology- are there analogous nilpotent theorems in Other contexts? In Alg Topology this nilpotence theorem provides a stratification for our category, can we expect the same for other nilpotence theorems?

Oh Lord the Axioms

A *pre-triangulated category* is an additive category \mathcal{T} equipped with an auto-equivalence $\Sigma : \mathcal{T} \rightarrow \mathcal{T}$ (called *suspension*) and a class of diagrams (called *distinguished triangles*)

$a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ satisfying the axioms:

Oh Lord the Axioms

A *pre-triangulated category* is an additive category \mathcal{T} equipped with an auto-equivalence $\Sigma : \mathcal{T} \rightarrow \mathcal{T}$ (called *suspension*) and a class of diagrams (called *distinguished triangles*)

$a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ satisfying the axioms:

1. (*T0- Bookkeeping axiom*) The distinguished triangles are closed under isomorphisms and rotations; That is, a triangle is distinguished iff $b \xrightarrow{g} c \xrightarrow{h} \Sigma a \xrightarrow{-\Sigma f} \Sigma b$ is distinguished. Furthermore, For all $x \in \mathcal{T}$, the triangle $x \xrightarrow{\text{Id}} x \rightarrow 0 \rightarrow \Sigma x$ is distinguished;

Oh Lord the Axioms

A *pre-triangulated category* is an additive category \mathcal{T} equipped with an auto-equivalence $\Sigma : \mathcal{T} \rightarrow \mathcal{T}$ (called *suspension*) and a class of diagrams (called *distinguished triangles*)

$a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ satisfying the axioms:

1. (*T0- Bookkeeping axiom*) The distinguished triangles are closed under isomorphisms and rotations; That is, a triangle is distinguished iff $b \xrightarrow{g} c \xrightarrow{h} \Sigma a \xrightarrow{-\Sigma f} \Sigma b$ is distinguished. Furthermore, For all $x \in \mathcal{T}$, the triangle $x \xrightarrow{\text{Id}} x \rightarrow 0 \rightarrow \Sigma x$ is distinguished;
2. (*T1- Existence Axiom*) Every morphism $a \xrightarrow{f} b$ fits into a distinguished triangle $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$. We call this c the "Cone of f "

The Axioms

Oh Lord the Axioms

A *pre-triangulated category* is an additive category \mathcal{T} equipped with an auto-equivalence $\Sigma : \mathcal{T} \rightarrow \mathcal{T}$ (called *suspension*) and a class of diagrams (called *distinguished triangles*)

$a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ satisfying the axioms:

1. (T0- Bookkeeping axiom) The distinguished triangles are closed under isomorphisms and rotations; That is, a triangle is distinguished iff $b \xrightarrow{g} c \xrightarrow{h} \Sigma a \xrightarrow{-\Sigma f} \Sigma b$ is distinguished. Furthermore, For all $x \in \mathcal{T}$, the triangle $x \xrightarrow{\text{Id}} x \rightarrow 0 \rightarrow \Sigma x$ is distinguished;
2. (T1- Existence Axiom) Every morphism $a \xrightarrow{f} b$ fits into a distinguished triangle $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$. We call this c the "Cone of f "
3. (T2- Morphism Axiom) Given the diagram whose rows are distinguished triangles and whose left square commutes

$$\begin{array}{ccccccc}
 a & \xrightarrow{f} & b & \xrightarrow{g} & c & \xrightarrow{h} & \Sigma a \\
 \alpha \downarrow & & \beta \downarrow & & \gamma \downarrow & & \downarrow \Sigma \alpha \\
 a' & \xrightarrow{f'} & b' & \xrightarrow{g'} & c' & \xrightarrow{h'} & \Sigma a'
 \end{array}$$


there exists $\gamma : c \rightarrow c'$ such that the entire diagram commutes.

The Axioms Part 2

The dreaded Octahedral Axiom

A *Triangulated Category* is a pre-Triangulated Category that satisfies the further axiom

4. (T3)- *The Octahedral Axiom* Any two morphisms $a \xrightarrow{f} a' \xrightarrow{f'} a''$ fit into a diagram



where the dashed lines represent maps into the suspension and where the four triangles inside are distinguished.

CW-Complexes

1. In a beginning Algebraic Topology class, one studies so called CW-Complexes. Now given any morphism of $X \xrightarrow{f} Y$ of CW-complexes one can form another CW-Complex, called the "**Cone of f** ", and denoted $C(f)$, that comes with a map $Y \rightarrow C(f)$.

CW-Complexes

1. In a beginning Algebraic Topology class, one studies so called CW-Complexes. Now given any morphism of $X \xrightarrow{f} Y$ of CW-complexes one can form another CW-Complex, called the "**Cone of f** ", and denoted $C(f)$, that comes with a map $Y \rightarrow C(f)$.
2. Furthermore, for any CW-complex X one can form the **suspension of X** , denoted $\Sigma X = \mathbb{S}^1 \wedge X$.

CW-Complexes

1. In a beginning Algebraic Topology class, one studies so called CW-Complexes. Now given any morphism of $X \xrightarrow{f} Y$ of CW-complexes one can form another CW-Complex, called the "**Cone of f** ", and denoted $C(f)$, that comes with a map $Y \rightarrow C(f)$.
2. Furthermore, for any CW-complex X one can form the **suspension of X** , denoted $\Sigma X = \mathbb{S}^1 \wedge X$.
3. Finally, for any map $X \xrightarrow{f} Y$ of CW-complexes there is a natural map $C(f) \xrightarrow{h} \Sigma X$.

CW-Complexes

1. In a beginning Algebraic Topology class, one studies so called CW-Complexes. Now given any morphism of $X \xrightarrow{f} Y$ of CW-complexes one can form another CW-Complex, called the "**Cone of f** ", and denoted $C(f)$, that comes with a map $Y \rightarrow C(f)$.
2. Furthermore, for any CW-complex X one can form the **suspension of X** , denoted $\Sigma X = \mathbb{S}^1 \wedge X$.
3. Finally, for any map $X \xrightarrow{f} Y$ of CW-complexes there is a natural map $C(f) \xrightarrow{h} \Sigma X$.
4. Putting this all together, for any map of CW-complexes, we get morphisms $X \xrightarrow{f} Y \xrightarrow{g} C(f) \xrightarrow{h} \Sigma X$.

Complexes of R-Modules

1. Recall that the Category of Chain Complexes of R-Modules has objects

$$A = \cdots \rightarrow A_n \xrightarrow{d_n} A_{n-1} \xrightarrow{d_{n-1}} A_{n-2} \rightarrow \cdots$$

with A_i an R-Module and where the d_i are R-Module Homomorphisms called the differentials.

Complexes of R-Modules

1. Recall that the Category of Chain Complexes of R-Modules has objects

$$A = \cdots \rightarrow A_n \xrightarrow{d_n} A_{n-1} \xrightarrow{d_{n-1}} A_{n-2} \rightarrow \cdots$$

with A_i an R-Module and where the d_i are R-Module Homomorphisms called the differentials.

2. A morphism $f : A \rightarrow B$ of Chain Complexes is a map from $A_n \rightarrow B_n$ that makes each square with the differentials commute.

Complexes of R-Modules

1. Recall that the Category of Chain Complexes of R-Modules has objects

$$A = \cdots \rightarrow A_n \xrightarrow{d_n} A_{n-1} \xrightarrow{d_{n-1}} A_{n-2} \rightarrow \cdots$$

with A_i an R-Module and where the d_i are R-Module Homomorphisms called the differentials.

2. A morphism $f : A \rightarrow B$ of Chain Complexes is a map from $A_n \rightarrow B_n$ that makes each square with the differentials commute.
3. Given any morphism $f : A \rightarrow B$ of chain complexes one can form the chain complex, **$C(f)$ called the cone of f** , where $(C(f))_n = A_{n-1} \oplus B_n$ and whose differential $d_n : A_{n-1} \oplus B_n \rightarrow A_{n-2} \oplus B_{n-1}$ is given as $d_n = \begin{pmatrix} -d_{n-1}^A & 0 \\ -f & d_n^B \end{pmatrix}$

Complexes of R-Modules

1. Recall that the Category of Chain Complexes of R-Modules has objects

$$A = \cdots \rightarrow A_n \xrightarrow{d_n} A_{n-1} \xrightarrow{d_{n-1}} A_{n-2} \rightarrow \cdots$$

with A_i an R-Module and where the d_i are R-Module Homomorphisms called the differentials.

2. A morphism $f : A \rightarrow B$ of Chain Complexes is a map from $A_n \rightarrow B_n$ that makes each square with the differentials commute.
3. Given any morphism $f : A \rightarrow B$ of chain complexes one can form the chain complex, **$C(f)$ called the cone of f** , where $(C(f))_n = A_{n-1} \oplus B_n$ and whose differential $d_n : A_{n-1} \oplus B_n \rightarrow A_{n-2} \oplus B_{n-1}$ is given as $d_n = \begin{pmatrix} -d_{n-1}^A & 0 \\ -f & d_n^B \end{pmatrix}$
4. Now given any Chain complex A , we let ΣA be the chain complex shifted down in one dimension- that is $\Sigma A_n = A_{n-1}$.

Complexes of R-Modules

1. Recall that the Category of Chain Complexes of R-Modules has objects

$$A = \cdots \rightarrow A_n \xrightarrow{d_n} A_{n-1} \xrightarrow{d_{n-1}} A_{n-2} \rightarrow \cdots$$

with A_i an R-Module and where the d_i are R-Module Homomorphisms called the differentials.

2. A morphism $f : A \rightarrow B$ of Chain Complexes is a map from $A_n \rightarrow B_n$ that makes each square with the differentials commute.
3. Given any morphism $f : A \rightarrow B$ of chain complexes one can form the chain complex, **$C(f)$ called the cone of f** , where $(C(f))_n = A_{n-1} \oplus B_n$ and whose differential $d_n : A_{n-1} \oplus B_n \rightarrow A_{n-2} \oplus B_{n-1}$ is given as $d_n = \begin{pmatrix} -d_{n-1}^A & 0 \\ -f & d_n^B \end{pmatrix}$
4. Now given any Chain complex A , we let ΣA be the chain complex shifted down in one dimension- that is $\Sigma A_n = A_{n-1}$.
5. Then putting it all together we get a sequence of morphisms of chain complexes

$$A \xrightarrow{f} B \xrightarrow{i} C(f) \xrightarrow{p} \Sigma A$$

where i and p are the canonical inclusion and projection maps.

Building Intuition

The intuition one should have is that Triangulated Categories behave almost like an abelian category- and that Triangles are analogues of exact sequences.

Building Intuition

The intuition one should have is that Triangulated Categories behave almost like an abelian category- and that Triangles are analogues of exact sequences.

1. Given a triangle $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ we have $g \circ f = h \circ g = \Sigma f \circ h = 0$

Building Intuition

The intuition one should have is that Triangulated Categories behave almost like an abelian category- and that Triangles are analogues of exact sequences.

1. Given a triangle $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ we have $g \circ f = h \circ g = \Sigma f \circ h = 0$
2. Applying $\text{Hom}(y, -)$ to a triangle $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ gives a long exact sequence . We say $\text{Hom}(y, -)$ is a **Homological functor**

Building Intuition

The intuition one should have is that Triangulated Categories behave almost like an abelian category- and that Triangles are analogues of exact sequences.

1. Given a triangle $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ we have $g \circ f = h \circ g = \Sigma f \circ h = 0$
2. Applying $\text{Hom}(y, -)$ to a triangle $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ gives a long exact sequence . We say $\text{Hom}(y, -)$ is a **Homological functor**
3. Dually, applying $\text{Hom}(-, y)$ gives a long exact sequence with appropriate arrows flipped. We call such a functor **cohomological**.

Building Intuition

The intuition one should have is that Triangulated Categories behave almost like an abelian category- and that Triangles are analogues of exact sequences.

1. Given a triangle $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ we have $g \circ f = h \circ g = \Sigma f \circ h = 0$
2. Applying $\text{Hom}(y, -)$ to a triangle $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ gives a long exact sequence . We say $\text{Hom}(y, -)$ is a **Homological functor**
3. Dually, applying $\text{Hom}(-, y)$ gives a long exact sequence with appropriate arrows flipped. We call such a functor **cohomological**.
4. Taking (co)products of triangles gives again a distinguished triangle

Building Intuition

The intuition one should have is that Triangulated Categories behave almost like an abelian category- and that Triangles are analogues of exact sequences.

1. Given a triangle $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ we have $g \circ f = h \circ g = \Sigma f \circ h = 0$
2. Applying $\text{Hom}(y, -)$ to a triangle $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ gives a long exact sequence . We say $\text{Hom}(y, -)$ is a **Homological functor**
3. Dually, applying $\text{Hom}(-, y)$ gives a long exact sequence with appropriate arrows flipped. We call such a functor **cohomological**.
4. Taking (co)products of triangles gives again a distinguished triangle
5. If $a_1 \oplus a_2 \xrightarrow{f} b_1 \oplus b_2 \xrightarrow{g} c_1 \oplus c_2 \xrightarrow{h} \Sigma(a_1 \oplus a_2)$ is a triangle, so are $a_i \xrightarrow{f_i} b_i \xrightarrow{g_i} c_i \xrightarrow{h_i} \Sigma a_i$

More Intuition

1. "The Five Lemma of Triangles"- Suppose we are given a morphism of triangles

$$\begin{array}{ccccccc} a & \xrightarrow{f} & b & \xrightarrow{g} & c & \xrightarrow{h} & \Sigma a \\ \alpha \downarrow & & \beta \downarrow & & \gamma \downarrow & & \downarrow \Sigma \alpha \\ a' & \xrightarrow{f'} & b' & \xrightarrow{g'} & c' & \xrightarrow{h'} & \Sigma a' \end{array}$$

where α, β are isomorphisms. Then so is γ

More Intuition

1. "The Five Lemma of Triangles"- Suppose we are given a morphism of triangles

$$\begin{array}{ccccccc} a & \xrightarrow{f} & b & \xrightarrow{g} & c & \xrightarrow{h} & \Sigma a \\ \alpha \downarrow & & \beta \downarrow & & \gamma \downarrow & & \downarrow \Sigma \alpha \\ a' & \xrightarrow{f'} & b' & \xrightarrow{g'} & c' & \xrightarrow{h'} & \Sigma a' \end{array}$$

where α, β are isomorphisms. Then so is γ

2. One might imagine that given a map $a \xrightarrow{f} b$ we could complete it into 2 different triangles. The above result actually shows such a triangle is unique up to (a non-unique) isomorphism. (take $a = a', b = b', \alpha = \beta = 1$)

Last bit on Intuition

1. "Split Triangles". Suppose $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ is a triangle such that there exists a retraction of g . Then there exists an isomorphism of triangles

$$\begin{array}{ccccccc} a & \xrightarrow{f} & b & \xrightarrow{g} & c & \xrightarrow{h} & \Sigma a \\ \downarrow 1 & & \downarrow \beta & & \downarrow 1 & & \downarrow 1 \\ a & \xrightarrow{f'} & a \oplus c & \xrightarrow{g'} & c & \xrightarrow{h'} & \Sigma a' \end{array}$$

Last bit on Intuition

1. "Split Triangles". Suppose $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ is a triangle such that there exists a retraction of g . Then there exists an isomorphism of triangles

$$\begin{array}{ccccccc}
 a & \xrightarrow{f} & b & \xrightarrow{g} & c & \xrightarrow{h} & \Sigma a \\
 \downarrow 1 & & \downarrow \beta & & \downarrow 1 & & \downarrow 1 \\
 a & \xrightarrow{f'} & a \oplus c & \xrightarrow{g'} & c & \xrightarrow{h'} & \Sigma a'
 \end{array}$$

2. The dual of the above is true too- if there is a retraction of f , then the triangle splits again

Last bit on Intuition

1. "Split Triangles". Suppose $a \xrightarrow{f} b \xrightarrow{g} c \xrightarrow{h} \Sigma a$ is a triangle such that there exists a retraction of g . Then there exists an isomorphism of triangles

$$\begin{array}{ccccccc} a & \xrightarrow{f} & b & \xrightarrow{g} & c & \xrightarrow{h} & \Sigma a \\ \downarrow 1 & & \downarrow \beta & & \downarrow 1 & & \downarrow 1 \\ a & \xrightarrow{f'} & a \oplus c & \xrightarrow{g'} & c & \xrightarrow{h'} & \Sigma a' \end{array}$$

2. The dual of the above is true too- if there is a retraction of f , then the triangle splits again
3. Pullbacks and Pushouts exist. (up to a weak version- the maps out of/into them are unique but up to a non-unique isomorphism). They are often called **Homotopy Cartesian Squares**. Moreover, the completion of the triangle guaranteed in Axiom T2 can be chosen in such a way that makes the middle square Homotopy Cartesian.

Inverting Morphisms= Killing Objects

This small easy lemma below will be the key to much of the discussion after this.

Inverting Morphisms= Killing Objects

This small easy lemma below will be the key to much of the discussion after this.

1. The morphism $a \xrightarrow{f} b$ is an isomorphism iff the cone of f is isomorphic to 0. That is we have a distinguished triangle $a \xrightarrow{f} b \xrightarrow{0} 0 \xrightarrow{0} \Sigma a$

Inverting Morphisms= Killing Objects

This small easy lemma below will be the key to much of the discussion after this.

1. The morphism $a \xrightarrow{f} b$ is an isomorphism iff the cone of f is isomorphic to 0. That is we have a distinguished triangle $a \xrightarrow{f} b \xrightarrow{0} 0 \xrightarrow{0} \Sigma a$
2. Now is a good time to mention- no results above used the Octahedral Axiom. In fact, Neeman doesn't even give the 4th Axiom in his textbook on Triangulated Categories until after the sections covering the material above. His 4th axiom is actually different than the Octahedral Axiom(his involves so called mapping cones) but he eventually proves they are equivalent.
3. We will need it for the slides to come however.

Killing Cones

1. Now we get to the some of the first major results. We say a full, additive subcategory \mathcal{S} is a **triangulated subcategory** if, given $x, y \in \mathcal{S}$ the cone of any morphism between x and y is in \mathcal{S} .

Killing Cones

1. Now we get to the some of the first major results. We say a full, additive subcategory \mathcal{S} is a **triangulated subcategory** if, given $x, y \in \mathcal{S}$ the cone of any morphism between x and y is in \mathcal{S} .
2. We let $A_{\mathcal{S}}$ be the sub collection of morphisms of \mathcal{T} be given by all morphisms whose cone lies in \mathcal{S} . That is, $f \in A_{\mathcal{S}}$ iff there is some x, y such that $x \xrightarrow{f} y \xrightarrow{g} z \xrightarrow{h} \Sigma x$ is a triangle with $z \in \mathcal{S}$. Then note,

Killing Cones

1. Now we get to the some of the first major results. We say a full, additive subcategory \mathcal{S} is a **triangulated subcategory** if, given $x, y \in \mathcal{S}$ the cone of any morphism between x and y is in \mathcal{S} .
2. We let $A_{\mathcal{S}}$ be the sub collection of morphisms of \mathcal{T} be given by all morphisms whose cone lies in \mathcal{S} . That is, $f \in A_{\mathcal{S}}$ iff there is some x, y such that $x \xrightarrow{f} y \xrightarrow{g} z \xrightarrow{h} \Sigma x$ is a triangle with $z \in \mathcal{S}$. Then note,
 - 2.1 Every isomorphism f is in $A_{\mathcal{S}}$.
 - 2.2 While we can consider this system of morphisms for any triangulated subcategory, we will be most interested in them for the following type.

Killing Cones

1. Now we get to the some of the first major results. We say a full, additive subcategory \mathcal{S} is a **triangulated subcategory** if, given $x, y \in \mathcal{S}$ the cone of any morphism between x and y is in \mathcal{S} .
2. We let $A_{\mathcal{S}}$ be the sub collection of morphisms of \mathcal{T} be given by all morphisms whose cone lies in \mathcal{S} . That is, $f \in A_{\mathcal{S}}$ iff there is some x, y such that $x \xrightarrow{f} y \xrightarrow{g} z \xrightarrow{h} \Sigma x$ is a triangle with $z \in \mathcal{S}$. Then note,
 - 2.1 Every isomorphism f is in $A_{\mathcal{S}}$.
 - 2.2 While we can consider this system of morphisms for any triangulated subcategory, we will be most interested in them for the following type.
3. A **triangulated subcategory** \mathcal{D} is called **Thick** if it is closed under direct summands.

Killing Cones

1. Now we get to some of the first major results. We say a full, additive subcategory \mathcal{S} is a **triangulated subcategory** if, given $x, y \in \mathcal{S}$ the cone of any morphism between x and y is in \mathcal{S} .
2. We let $\mathbf{A}_{\mathcal{S}}$ be the sub collection of morphisms of \mathcal{T} be given by all morphisms whose cone lies in \mathcal{S} . That is, $f \in \mathbf{A}_{\mathcal{S}}$ iff there is some x, y such that $x \xrightarrow{f} y \xrightarrow{g} z \xrightarrow{h} \Sigma x$ is a triangle with $z \in \mathcal{S}$. Then note,
 - 2.1 Every isomorphism f is in $\mathbf{A}_{\mathcal{S}}$.
 - 2.2 While we can consider this system of morphisms for any triangulated subcategory, we will be most interested in them for the following type.
3. A **triangulated subcategory** \mathcal{D} is called **Thick** if it is closed under direct summands.
 - 3.1 One should think of thick subcategories as being the analogue of normal subgroups. In group Theory, we have a correspondence between normal subgroups and kernels of group homomorphisms. Is something like that true here?

Triangulated Functors

Continuing the analogy above, we want to define something like a group homomorphism. Our version is the following: **A triangulated functor** is

1. An additive functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between triangulated categories

Triangulated Functors

Continuing the analogy above, we want to define something like a group homomorphism. Our version is the following: **A triangulated functor** is

1. An additive functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between triangulated categories
2. That commutes with suspensions, and
3. Takes triangles in \mathcal{C} to triangles in \mathcal{D} . Furthermore,

Triangulated Functors

Continuing the analogy above, we want to define something like a group homomorphism. Our version is the following: A **triangulated functor** is

1. An additive functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between triangulated categories
2. That commutes with suspensions, and
3. Takes triangles in \mathcal{C} to triangles in \mathcal{D} . Furthermore,
4. The **kernal of a triangulated functor** is the class of objects in \mathcal{C} that get mapped to an object isomorphic to 0 in \mathcal{D} .

Triangulated Functors

Continuing the analogy above, we want to define something like a group homomorphism. Our version is the following: A **triangulated functor** is

1. An additive functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between triangulated categories
2. That commutes with suspensions, and
3. Takes triangles in \mathcal{C} to triangles in \mathcal{D} . Furthermore,
4. The **kernal of a triangulated functor** is the class of objects in \mathcal{C} that get mapped to an object isomorphic to 0 in \mathcal{D} .
5. It should come as no surprise that
 - 5.1 $\ker(F)$ is a thick subcategory

Triangulated Functors

Continuing the analogy above, we want to define something like a group homomorphism. Our version is the following: A **triangulated functor** is

1. An additive functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between triangulated categories
2. That commutes with suspensions, and
3. Takes triangles in \mathcal{C} to triangles in \mathcal{D} . Furthermore,
4. The **kernal of a triangulated functor** is the class of objects in \mathcal{C} that get mapped to an object isomorphic to 0 in \mathcal{D} .
5. It should come as no surprise that
 - 5.1 $\ker(F)$ is a thick subcategory
 - 5.2 Is the converse true? Are all thick subcategories kernels of a triangulated functor?

Main Thrm

Let \mathcal{D} be a thick subcategory of a triangulated category \mathcal{T} . Then there exists a triangulated category, denoted \mathcal{T}/\mathcal{D} and a universal triangulated functor $Q : \mathcal{T} \rightarrow \mathcal{T}/\mathcal{D}$ such that $\ker(Q) = \mathcal{D}$. This pair is universal in the sense that if there is a triangulated category \mathcal{F} and functor $G : \mathcal{T} \rightarrow \mathcal{F}$ with $\mathcal{D} \subseteq \ker(G)$, then G factors uniquely as $\mathcal{T} \xrightarrow{Q} \mathcal{T}/\mathcal{D} \xrightarrow{\bar{G}} \mathcal{F}$.
Indeed

Main Thrm

Let \mathcal{D} be a thick subcategory of a triangulated category \mathcal{T} . Then there exists a triangulated category, denoted \mathcal{T}/\mathcal{D} and a universal triangulated functor $Q : \mathcal{T} \rightarrow \mathcal{T}/\mathcal{D}$ such that $\ker(Q) = \mathcal{D}$. This pair is universal in the sense that if there is a triangulated category \mathcal{F} and functor $G : \mathcal{T} \rightarrow \mathcal{F}$ with $\mathcal{D} \subseteq \ker(G)$, then G factors uniquely as $\mathcal{T} \xrightarrow{Q} \mathcal{T}/\mathcal{D} \xrightarrow{\bar{G}} \mathcal{F}$.
Indeed

1. $\text{Obj}(\mathcal{T}/\mathcal{D}) = \text{Obj}(\mathcal{T})$ and Q is just the identity on Objects.

Main Thrm

Let \mathcal{D} be a thick subcategory of a triangulated category \mathcal{T} . Then there exists a triangulated category, denoted \mathcal{T}/\mathcal{D} and a universal triangulated functor $Q : \mathcal{T} \rightarrow \mathcal{T}/\mathcal{D}$ such that $\ker(Q) = \mathcal{D}$. This pair is universal in the sense that if there is a triangulated category \mathcal{F} and functor $G : \mathcal{T} \rightarrow \mathcal{F}$ with $\mathcal{D} \subseteq \ker(G)$, then G factors uniquely as $\mathcal{T} \xrightarrow{Q} \mathcal{T}/\mathcal{D} \xrightarrow{\bar{G}} \mathcal{F}$.
Indeed

1. $\text{Obj}(\mathcal{T}/\mathcal{D}) = \text{Obj}(\mathcal{T})$ and Q is just the identity on Objects.
2. The tricky part is defining Q on morphisms. The way one does this is by formally inverting all morphisms whose cone lies in \mathcal{D} . That is, we formally invert those morphisms in $A_{\mathcal{D}}$. This procedure is called "**calculus of fractions**" and it is a very gross procedure.

Main Thrm

Let \mathcal{D} be a thick subcategory of a triangulated category \mathcal{T} . Then there exists a triangulated category, denoted \mathcal{T}/\mathcal{D} and a universal triangulated functor $Q : \mathcal{T} \rightarrow \mathcal{T}/\mathcal{D}$ such that $\ker(Q) = \mathcal{D}$. This pair is universal in the sense that if there is a triangulated category \mathcal{F} and functor $G : \mathcal{T} \rightarrow \mathcal{F}$ with $\mathcal{D} \subseteq \ker(G)$, then G factors uniquely as $\mathcal{T} \xrightarrow{Q} \mathcal{T}/\mathcal{D} \xrightarrow{\bar{G}} \mathcal{F}$. Indeed

1. $\text{Obj}(\mathcal{T}/\mathcal{D}) = \text{Obj}(\mathcal{T})$ and Q is just the identity on Objects.
2. The tricky part is defining Q on morphisms. The way one does this is by formally inverting all morphisms whose cone lies in \mathcal{D} . That is, we formally invert those morphisms in $A_{\mathcal{D}}$. This procedure is called "**calculus of fractions**" and it is a very gross procedure.
3. Since we are inverting morphisms whose cone lands in \mathcal{D} , one can show that every object of \mathcal{D} is isomorphic to 0 in \mathcal{T}/\mathcal{D} .

Main Thrm

Let \mathcal{D} be a thick subcategory of a triangulated category \mathcal{T} . Then there exists a triangulated category, denoted \mathcal{T}/\mathcal{D} and a universal triangulated functor $Q : \mathcal{T} \rightarrow \mathcal{T}/\mathcal{D}$ such that $\ker(Q) = \mathcal{D}$. This pair is universal in the sense that if there is a triangulated category \mathcal{F} and functor $G : \mathcal{T} \rightarrow \mathcal{F}$ with $\mathcal{D} \subseteq \ker(G)$, then G factors uniquely as $\mathcal{T} \xrightarrow{Q} \mathcal{T}/\mathcal{D} \xrightarrow{\bar{G}} \mathcal{F}$. Indeed

1. $\text{Obj}(\mathcal{T}/\mathcal{D}) = \text{Obj}(\mathcal{T})$ and Q is just the identity on Objects.
2. The tricky part is defining Q on morphisms. The way one does this is by formally inverting all morphisms whose cone lies in \mathcal{D} . That is, we formally invert those morphisms in $\mathcal{A}_{\mathcal{D}}$. This procedure is called "**calculus of fractions**" and it is a very gross procedure.
3. Since we are inverting morphisms whose cone lands in \mathcal{D} , one can show that every object of \mathcal{D} is isomorphic to 0 in \mathcal{T}/\mathcal{D} .
4. The process is very gross and one loses control over the morphisms in the localized category pretty easily. In fact, the collection of morphisms will in general be a class (and not a set) in the localized category. We can get around this in some sneaky ways however- hopefully we can discuss this soon.
5. Side note- The way in which one formally inverts morphisms is just a massive generalization of how one localizes a ring.

Derived Category of a Ring

An example of the above Thrm

Let us show this powerful theorem in action. Let R be a ring and let $\mathcal{T} = K(R\text{-Mod})$ denote the homotopy category of chain complexes of R -modules. This is a classic example of a triangulated category. Now consider the subcategory of so called **Acyclic** objects, defined as $\text{Acy}(\mathcal{T}) = \{A \in \mathcal{T} : H^i(A) = 0 \text{ for all } i\}$. That is, the collection of all chain complexes with trivial Homology. Then

An example of the above Thrm

Let us show this powerful theorem in action. Let R be a ring and let $\mathcal{T} = K(R\text{-Mod})$ denote the homotopy category of chain complexes of R -modules. This is a classic example of a triangulated category. Now consider the subcategory of so called **Acyclic** objects, defined as $\text{Acy}(\mathcal{T}) = \{A \in \mathcal{T} : H^i(A) = 0 \text{ for all } i\}$. That is, the collection of all chain complexes with trivial Homology. Then

1. $\text{Acy}(\mathcal{T})$ is a thick subcategory of \mathcal{T}

Derived Category of a Ring

An example of the above Thrm

Let us show this powerful theorem in action. Let R be a ring and let $\mathcal{T} = K(R\text{-Mod})$ denote the homotopy category of chain complexes of R -modules. This is a classic example of a triangulated category. Now consider the subcategory of so called **Acyclic** objects, defined as $\text{Acy}(\mathcal{T}) = \{A \in \mathcal{T} : H^i(A) = 0 \text{ for all } i\}$. That is, the collection of all chain complexes with trivial Homology. Then

1. $\text{Acy}(\mathcal{T})$ is a thick subcategory of \mathcal{T}
2. The corresponding quotient $\mathcal{T} / \text{Acy}(\mathcal{T})$ is a triangulated category **called the Derived Category and denoted $D(R)$** .

Derived Category of a Ring

An example of the above Thrm

Let us show this powerful theorem in action. Let R be a ring and let $\mathcal{T} = K(R\text{-Mod})$ denote the homotopy category of chain complexes of R -modules. This is a classic example of a triangulated category. Now consider the subcategory of so called **Acyclic** objects, defined as $\text{Acy}(\mathcal{T}) = \{A \in \mathcal{T} : H^i(A) = 0 \text{ for all } i\}$. That is, the collection of all chain complexes with trivial Homology. Then

1. $\text{Acy}(\mathcal{T})$ is a thick subcategory of \mathcal{T}
2. The corresponding quotient $\mathcal{T} / \text{Acy}(\mathcal{T})$ is a triangulated category **called the Derived Category and denoted $D(R)$** .
3. The collection of morphisms $A_{\text{Acy}(\mathcal{T})}$ is the collection of all **"quasi-isomorphisms"**-morphisms that are invertible under every Homology functor. Hence, in the derived category, all quasi-isomorphisms are invertible.

Derived Category of a Ring

An example of the above Thrm

Let us show this powerful theorem in action. Let R be a ring and let $\mathcal{T} = K(R\text{-Mod})$ denote the homotopy category of chain complexes of R -modules. This is a classic example of a triangulated category. Now consider the subcategory of so called **Acyctic** objects, defined as $\text{Acy}(\mathcal{T}) = \{A \in \mathcal{T} : H^i(A) = 0 \text{ for all } i\}$. That is, the collection of all chain complexes with trivial Homology. Then

1. $\text{Acy}(\mathcal{T})$ is a thick subcategory of \mathcal{T}
2. The corresponding quotient $\mathcal{T} / \text{Acy}(\mathcal{T})$ is a triangulated category **called the Derived Category and denoted $D(R)$** .
3. The collection of morphisms $A_{\text{Acy}(\mathcal{T})}$ is the collection of all **"quasi-isomorphisms"**-morphisms that are invertible under every Homology functor. Hence, in the derived category, all quasi-isomorphisms are invertible.
4. Much of the historical motivation of the work above was to find a category in which these quasi-isomorphisms actually are isomorphisms- the above does just that.

Similar examples

1. Often, we are focused only on chain complexes that are either bounded above, below, or just plain old bounded. The collection of all those form triangulated categories in their own right, denoted as $K^+(\mathcal{R} - \text{Mod})$, $K^-(\mathcal{R} - \text{Mod})$, $K^b(\mathcal{R} - \text{Mod})$. respectively.

Similar examples

1. Often, we are focused only on chain complexes that are either bounded above, below, or just plain old bounded. The collection of all those form triangulated categories in their own right, denoted as $K^+(R - \text{Mod})$, $K^-(R - \text{Mod})$, $K^b(R - \text{Mod})$. respectively.
2. One then looks at the acyclic objects in each of these categories and again forms the Verdier quotient for all of these.
3. Alternatively, recall that $D(R)$ has the same objects as $K(R - \text{Mod})$, so we could talk about bounded (above/below) complexes in $D(R)$ - again denoted $D^+(R)$, $D^-(R)$, $D^b(R)$.

Similar examples

1. Often, we are focused only on chain complexes that are either bounded above, below, or just plain old bounded. The collection of all those form triangulated categories in their own right, denoted as $K^+(R - \text{Mod})$, $K^-(R - \text{Mod})$, $K^b(R - \text{Mod})$. respectively.
2. One then looks at the acyclic objects in each of these categories and again forms the Verdier quotient for all of these.
3. Alternatively, recall that $D(R)$ has the same objects as $K(R - \text{Mod})$, so we could talk about bounded (above/below) complexes in $D(R)$ - again denoted $D^+(R)$, $D^-(R)$, $D^b(R)$.
4. It is a non trivial fact that these are the verdier quotients of the above. That is
 - 4.1 $D^+(R) \cong K^+(R - \text{Mod}) / \text{Acy}(K^+(R - \text{Mod}))$
 - 4.2 $D^-(R) \cong K^-(R - \text{Mod}) / \text{Acy}(K^-(R - \text{Mod}))$
 - 4.3 $D^b(R) \cong K^b(R - \text{Mod}) / \text{Acy}(K^b(R - \text{Mod}))$

Ext Groups

1. Recall that one of the major tools in group cohomology theory are the so called Ext Groups. These are the Right Derived Functors of Hom and one goes through a lot of work showing that they "fix" the non-exactness of Hom (that is, for any pair of morphisms $A \rightarrow B \rightarrow C$, and any $R\text{-Mod } X$, there is a LES $0 \rightarrow \text{Hom}(X, A) \rightarrow \text{Hom}(X, B) \rightarrow \text{Hom}(X, C) \rightarrow \text{Ext}^1(X, A) \rightarrow \dots$)

Ext Groups

1. Recall that one of the major tools in group cohomology theory are the so called Ext Groups. These are the Right Derived Functors of Hom and one goes through a lot of work showing that they "fix" the non-exactness of Hom (that is, for any pair of morphisms $A \rightarrow B \rightarrow C$, and any $R\text{-Mod } X$, there is a LES $0 \rightarrow \text{Hom}(X, A) \rightarrow \text{Hom}(X, B) \rightarrow \text{Hom}(X, C) \rightarrow \text{Ext}^1(X, A) \rightarrow \dots$)
2. Now in the Derived Category, one makes the simple definition that the i th Ext group is $\text{Ext}^i(A, B) := \text{Hom}_{\mathcal{D}(R)}(\Sigma^{-i} A, B)$.

Ext Groups

1. Recall that one of the major tools in group cohomology theory are the so called Ext Groups. These are the Right Derived Functors of Hom and one goes through a lot of work showing that they "fix" the non-exactness of Hom (that is, for any pair of morphisms $A \rightarrow B \rightarrow C$, and any $R\text{-Mod } X$, there is a LES $0 \rightarrow \text{Hom}(X, A) \rightarrow \text{Hom}(X, B) \rightarrow \text{Hom}(X, C) \rightarrow \text{Ext}^1(X, A) \rightarrow \dots$)
2. Now in the Derived Category, one makes the simple definition that the i th Ext group is $\text{Ext}^i(A, B) := \text{Hom}_{D(R)}(\Sigma^{-i} A, B)$.
3. Ok, but given any two R -modules, A and B , you can consider them as chain complexes concentrated in degree 0 in the derived category- so one can ask, if you apply the above definition of Ext to these modules A and B , do you get an isomorphic result to the classic Ext group?

Quick Applications of Derived Category

Ext Groups

1. Recall that one of the major tools in group cohomology theory are the so called Ext Groups. These are the Right Derived Functors of Hom and one goes through a lot of work showing that they "fix" the non-exactness of Hom (that is, for any pair of morphisms $A \rightarrow B \rightarrow C$, and any R -Mod X , there is a LES $0 \rightarrow \text{Hom}(X, A) \rightarrow \text{Hom}(X, B) \rightarrow \text{Hom}(X, C) \rightarrow \text{Ext}^1(X, A) \rightarrow \dots$)
2. Now in the Derived Category, one makes the simple definition that the i th Ext group is $\text{Ext}^i(A, B) := \text{Hom}_{D(R)}(\Sigma^{-i}A, B)$.
3. Ok, but given any two R -modules, A and B , you can consider them as chain complexes concentrated in degree 0 in the derived category- so one can ask, if you apply the above definition of Ext to these modules A and B , do you get an isomorphic result to the classic Ext group?
4. YUP!!! Moreover, recall that $\text{Hom}_{D(R)}(X, -), \text{Hom}_{D(R)}(-, Y)$ are Homological/Cohomological, so they automatically turn any triangle into a LES. In particular, if $0 \rightarrow A \rightarrow A' \rightarrow A'' \rightarrow 0$ is a SES, then this gives an exact triangle in $D(R)$. So one recovers the LES expected from Ext by applying $\text{Hom}_{D(R)}(\Sigma^{-1}X, -)$ to this triangle. Moreover, this shows one can construct these sequences even without enough injective/projectives! (A similar Statement is true for Tor as well)

An Eye towards Bausfield Localization 1

Hints of a relationship to Modular Representation Theory

The talk has been dominated by Algebraic Geometric discussion so far. However, I mentioned that there are connections to representation theory and homotopy theory. Let us give a taste of those connections now. First, for Representation Theory-

Hints of a relationship to Modular Representation Theory

The talk has been dominated by Algebraic Geometric discussion so far. However, I mentioned that there are connections to representation theory and homotopy theory. Let us give a taste of those connections now. First, for Representation Theory-

1. Recall the category **$\text{Stab}(\mathbf{k}G)$** is the **stable module category** whose objects are $\mathbf{k}G$ -Modules, and where you kill the morphisms that factor through a projective module. This is a triangulated category, whose suspension functor Ω is given as the kernel of the projective cover.

Hints of a relationship to Modular Representation Theory

The talk has been dominated by Algebraic Geometric discussion so far. However, I mentioned that there are connections to representation theory and homotopy theory. Let us give a taste of those connections now. First, for Representation Theory-

1. Recall the category **$\text{Stab}(\mathbf{kG})$** is the **stable module category** whose objects are \mathbf{kG} -Modules, and where you kill the morphisms that factor through a projective module. This is a triangulated category, whose suspension functor Ω is given as the kernel of the projective cover.
2. Sitting inside $\text{Stab}(\mathbf{kG})$ is the **triangulated category $\text{stab}(\mathbf{kG})$** consisting of all finite dimensional \mathbf{kG} -Modules.

An Eye towards Bausfield Localization 1

Hints of a relationship to Modular Representation Theory

The talk has been dominated by Algebraic Geometric discussion so far. However, I mentioned that there are connections to representation theory and homotopy theory. Let us give a taste of those connections now. First, for Representation Theory-

1. Recall the category **$\text{Stab}(kG)$** is the **stable module category** whose objects are kG -Modules, and where you kill the morphisms that factor through a projective module. This is a triangulated category, whose suspension functor Ω is given as the kernel of the projective cover.
2. Sitting inside $\text{Stab}(kG)$ is the **triangulated category $\text{stab}(kG)$** consisting of all finite dimensional kG -Modules.
3. As a side note - Interestingly enough, one would probably think the relationship of $\text{stab}(kG)$ to $\text{Stab}(kG)$ should be the same as the relationship of $D^b(R)$ to $D(R)$. This is actually incorrect in a precise sense- and hints at a possible "failure" for $D(R)$.

An Eye towards Bausfield Localization 1

Hints of a relationship to Modular Representation Theory

The talk has been dominated by Algebraic Geometric discussion so far. However, I mentioned that there are connections to representation theory and homotopy theory. Let us give a taste of those connections now. First, for Representation Theory-

1. Recall the category **Stab(kG) is the stable module category** whose objects are kG -Modules, and where you kill the morphisms that factor through a projective module. This is a triangulated category, whose suspension functor Ω is given as the kernel of the projective cover.
2. Sitting inside $\text{Stab}(kG)$ is the **triangulated category $\text{stab}(kG)$** consisting of all finite dimensional kG -Modules.
3. As a side note - Interestingly enough, one would probably think the relationship of $\text{stab}(kG)$ to $\text{Stab}(kG)$ should be the same as the relationship of $D^b(R)$ to $D(R)$. This is actually incorrect in a precise sense- and hints at a possible "failure" for $D(R)$.
4. Now on the face of it, this seems disjoint from the previous topic. However, it is a theorem of Rickard (1989) that one can realize $\text{stab}(kG)$ as a Verdier Quotient of a derived Category, namely

$$\text{stab}(kG) \cong D^b(kG) / K^b(\mathbf{P}\text{-Mod}).$$

where **P-Mod is the full subcategory of projective modules**. In fact this is true for any self injective algebra.

Stable Homotopy Theory

The last example I wanted to mention is how this plays out in Stable Homotopy Theory. The theory is actually much nicer in many cases in this context. In this setting, we are more interested in **Localization Functors**.

Stable Homotopy Theory

The last example I wanted to mention is how this plays out in Stable Homotopy Theory. The theory is actually much nicer in many cases in this context. In this setting, we are more interested in **Localization Functors**.

1. An exact (triangulated) functor $L : \mathcal{T} \rightarrow \mathcal{T}$ is called a **Localization functor** if there is a natural transformation $\eta : 1 \rightarrow L$ such that $L(\eta(X)) = \eta(LX)$ for all X .

Stable Homotopy Theory

The last example I wanted to mention is how this plays out in Stable Homotopy Theory. The theory is actually much nicer in many cases in this context. In this setting, we are more interested in **Localization Functors**.

1. An exact (triangulated) functor $L : \mathcal{T} \rightarrow \mathcal{T}$ is called a **Localization functor** if there is a natural transformation $\eta : 1 \rightarrow L$ such that $L(\eta(X)) = \eta(LX)$ for all X .
2. Given a Localization functor we define the **subcategory of L-Acyclic** objects as the kernel of L and **L-Local objects** as the Essential Image of L , denoted L^\perp .

Stable Homotopy Theory

The last example I wanted to mention is how this plays out in Stable Homotopy Theory. The theory is actually much nicer in many cases in this context. In this setting, we are more interested in **Localization Functors**.

1. An exact (triangulated) functor $L : \mathcal{T} \rightarrow \mathcal{T}$ is called a **Localization functor** if there is a natural transformation $\eta : 1 \rightarrow L$ such that $L(\eta(X)) = \eta(LX)$ for all X .
2. Given a Localization functor we define the subcategory of **L-Acyclic** objects as the kernel of L and **L-Local objects** as the Essential Image of L , denoted L^\perp .
3. It is a Theorem that L induces an equivalence of Categories $\mathcal{T}/\ker(L) \cong L^\perp$. More specifically, one realizes the Localization Functor as the composite of a Verdier Quotient and a fully faithful right adjoint.

Stable Homotopy Theory

The last example I wanted to mention is how this plays out in Stable Homotopy Theory. The theory is actually much nicer in many cases in this context. In this setting, we are more interested in **Localization Functors**.

1. An exact (triangulated) functor $L : \mathcal{T} \rightarrow \mathcal{T}$ is called a **Localization functor** if there is a natural transformation $\eta : 1 \rightarrow L$ such that $L(\eta(X)) = \eta(LX)$ for all X .
2. Given a Localization functor we define the subcategory of **L-Acyclic** objects as the kernel of L and **L-Local objects** as the Essential Image of L , denoted L^\perp .
3. It is a Theorem that L induces an equivalence of Categories $\mathcal{T}/\ker(L) \cong L^\perp$. More specifically, one realizes the Localization Functor as the composite of a Verdier Quotient and a fully faithful right adjoint.
4. This fully faithful right adjoint realizes the quotient as a subcategory of \mathcal{T} allowing one to avoid the messy Calculus of Fractions, and Set-Theoretic Issues associated with the construction of Verdier Quotients. (An interesting fact is that all examples of Verdier Quotients given above are actually Bausfield Localizations!)

Stable Homotopy Theory

The last example I wanted to mention is how this plays out in Stable Homotopy Theory. The theory is actually much nicer in many cases in this context. In this setting, we are more interested in **Localization Functors**.

1. An exact (triangulated) functor $L : \mathcal{T} \rightarrow \mathcal{T}$ is called a **Localization functor** if there is a natural transformation $\eta : 1 \rightarrow L$ such that $L(\eta(X)) = \eta(LX)$ for all X .
2. Given a Localization functor we define the subcategory of **L-Acyclic** objects as the kernel of L and **L-Local objects** as the Essential Image of L , denoted L^\perp .
3. It is a Theorem that L induces an equivalence of Categories $\mathcal{T}/\ker(L) \cong L^\perp$. More specifically, one realizes the Localization Functor as the composite of a Verdier Quotient and a fully faithful right adjoint.
4. This fully faithful right adjoint realizes the quotient as a subcategory of \mathcal{T} allowing one to avoid the messy Calculus of Fractions, and Set-Theoretic Issues associated with the construction of Verdier Quotients. (An interesting fact is that all examples of Verdier Quotients given above are actually Bausfield Localizations!)
5. This is the type of Localization one does in SH. It was Bausfield who realized you could do this game for any Homology Functor out of SH (that is, for any Homology functor with kernel S , there is a Localization functor with that kernel too)

What about the Tensor

1. All of the above has been about so called Triangulated Categories- and one can get pretty far working just in that world.
2. However, most of the examples "in nature," and indeed all of the examples given in this talk have an added "tensor product" structure on them.
3. It is when this tensor structure is added to the mix that things get really interesting- and including the tensor opens the door to so called "Tensor Triangulated Geometry", and with that a world of unification.