

An Introduction to Stability and Banach Algebras

Ryan Pugh

February 11, 2021

Toeplitz and Hankel Matrices

Toeplitz Matrices

A Toeplitz matrix is a matrix (usually infinite but sometimes finite) that is constant along the diagonals:

$$A = \begin{pmatrix} a_0 & a_{-1} & a_{-2} & a_{-3} & \dots \\ a_1 & a_0 & a_{-1} & a_{-2} & \dots \\ a_2 & a_1 & a_0 & a_{-1} & \dots \\ a_3 & a_2 & a_1 & a_0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

We think of them as operators acting on the space $\ell^2(\mathbb{N})$, i.e, the space of sequences indexed by the natural numbers.

Toeplitz Matrices

A Toeplitz matrix is a matrix (usually infinite but sometimes finite) that is constant along the diagonals:

$$A = \begin{pmatrix} a_0 & a_{-1} & a_{-2} & a_{-3} & \dots \\ a_1 & a_0 & a_{-1} & a_{-2} & \dots \\ a_2 & a_1 & a_0 & a_{-1} & \dots \\ a_3 & a_2 & a_1 & a_0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

We think of them as operators acting on the space $\ell^2(\mathbb{N})$, i.e, the space of sequences indexed by the natural numbers.

One question we can ask about them is: When are these matrices bounded (= continuous)?

Toeplitz Matrices

$$A = \begin{pmatrix} a_0 & a_{-1} & a_{-2} & a_{-3} & \dots \\ a_1 & a_0 & a_{-1} & a_{-2} & \dots \\ a_2 & a_1 & a_0 & a_{-1} & \dots \\ a_3 & a_2 & a_1 & a_0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

Theorem

The Toeplitz matrix above generates a bounded operator on ℓ^2 if and only if there exists a function $a \in L^\infty(\mathbb{T})$ whose sequence of Fourier coefficients is the sequence $\{a_n\}_{n \in \mathbb{Z}}$

Toeplitz Matrices

$$A = \begin{pmatrix} a_0 & a_{-1} & a_{-2} & a_{-3} & \dots \\ a_1 & a_0 & a_{-1} & a_{-2} & \dots \\ a_2 & a_1 & a_0 & a_{-1} & \dots \\ a_3 & a_2 & a_1 & a_0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

Theorem

The Toeplitz matrix above generates a bounded operator on ℓ^2 if and only if there exists a function $a \in L^\infty(\mathbb{T})$ whose sequence of Fourier coefficients is the sequence $\{a_n\}_{n \in \mathbb{Z}}$

This function $a \in L^\infty$ is called the *symbol* for the matrix and we refer to the matrix by $T(a)$.

Whenever we're introduced to any new object, there are lots of good questions we can ask about them. We might ask

1. If the matrix is bounded, what is its norm? (this is, pretty satisfyingly, equal to the L^∞ norm of its symbol)

Whenever we're introduced to any new object, there are lots of good questions we can ask about them. We might ask

1. If the matrix is bounded, what is its norm? (this is, pretty satisfyingly, equal to the L^∞ norm of its symbol)
2. What can we say about the spectrum? (hard question)

Whenever we're introduced to any new object, there are lots of good questions we can ask about them. We might ask

1. If the matrix is bounded, what is its norm? (this is, pretty satisfyingly, equal to the L^∞ norm of its symbol)
2. What can we say about the spectrum? (hard question)
3. In what context do they arise?

Hankel Matrices

Closely related to Toeplitz matrices are these things called Hankel matrices. They look like this:

Hankel Matrices

Closely related to Toeplitz matrices are these things called Hankel matrices. They look like this:

$$A = \begin{pmatrix} a_1 & a_2 & a_3 & \dots \\ a_2 & a_3 & a_4 & \dots \\ a_3 & a_4 & a_5 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \tilde{A} = \begin{pmatrix} a_{-1} & a_{-2} & a_{-3} & \dots \\ a_{-2} & a_{-3} & a_{-4} & \dots \\ a_{-3} & a_{-4} & a_{-5} & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

Hankel Matrices

Closely related to Toeplitz matrices are these things called Hankel matrices. They look like this:

$$A = \begin{pmatrix} a_1 & a_2 & a_3 & \dots \\ a_2 & a_3 & a_4 & \dots \\ a_3 & a_4 & a_5 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \tilde{A} = \begin{pmatrix} a_{-1} & a_{-2} & a_{-3} & \dots \\ a_{-2} & a_{-3} & a_{-4} & \dots \\ a_{-3} & a_{-4} & a_{-5} & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

Similar to Toeplitz matrices, we have the following fact:

Theorem

The matrix A (resp. \tilde{A}) generates a bounded operator on ℓ^2 if and only if there exists a function $b \in L^\infty$ such that $b_n = a_n$ (resp. $b_{-n} = a_{-n}$) for all $n \geq 1$.

Connecting the Two

There is a fact that relates Topelitz and Hankel matrices! It is seemingly simple, yet very powerful when trying to prove statements.

Connecting the Two

There is a fact that relates Topelitz and Hankel matrices! It is seemingly simple, yet very powerful when trying to prove statements.

Theorem

If $a, b \in L^\infty$, then

$$T(ab) = T(a)T(b) + H(a)H(\tilde{b}).$$

Connecting the Two

There is a fact that relates Toeplitz and Hankel matrices! It is seemingly simple, yet very powerful when trying to prove statements.

Theorem

If $a, b \in L^\infty$, then

$$T(ab) = T(a)T(b) + H(a)H(\tilde{b}).$$

Worth noting is that if we further require a and b to be continuous, then we can actually conclude that the Hankel matrices generated by them are *compact*. This is useful for several reasons.

Some uses for the last theorem

Some uses for the last theorem

(Recall?) Being Fredholm is equivalent to being invertible modulo compact operators.

Some uses for the last theorem

(Recall?) Being Fredholm is equivalent to being invertible modulo compact operators. Knowing that

$$T(ab) = T(a)T(b) + H(a)H(\tilde{b})$$

i.e.,

$$T(a)T(b) = T(ab) - H(a)H(\tilde{b}).$$

can tell us things about the Fredholmness of $T(a)$ or $T(b)$ if we know things about a and b .

Some uses for the last theorem

(Recall?) Being Fredholm is equivalent to being invertible modulo compact operators. Knowing that

$$T(ab) = T(a)T(b) + H(a)H(\tilde{b})$$

i.e.,

$$T(a)T(b) = T(ab) - H(a)H(\tilde{b}).$$

can tell us things about the Fredholmness of $T(a)$ or $T(b)$ if we know things about a and b . In particular, if $a \in C$ is invertible with continuous inverse, then

$$T(a)T(a^{-1}) = Id + K_1$$

and

$$T(a^{-1})T(a) = Id + K_2$$

so we can conclude right away $T(a)$ and $T(a^{-1})$ are Fredholm.

Some uses for the last theorem

Wiener-Hopf Factorization: writing a symbol a in a Banach algebra A as $a = a_- \chi_n a_+$ where

$$a_- \in A_- = \{a \in A : a_n = 0 \ \forall n > 0\},$$

$$a_+ \in A_+ = \{a \in A : a_n = 0 \ \forall n < 0\}.$$

Some uses for the last theorem

Wiener-Hopf Factorization: writing a symbol a in a Banach algebra A as $a = a_- \chi_n a_+$ where

$$a_- \in A_- = \{a \in A : a_n = 0 \ \forall n > 0\},$$

$$a_+ \in A_+ = \{a \in A : a_n = 0 \ \forall n < 0\}.$$

Notice that $H(a_-) = H(\tilde{a}_+) = 0$.

Some uses for the last theorem

Wiener-Hopf Factorization: writing a symbol a in a Banach algebra A as $a = a_- \chi_n a_+$ where

$$a_- \in A_- = \{a \in A : a_n = 0 \ \forall n > 0\},$$

$$a_+ \in A_+ = \{a \in A : a_n = 0 \ \forall n < 0\}.$$

Notice that $H(a_-) = H(\tilde{a}_+) = 0$.

Now iterate the previous theorem twice:

$$\begin{aligned} T(a) &= T(a_- \chi_n a_+) \\ &= T(a_-) T(\chi_n a_+) + H(a_-) H(\widetilde{\chi_n a_+}) \\ &= T(a_-) (T(\chi_n) T(a_+) + H(\chi_n) H(\tilde{a}_+)) \\ &= T(a_-) T(\chi_n) T(a_+)! \end{aligned}$$

Some uses for the last theorem

Wiener-Hopf Factorization: writing a symbol a in a Banach algebra A as $a = a_- \chi_n a_+$ where

$$a_- \in A_- = \{a \in A : a_n = 0 \ \forall n > 0\},$$

$$a_+ \in A_+ = \{a \in A : a_n = 0 \ \forall n < 0\}.$$

Notice that $H(a_-) = H(\tilde{a}_+) = 0$.

Now iterate the previous theorem twice:

$$\begin{aligned} T(a) &= T(a_- \chi_n a_+) \\ &= T(a_-) T(\chi_n a_+) + H(a_-) H(\widetilde{\chi_n a_+}) \\ &= T(a_-) (T(\chi_n) T(a_+) + H(\chi_n) H(\tilde{a}_+)) \\ &= T(a_-) T(\chi_n) T(a_+)! \end{aligned}$$

Nothing special about χ_n in this equality, it just becomes relevant in WHF.

Some uses for the last theorem

Wiener-Hopf Factorization: writing a symbol a in a Banach algebra A as $a = a_- \chi_n a_+$ where

$$a_- \in A_- = \{a \in A : a_n = 0 \ \forall n > 0\},$$

$$a_+ \in A_+ = \{a \in A : a_n = 0 \ \forall n < 0\}.$$

Notice that $H(a_-) = H(\tilde{a}_+) = 0$.

Now iterate the previous theorem twice:

$$\begin{aligned} T(a) &= T(a_- \chi_n a_+) \\ &= T(a_-) T(\chi_n a_+) + H(a_-) H(\widetilde{\chi_n a_+}) \\ &= T(a_-) (T(\chi_n) T(a_+) + H(\chi_n) H(\tilde{a}_+)) \\ &= T(a_-) T(\chi_n) T(a_+)! \end{aligned}$$

Nothing special about χ_n in this equality, it just becomes relevant in WHF. There's lots of theory in WHF, this isn't even scratching the surface !

Playground for Problems: Approximation Methods

Let A be an infinite matrix that generates a bounded operator on ℓ^2 (e.g., a Toeplitz matrix with essentially bounded, measurable symbol if you like).

Let A be an infinite matrix that generates a bounded operator on ℓ^2 (e.g., a Toeplitz matrix with essentially bounded, measurable symbol if you like). To solve $Ax = y$, we consider the truncated systems where we truncate A to an $n \times n$ matrix and truncate x and y as well.

Let A be an infinite matrix that generates a bounded operator on ℓ^2 (e.g., a Toeplitz matrix with essentially bounded, measurable symbol if you like). To solve $Ax = y$, we consider the truncated systems where we truncate A to an $n \times n$ matrix and truncate x and y as well. Specifically, we can define

$P_n : \{x_1, x_2, x_3, \dots\} \mapsto \{x_1, x_2, \dots, x_n, 0, 0, \dots\}$ and instead of solving $Ax = y$ we aim to solve $P_n A x^{(n)} = P_n y$ where $x^{(n)}$ denotes any vector in the image of P_n .

Let A be an infinite matrix that generates a bounded operator on ℓ^2 (e.g., a Toeplitz matrix with essentially bounded, measurable symbol if you like). To solve $Ax = y$, we consider the truncated systems where we truncate A to an $n \times n$ matrix and truncate x and y as well. Specifically, we can define

$P_n : \{x_1, x_2, x_3, \dots\} \mapsto \{x_1, x_2, \dots, x_n, 0, 0, \dots\}$ and instead of solving $Ax = y$ we aim to solve $P_n A x^{(n)} = P_n y$ where $x^{(n)}$ denotes any vector in the image of P_n . Said differently, we are considering the operator $P_n A P_n$ restricted to the image of P_n . Restricting to this is called the *finite section method*.

Some Things to Consider

In Numerical Analysis, they hold the following principle closely to heart:

$$\text{convergence} = \text{approximation} + \text{stability}$$

Some Things to Consider

In Numerical Analysis, they hold the following principle closely to heart:

$$\text{convergence} = \text{approximation} + \text{stability}$$

We will almost always require “approximation” in our assumptions.

Some Things to Consider

In Numerical Analysis, they hold the following principle closely to heart:

$$\text{convergence} = \text{approximation} + \text{stability}$$

We will almost always require “approximation” in our assumptions. The questions we ask here are:

1. Does the sequence $\{P_n A P_n\}$ approximate A *well*? (and what does “well” mean?)

Some Things to Consider

In Numerical Analysis, they hold the following principle closely to heart:

$$\text{convergence} = \text{approximation} + \text{stability}$$

We will almost always require “approximation” in our assumptions. The questions we ask here are:

1. Does the sequence $\{P_n A P_n\}$ approximate A *well*? (and what does “well” mean?)
2. Is the sequence stable?

Appropriate Approximations

We say that a sequence $\{A_n\}$ is an *appropriate approximation* for A if

1. There exists an $n_0 \in \mathbb{N}$ such that A_n is invertible for all $n \geq n_0$ and
2. For all $y \in \ell^2$, the (unique) solutions to $A_n x^{(n)} = P_n y$ converge in ℓ^2 to a solution $x \in \ell^2$ of the equation $Ax = y$.

Appropriate Approximations

We say that a sequence $\{A_n\}$ is an *appropriate approximation* for A if

1. There exists an $n_0 \in \mathbb{N}$ such that A_n is invertible for all $n \geq n_0$ and
2. For all $y \in \ell^2$, the (unique) solutions to $A_n x^{(n)} = P_n y$ converge in ℓ^2 to a solution $x \in \ell^2$ of the equation $Ax = y$.

We have the following:

Theorem

Let $A \in B(\ell^2)$ and let $\{A_n\}$ be an approximating sequence for A . Then $\{A_n\}$ is an appropriate approximating sequence for A if and only if A is invertible, the matrices A_n are all invertible for sufficiently large n , and $A_n^{-1} := A_n^{-1}P_n$ converges strongly to A^{-1} .

Closely related to this idea of appropriate approximations is the notion of stability. We call a sequence of matrices $\{A_n\}$ stable if A_n is invertible for all sufficiently large n (say $n \geq n_0$) and $\sup_{n \geq n_0} \|A_n^{-1}\| < \infty$.

Closely related to this idea of appropriate approximations is the notion of stability. We call a sequence of matrices $\{A_n\}$ stable if A_n is invertible for all sufficiently large n (say $n \geq n_0$) and $\sup_{n \geq n_0} \|A_n^{-1}\| < \infty$. The following theorem relates the two notions we've discussed so far:

Theorem

Let $A \in B(\ell^2)$ and let $\{A_n\}$ be an approximating sequence for A . Then $\{A_n\}$ is an appropriate approximating sequence for A if and only if A is invertible and $\{A_n\}$ is a stable sequence.

Methods for Determining Stability

- The tools used to determine stability are functional analytic in nature, but there are also more algebraic tools used to determine this.

Methods for Determining Stability

- The tools used to determine stability are functional analytic in nature, but there are also more algebraic tools used to determine this.
- The more algebraic approach follows this strategy: Can we build a Banach algebra G so that $\{A_n\}$ being stable is equivalent to something being invertible in G ?

Methods for Determining Stability

- The tools used to determine stability are functional analytic in nature, but there are also more algebraic tools used to determine this.
- The more algebraic approach follows this strategy: Can we build a Banach algebra G so that $\{A_n\}$ being stable is equivalent to something being invertible in G ?
- We'll revisit this notion soon without getting too technical, but for now let's have a bit of fun.

Some Tools for the Toolbox

Banach Algebras

A Banach algebra is a Banach space equipped with a multiplication whose norm satisfies the property that $\|ab\| \leq \|a\| \cdot \|b\|$. Notice that this automatically gives continuity of multiplication.

Banach Algebras

A Banach algebra is a Banach space equipped with a multiplication whose norm satisfies the property that $\|ab\| \leq \|a\| \cdot \|b\|$. Notice that this automatically gives continuity of multiplication. They don't have to be unital, but we can always embed them into a unital one (and they will be of codimension 1).

Banach Algebras

A Banach algebra is a Banach space equipped with a multiplication whose norm satisfies the property that $\|ab\| \leq \|a\| \cdot \|b\|$. Notice that this automatically gives continuity of multiplication. They don't have to be unital, but we can always embed them into a unital one (and they will be of codimension 1). Examples of Banach algebras:

Banach Algebras

A Banach algebra is a Banach space equipped with a multiplication whose norm satisfies the property that $\|ab\| \leq \|a\| \cdot \|b\|$. Notice that this automatically gives continuity of multiplication. They don't have to be unital, but we can always embed them into a unital one (and they will be of codimension 1). Examples of Banach algebras:

1. For any Banach space X , the space $B(X)$ of bounded linear operators on X with multiplication given by composition and equipped with operator norm

Banach Algebras

A Banach algebra is a Banach space equipped with a multiplication whose norm satisfies the property that $\|ab\| \leq \|a\| \cdot \|b\|$. Notice that this automatically gives continuity of multiplication. They don't have to be unital, but we can always embed them into a unital one (and they will be of codimension 1). Examples of Banach algebras:

1. For any Banach space X , the space $B(X)$ of bounded linear operators on X with multiplication given by composition and equipped with operator norm
2. $C(K)$ for any compact topological space K equipped with the sup norm

Banach Algebras

A Banach algebra is a Banach space equipped with a multiplication whose norm satisfies the property that $\|ab\| \leq \|a\| \cdot \|b\|$. Notice that this automatically gives continuity of multiplication. They don't have to be unital, but we can always embed them into a unital one (and they will be of codimension 1). Examples of Banach algebras:

1. For any Banach space X , the space $B(X)$ of bounded linear operators on X with multiplication given by composition and equipped with operator norm
2. $C(K)$ for any compact topological space K equipped with the sup norm
3. \mathbb{R} and \mathbb{C} (fan favorite?)

Banach Algebras

A Banach algebra is a Banach space equipped with a multiplication whose norm satisfies the property that $\|ab\| \leq \|a\| \cdot \|b\|$. Notice that this automatically gives continuity of multiplication. They don't have to be unital, but we can always embed them into a unital one (and they will be of codimension 1). Examples of Banach algebras:

1. For any Banach space X , the space $B(X)$ of bounded linear operators on X with multiplication given by composition and equipped with operator norm
2. $C(K)$ for any compact topological space K equipped with the sup norm
3. \mathbb{R} and \mathbb{C} (fan favorite?)
4. The Wiener algebra $W = \{f \in C(\mathbb{T}) : \sum_{n \in \mathbb{Z}} |\hat{f}_n| < \infty\}$. Can you guess what the norm is?

Another Example

Define the set \mathcal{F} to be the set of all sequences $\{A_n\}$ of operators (matrices) $A_n \in B(\text{Im}(P_n))(\cong \mathbb{C}^{n \times n})$ for which

$$\|\{A_n\}\| := \sup_{n \geq 1} \|A_n\| < \infty$$

Another Example

Define the set \mathcal{F} to be the set of all sequences $\{A_n\}$ of operators (matrices) $A_n \in B(\text{Im}(P_n))(\cong \mathbb{C}^{n \times n})$ for which

$$\|\{A_n\}\| := \sup_{n \geq 1} \|A_n\| < \infty$$

This set is a Banach algebra when equipped with the above norm and the following operations:

$$\{A_n\} + \{B_n\} := \{A_n + B_n\}, \lambda\{A_n\} := \{\lambda A_n\}, \{A_n\}\{B_n\} := \{A_n B_n\}$$

Another Example

Define the set \mathcal{F} to be the set of all sequences $\{A_n\}$ of operators (matrices) $A_n \in B(\text{Im}(P_n)) (\cong \mathbb{C}^{n \times n})$ for which

$$\|\{A_n\}\| := \sup_{n \geq 1} \|A_n\| < \infty$$

This set is a Banach algebra when equipped with the above norm and the following operations:

$$\{A_n\} + \{B_n\} := \{A_n + B_n\}, \lambda\{A_n\} := \{\lambda A_n\}, \{A_n\}\{B_n\} := \{A_n B_n\}$$

Denote by \mathcal{N} the subset of \mathcal{F} consisting of all $\{C_n\} \in \mathcal{F}$ such that $\|C_n\| \rightarrow 0$ as $n \rightarrow \infty$.

Another Example

Define the set \mathcal{F} to be the set of all sequences $\{A_n\}$ of operators (matrices) $A_n \in B(\text{Im}(P_n))(\cong \mathbb{C}^{n \times n})$ for which

$$\|\{A_n\}\| := \sup_{n \geq 1} \|A_n\| < \infty$$

This set is a Banach algebra when equipped with the above norm and the following operations:

$$\{A_n\} + \{B_n\} := \{A_n + B_n\}, \lambda\{A_n\} := \{\lambda A_n\}, \{A_n\}\{B_n\} := \{A_n B_n\}$$

Denote by \mathcal{N} the subset of \mathcal{F} consisting of all $\{C_n\} \in \mathcal{F}$ such that $\|C_n\| \rightarrow 0$ as $n \rightarrow \infty$. This is a closed 2-sided ideal of \mathcal{F} and hence \mathcal{F}/\mathcal{N} is a Banach algebra.

Our First Pass at Algebraization of Stability

Theorem

A sequence $\{A_n\} \in \mathcal{F}$ is stable if and only if the coset $\{A_n\} + \mathcal{N}$ is invertible in \mathcal{F}/\mathcal{N} .

Our First Pass at Algebraization of Stability

Theorem

A sequence $\{A_n\} \in \mathcal{F}$ is stable if and only if the coset $\{A_n\} + \mathcal{N}$ is invertible in \mathcal{F}/\mathcal{N} .

Problem: \mathcal{N} is in a sense too small and \mathcal{F} is too big.

Our First Pass at Algebraization of Stability

Theorem

A sequence $\{A_n\} \in \mathcal{F}$ is stable if and only if the coset $\{A_n\} + \mathcal{N}$ is invertible in \mathcal{F}/\mathcal{N} .

Problem: \mathcal{N} is in a sense too small and \mathcal{F} is too big.

For finite Toeplitz matrices, we have

$$T_n(a)T_n(b) = T_n(ab) - P_n H(a)H(\tilde{b})P_n - W_n H(\tilde{a})H(b)W_n$$

Our First Pass at Algebraization of Stability

Theorem

A sequence $\{A_n\} \in \mathcal{F}$ is stable if and only if the coset $\{A_n\} + \mathcal{N}$ is invertible in \mathcal{F}/\mathcal{N} .

Problem: \mathcal{N} is in a sense too small and \mathcal{F} is too big.

For finite Toeplitz matrices, we have

$$T_n(a)T_n(b) = T_n(ab) - P_n H(a)H(\tilde{b})P_n - W_n H(\tilde{a})H(b)W_n$$

The next step would be to construct a smaller algebra that contains what we want while also containing all elements of the form $\{T_n(a)\}$.

Our First Pass at Algebraization of Stability

Theorem

A sequence $\{A_n\} \in \mathcal{F}$ is stable if and only if the coset $\{A_n\} + \mathcal{N}$ is invertible in \mathcal{F}/\mathcal{N} .

Problem: \mathcal{N} is in a sense too small and \mathcal{F} is too big.

For finite Toeplitz matrices, we have

$$T_n(a)T_n(b) = T_n(ab) - P_n H(a)H(\tilde{b})P_n - W_n H(\tilde{a})H(b)W_n$$

The next step would be to construct a smaller algebra that contains what we want while also containing all elements of the form $\{T_n(a)\}$. Instead of doing this, let's resume our journey through general Banach algebra theory.

Our First Pass at Algebraization of Stability

Theorem

A sequence $\{A_n\} \in \mathcal{F}$ is stable if and only if the coset $\{A_n\} + \mathcal{N}$ is invertible in \mathcal{F}/\mathcal{N} .

Problem: \mathcal{N} is in a sense too small and \mathcal{F} is too big.

For finite Toeplitz matrices, we have

$$T_n(a)T_n(b) = T_n(ab) - P_n H(a)H(\tilde{b})P_n - W_n H(\tilde{a})H(b)W_n$$

The next step would be to construct a smaller algebra that contains what we want while also containing all elements of the form $\{T_n(a)\}$. Instead of doing this, let's resume our journey through general Banach algebra theory. Next stop: Maximal Ideals!

- In a Banach algebra A , every maximal ideal (which is proper by definition) is necessarily closed.

Banach Algebras: Ideals

- In a Banach algebra A , every maximal ideal (which is proper by definition) is necessarily closed.
- Maximal ideals are closely related to *multiplicative linear functionals*, which are nonzero linear functionals $f : A \rightarrow \mathbb{C}$ satisfying $f(ab) = f(a)f(b)$.

Banach Algebras: Ideals

- In a Banach algebra A , every maximal ideal (which is proper by definition) is necessarily closed.
- Maximal ideals are closely related to *multiplicative linear functionals*, which are nonzero linear functionals $f : A \rightarrow \mathbb{C}$ satisfying $f(ab) = f(a)f(b)$.
- Thanks to Gelfand-Mazur, we know there is a one-to-one correspondence between multiplicative linear functionals and maximal ideals of a Banach algebra (via associating with each functional its kernel).

Maximal Ideal Space

Perhaps a bit misleading from the name, but justified from the last slide, the maximal ideal space M of a (commutative, unital) Banach algebra is actually the space of nonzero multiplicative linear functionals.

Maximal Ideal Space

Perhaps a bit misleading from the name, but justified from the last slide, the maximal ideal space M of a (commutative, unital) Banach algebra is actually the space of nonzero multiplicative linear functionals.

Do you think we can equip it with a topology?

Maximal Ideal Space: Giving it a Topology

For each element $a \in A$ we can assign it a function $\hat{a} : M \rightarrow \mathbb{C}$ defined by $\hat{a}(f) = f(a)$. This function is called the *Gelfand transform* of a . The map $\gamma : A \rightarrow C(M)$ is called the *Gelfand map*.

Maximal Ideal Space: Giving it a Topology

For each element $a \in A$ we can assign it a function $\hat{a} : M \rightarrow \mathbb{C}$ defined by $\hat{a}(f) = f(a)$. This function is called the *Gelfand transform* of a . The map $\gamma : A \rightarrow C(M)$ is called the *Gelfand map*.

Define the set $\hat{A} := \{\hat{a} : a \in A\}$. The *Gelfand topology* is the coarsest (weakest) topology on M such that all functions $\hat{a} \in \hat{A}$ are continuous. If you prefer, we can think of M as a subspace of A^* and then the Gelfand topology is nothing but the subspace topology where A^* is equipped with the weak-* topology.

Maximal Ideal Space: Giving it a Topology

For each element $a \in A$ we can assign it a function $\hat{a} : M \rightarrow \mathbb{C}$ defined by $\hat{a}(f) = f(a)$. This function is called the *Gelfand transform* of a . The map $\gamma : A \rightarrow C(M)$ is called the *Gelfand map*.

Define the set $\hat{A} := \{\hat{a} : a \in A\}$. The *Gelfand topology* is the coarsest (weakest) topology on M such that all functions $\hat{a} \in \hat{A}$ are continuous. If you prefer, we can think of M as a subspace of A^* and then the Gelfand topology is nothing but the subspace topology where A^* is equipped with the weak-* topology.

Can show that M is a compact Hausdorff space.

Why is M compact?

(Outline)

1. First show it is closed using the definition of the open sets in M : For $\phi \in M$, the open ball of radius $\epsilon > 0$ centered at ϕ is

$$B_{x_1, x_2, \dots, x_n, \epsilon}(\phi) = \{\psi \in M : |\phi(x_i) - \psi(x_i)| < \epsilon \ \forall i\}$$

Why is M compact?

(Outline)

1. First show it is closed using the definition of the open sets in M : For $\phi \in M$, the open ball of radius $\epsilon > 0$ centered at ϕ is

$$B_{x_1, x_2, \dots, x_n, \epsilon}(\phi) = \{\psi \in M : |\phi(x_i) - \psi(x_i)| < \epsilon \ \forall i\}$$

2. Next show that for any $\phi \in M$, $\|\phi\| \leq 1$ (uses Neumann series)

Why is M compact?

(Outline)

1. First show it is closed using the definition of the open sets in M : For $\phi \in M$, the open ball of radius $\epsilon > 0$ centered at ϕ is

$$B_{x_1, x_2, \dots, x_n, \epsilon}(\phi) = \{\psi \in M : |\phi(x_i) - \psi(x_i)| < \epsilon \ \forall i\}$$

2. Next show that for any $\phi \in M$, $\|\phi\| \leq 1$ (uses Neumann series)
3. Then sit and fondly remember Banach-Alaoglu (the closed unit ball is compact in the weak* – topology)

Why is M compact?

(Outline)

1. First show it is closed using the definition of the open sets in M : For $\phi \in M$, the open ball of radius $\epsilon > 0$ centered at ϕ is

$$B_{x_1, x_2, \dots, x_n, \epsilon}(\phi) = \{\psi \in M : |\phi(x_i) - \psi(x_i)| < \epsilon \ \forall i\}$$

2. Next show that for any $\phi \in M$, $\|\phi\| \leq 1$ (uses Neumann series)
3. Then sit and fondly remember Banach-Alaoglu (the closed unit ball is compact in the weak* – topology)
4. Finally, since it is a closed subset of a compact set, it is itself compact

Theorem

Let A be a commutative Banach algebra with unit and let M be the maximal ideal space of A . An element $a \in A$ is invertible if and only if $\hat{a}(m) \neq 0$ for all $m \in M$.

Theorem

Let A be a commutative Banach algebra with unit and let M be the maximal ideal space of A . An element $a \in A$ is invertible if and only if $\hat{a}(m) \neq 0$ for all $m \in M$.

In English: The Gelfand map is a Banach algebra homomorphism of A into $C(M)$ which preserves spectra. This is seen by showing that the range of \hat{a} is equal to the spectrum of a .

Last Theorem in Action: The Wiener Algebra

Recall: The Wiener algebra is $W = \{a \in C(\mathbb{T}) : \sum_{n \in \mathbb{Z}} |\hat{a}_n| < \infty\}$

Last Theorem in Action: The Wiener Algebra

Recall: The Wiener algebra is $W = \{a \in C(\mathbb{T}) : \sum_{n \in \mathbb{Z}} |\hat{a}_n| < \infty\}$

Wiener's Theorem states that $a \in W$ is invertible in W if and only if $a(\tau) \neq 0$ for all $\tau \in \mathbb{T}$.

Last Theorem in Action: The Wiener Algebra

Recall: The Wiener algebra is $W = \{a \in C(\mathbb{T}) : \sum_{n \in \mathbb{Z}} |\hat{a}_n| < \infty\}$

Wiener's Theorem states that $a \in W$ is invertible in W if and only if $a(\tau) \neq 0$ for all $\tau \in \mathbb{T}$. The multiplicative linear functionals of W are the maps

$$\phi_\tau : W \rightarrow \mathbb{C}$$

that send $a \in W$ to $a(\tau)$ [$\tau \in \mathbb{T}$].

Last Theorem in Action: The Wiener Algebra

Recall: The Wiener algebra is $W = \{a \in C(\mathbb{T}) : \sum_{n \in \mathbb{Z}} |\hat{a}_n| < \infty\}$

Wiener's Theorem states that $a \in W$ is invertible in W if and only if $a(\tau) \neq 0$ for all $\tau \in \mathbb{T}$. The multiplicative linear functionals of W are the maps

$$\phi_\tau : W \rightarrow \mathbb{C}$$

that send $a \in W$ to $a(\tau)$ [$\tau \in \mathbb{T}$].

We can thus identify the maximal ideal space of W with \mathbb{T} , so here the Gelfand map is just the embedding $\gamma : W \rightarrow C(\mathbb{T})$.

Last Theorem in Action: The Wiener Algebra

Recall: The Wiener algebra is $W = \{a \in C(\mathbb{T}) : \sum_{n \in \mathbb{Z}} |\hat{a}_n| < \infty\}$

Wiener's Theorem states that $a \in W$ is invertible in W if and only if $a(\tau) \neq 0$ for all $\tau \in \mathbb{T}$. The multiplicative linear functionals of W are the maps

$$\phi_\tau : W \rightarrow \mathbb{C}$$

that send $a \in W$ to $a(\tau)$ [$\tau \in \mathbb{T}$].

We can thus identify the maximal ideal space of W with \mathbb{T} , so here the Gelfand map is just the embedding $\gamma : W \rightarrow C(\mathbb{T})$.

The last theorem then gives a proof of Wiener's Theorem, since $\hat{a} : M (\cong \mathbb{T}) \rightarrow \mathbb{C}$ is exactly the mapping $\hat{a}(\phi_\tau) = \phi_\tau(a) := a(\tau)$

What if we lack commutativity?

We now have a way to associate with every element a of a commutative, unital Banach algebra a collection of numbers in terms of which we can decide whether or not a is invertible. In the case where A is not commutative, we have the following theorem:

What if we lack commutativity?

We now have a way to associate with every element a of a commutative, unital Banach algebra a collection of numbers in terms of which we can decide whether or not a is invertible. In the case where A is not commutative, we have the following theorem:

Theorem

Let A be a Banach algebra with identity element e and let Z be a closed subalgebra of the center of A which contains e . Denote the maximal ideal space of Z by Ω and for each maximal ideal $\omega \in \Omega$ let J_ω be the smallest closed two-sided ideal of A which contains the set ω . Then an element $a \in A$ is invertible in A if and only if the coset $a + J_\omega$ is invertible in A/J_ω for every $\omega \in \Omega$.

Theorem

Let A be a Banach algebra with identity element e and let Z be a closed subalgebra of the center of A which contains e . Denote the maximal ideal space of Z by Ω and for each maximal ideal $\omega \in \Omega$ let J_ω be the smallest closed two-sided ideal of A which contains the set ω . Then an element $a \in A$ is invertible in A if and only if the coset $a + J_\omega$ is invertible in A/J_ω for every $\omega \in \Omega$.

Theorem

Let A be a Banach algebra with identity element e and let Z be a closed subalgebra of the center of A which contains e . Denote the maximal ideal space of Z by Ω and for each maximal ideal $\omega \in \Omega$ let J_ω be the smallest closed two-sided ideal of A which contains the set ω . Then an element $a \in A$ is invertible in A if and only if the coset $a + J_\omega$ is invertible in A/J_ω for every $\omega \in \Omega$.

- If A is commutative, we can take $Z = A$ and since $A/J_\omega = A/\omega \cong \mathbb{C}$ we end up in the theorem from before

Theorem

Let A be a Banach algebra with identity element e and let Z be a closed subalgebra of the center of A which contains e . Denote the maximal ideal space of Z by Ω and for each maximal ideal $\omega \in \Omega$ let J_ω be the smallest closed two-sided ideal of A which contains the set ω . Then an element $a \in A$ is invertible in A if and only if the coset $a + J_\omega$ is invertible in A/J_ω for every $\omega \in \Omega$.

- If A is commutative, we can take $Z = A$ and since $A/J_\omega = A/\omega \cong \mathbb{C}$ we end up in the theorem from before
- If the center is trivial, we get that a is invertible if and only if a is invertible (spicy, I know).

Example: Operators of Local Type

- The set $J := \{A \in B(\ell^2) : AT(c) - T(c)A \in K(\ell^2) \forall c \in C\}$ are called *operators of local type*. It contains the compact operators and it contains $T(a)$ with $a \in L^\infty$.

Example: Operators of Local Type

- The set $J := \{A \in B(\ell^2) : AT(c) - T(c)A \in K(\ell^2) \forall c \in C\}$ are called *operators of local type*. It contains the compact operators and it contains $T(a)$ with $a \in L^\infty$.
- Put $J^\pi = J/K(\ell^2)$. Then the algebra $\{T(c) + K(\ell^2) : c \in C\}$ is in the center of J^π .

Example: Operators of Local Type

- The set $J := \{A \in B(\ell^2) : AT(c) - T(c)A \in K(\ell^2) \forall c \in C\}$ are called *operators of local type*. It contains the compact operators and it contains $T(a)$ with $a \in L^\infty$.
- Put $J^\pi = J/K(\ell^2)$. Then the algebra $\{T(c) + K(\ell^2) : c \in C\}$ is in the center of J^π .
- J^π is inverse closed in $B(\ell^2)/K(\ell^2)$ (i.e., if $A \in J$ and $A + K(\ell^2)$ is invertible in $B(\ell^2)/K(\ell^2)$, then the inverse of $A + K(\ell^2)$ is in J^π .)

Example: Operators of Local Type

- The set $J := \{A \in B(\ell^2) : AT(c) - T(c)A \in K(\ell^2) \forall c \in C\}$ are called *operators of local type*. It contains the compact operators and it contains $T(a)$ with $a \in L^\infty$.
- Put $J^\pi = J/K(\ell^2)$. Then the algebra $\{T(c) + K(\ell^2) : c \in C\}$ is in the center of J^π .
- J^π is inverse closed in $B(\ell^2)/K(\ell^2)$ (i.e., if $A \in J$ and $A + K(\ell^2)$ is invertible in $B(\ell^2)/K(\ell^2)$, then the inverse of $A + K(\ell^2)$ is in J^π .)
- Altogether: $A \in J$ is Fredholm if and only if $A + K(\ell^2)$ is invertible in J^π

Example: Operators of Local Type

- The set $J := \{A \in B(\ell^2) : AT(c) - T(c)A \in K(\ell^2) \forall c \in C\}$ are called *operators of local type*. It contains the compact operators and it contains $T(a)$ with $a \in L^\infty$.
- Put $J^\pi = J/K(\ell^2)$. Then the algebra $\{T(c) + K(\ell^2) : c \in C\}$ is in the center of J^π .
- J^π is inverse closed in $B(\ell^2)/K(\ell^2)$ (i.e., if $A \in J$ and $A + K(\ell^2)$ is invertible in $B(\ell^2)/K(\ell^2)$, then the inverse of $A + K(\ell^2)$ is in J^π .)
- Altogether: $A \in J$ is Fredholm if and only if $A + K(\ell^2)$ is invertible in J^π

Can therefore use Allen-Douglass with $A = J^\pi$ and $Z = \{T(c) + K(\ell^2) : c \in C\}$ to study Fredholmness of operators of local type.

A C^* –Algebra is a Banach algebra A equipped with a map $*$: $A \rightarrow A$ (called an involution) satisfying the following:

A C^* –Algebra is a Banach algebra A equipped with a map $*$: $A \rightarrow A$ (called an involution) satisfying the following:

For each $a, b \in A$ and $\lambda \in \mathbb{C}$,

1. $a^{**} = a$

A C^* –Algebra is a Banach algebra A equipped with a map $*$: $A \rightarrow A$ (called an involution) satisfying the following:

For each $a, b \in A$ and $\lambda \in \mathbb{C}$,

1. $a^{**} = a$
2. $(a + b)^* = a^* + b^*$

A C^* –Algebra is a Banach algebra A equipped with a map $*$: $A \rightarrow A$ (called an involution) satisfying the following:

For each $a, b \in A$ and $\lambda \in \mathbb{C}$,

1. $a^{**} = a$
2. $(a + b)^* = a^* + b^*$
3. $(ab)^* = b^* a^*$

A C^* –Algebra is a Banach algebra A equipped with a map $*$: $A \rightarrow A$ (called an involution) satisfying the following:

For each $a, b \in A$ and $\lambda \in \mathbb{C}$,

1. $a^{**} = a$
2. $(a + b)^* = a^* + b^*$
3. $(ab)^* = b^* a^*$
4. $(\lambda a)^* = \bar{\lambda} a^*$

A C^* –Algebra is a Banach algebra A equipped with a map $*$: $A \rightarrow A$ (called an involution) satisfying the following:

For each $a, b \in A$ and $\lambda \in \mathbb{C}$,

1. $a^{**} = a$
2. $(a + b)^* = a^* + b^*$
3. $(ab)^* = b^* a^*$
4. $(\lambda a)^* = \bar{\lambda} a^*$
5. $\|a\|^2 = \|aa^*\|$

Examples

1. If X is compact Hausdorff, then $C(X)$ is a C^* -algebra with involution given by complex conjugation (i.e., $f^*(x) = \overline{f(x)}$)

Examples

1. If X is compact Hausdorff, then $C(X)$ is a C^* -algebra with involution given by complex conjugation (i.e., $f^*(x) = \overline{f(x)}$)
Similarly, L^∞ is a C^* -algebra.

Examples

1. If X is compact Hausdorff, then $C(X)$ is a C^* -algebra with involution given by complex conjugation (i.e., $f^*(x) = \overline{f(x)}$)
Similarly, L^∞ is a C^* -algebra.
2. Non-example: Complex conjugation defines an involution on the Wiener algebra W , but it fails property 5. Take $a(t) = -t^{-1} + 1 + t$ for example.

Examples

1. If X is compact Hausdorff, then $C(X)$ is a C^* -algebra with involution given by complex conjugation (i.e., $f^*(x) = \overline{f(x)}$). Similarly, L^∞ is a C^* -algebra.
2. Non-example: Complex conjugation defines an involution on the Wiener algebra W , but it fails property 5. Take $a(t) = -t^{-1} + 1 + t$ for example. Here, $(a\bar{a})(t) = -t^{-2} + 3 - t^2$, so $\|a\bar{a}\|_W = 5$ while $\|a\|_W^2 = 9$.

Examples

1. If X is compact Hausdorff, then $C(X)$ is a C^* -algebra with involution given by complex conjugation (i.e., $f^*(x) = \overline{f(x)}$). Similarly, L^∞ is a C^* -algebra.
2. Non-example: Complex conjugation defines an involution on the Wiener algebra W , but it fails property 5. Take $a(t) = -t^{-1} + 1 + t$ for example. Here, $(a\bar{a})(t) = -t^{-2} + 3 - t^2$, so $\|a\bar{a}\|_W = 5$ while $\|a\|_W^2 = 9$.
3. If H is a Hilbert space, then the algebras $B(H)$ and $K(H)$ are C^* -algebras with involution being passage to the adjoint.

Examples

1. If X is compact Hausdorff, then $C(X)$ is a C^* -algebra with involution given by complex conjugation (i.e., $f^*(x) = \overline{f(x)}$). Similarly, L^∞ is a C^* -algebra.
2. Non-example: Complex conjugation defines an involution on the Wiener algebra W , but it fails property 5. Take $a(t) = -t^{-1} + 1 + t$ for example. Here, $(a\bar{a})(t) = -t^{-2} + 3 - t^2$, so $\|a\bar{a}\|_W = 5$ while $\|a\|_W^2 = 9$.
3. If H is a Hilbert space, then the algebras $B(H)$ and $K(H)$ are C^* -algebras with involution being passage to the adjoint. $B(H)$ in a sense is *the* example due to Gelfand-Naimark.

Examples

1. If X is compact Hausdorff, then $C(X)$ is a C^* -algebra with involution given by complex conjugation (i.e., $f^*(x) = \overline{f(x)}$)
Similarly, L^∞ is a C^* -algebra.
2. Non-example: Complex conjugation defines an involution on the Wiener algebra W , but it fails property 5. Take $a(t) = -t^{-1} + 1 + t$ for example. Here,
 $(a\bar{a})(t) = -t^{-2} + 3 - t^2$, so $\|a\bar{a}\|_W = 5$ while $\|a\|_W^2 = 9$.
3. If H is a Hilbert space, then the algebras $B(H)$ and $K(H)$ are C^* -algebras with involution being passage to the adjoint.
 $B(H)$ in a sense is *the* example due to Gelfand-Naimark.
4. The quotient of any C^* -algebra with a self-adjoint, closed, 2-sided ideal will also be a C^* -algebra

We know in general, if B is a closed subalgebra of A , then for an element $b \in B$, the spectrum of b in B may be larger than its spectrum in A .

Inverse Closedness

We know in general, if B is a closed subalgebra of A , then for an element $b \in B$, the spectrum of b in B may be larger than its spectrum in A . In unital C^* -algebras, this never happens!

We know in general, if B is a closed subalgebra of A , then for an element $b \in B$, the spectrum of b in B may be larger than its spectrum in A . In unital C^* -algebras, this never happens!

Theorem

If A is a C^ -algebra with identity and B is a C^* -subalgebra of A which also contains identity, then for every $b \in B$*

$$sp_B b = sp_A b$$

Inverse Closedness

We know in general, if B is a closed subalgebra of A , then for an element $b \in B$, the spectrum of b in B may be larger than its spectrum in A . In unital C^* -algebras, this never happens!

Theorem

If A is a C^ -algebra with identity and B is a C^* -subalgebra of A which also contains identity, then for every $b \in B$*

$$sp_B b = sp_A b$$

This is more useful than it may seem...

- A C^* -algebra homomorphism f is a homomorphism in the algebraic sense between two C^* -algebras with the added property that $f(a^*) = f(a)^*$

C^* -algebra Homomorphisms

- A C^* -algebra homomorphism f is a homomorphism in the algebraic sense between two C^* -algebras with the added property that $f(a^*) = f(a)^*$
- It can be shown that the image $f(A)$ is a subalgebra of the codomain

C^* -algebra Homomorphisms

- A C^* -algebra homomorphism f is a homomorphism in the algebraic sense between two C^* -algebras with the added property that $f(a^*) = f(a)^*$
- It can be shown that the image $f(A)$ is a subalgebra of the codomain
- We say that such a homomorphism $f : A \rightarrow B$ between unital C^* -algebras *preserves spectra* if $spf(a) = spa$ for all $a \in A$ and we call it an isometry if $\|f(a)\| = \|a\|$ for all $a \in A$

The following fact is very useful:

Theorem

Let A and B be unital C^ -algebras and $f : A \rightarrow B$ be a C^* -algebra homomorphism.*

- 1. If f preserves spectra, then f also preserves norms.*
- 2. If f is injective, then f preserves spectra.*

Remember that when discussing stability from an algebraic perspective, we want to see if we can reduce stability of a sequence to invertibility in an algebra. The fact that injective $*$ -homomorphisms preserve spectra is huge in this regard !

Any questions?

