

Stratification of Tensor Triangular Categories

Applications to Motivic Categories

David Rubinstein

University of California, Santa Cruz

Outline of Presentation

- 1 Balmer Support and the Classification Theorem
- 2 Stratification
- 3 Derived Categories of Motives
- 4 My Research Problem

The Mathematical Landscape is Large

Categories \mathcal{C} are pervasive in all fields of mathematics

- Algebra
 - $\mathcal{C} = \text{Vect}_{\mathbb{F}}$;
 - $\mathcal{C} = \text{rep}(G)$;
 - ...

The Mathematical Landscape is Large

Categories \mathcal{C} are pervasive in all fields of mathematics

- Algebra
 - $\mathcal{C} = \text{Vect}_{\mathbb{F}}$;
 - $\mathcal{C} = \text{rep}(G)$;
 - ...
- Topology
 - $\mathcal{C} = \text{Top}$;
 - $\mathcal{C} = \text{CW}$;
 - ...

The Mathematical Landscape is Large

Categories \mathcal{C} are pervasive in all fields of mathematics

- Algebra
 - $\mathcal{C} = \text{Vect}_{\mathbb{F}}$;
 - $\mathcal{C} = \text{rep}(G)$;
 - ...
- Topology
 - $\mathcal{C} = \text{Top}$;
 - $\mathcal{C} = \text{CW}$;
 - ...
- Geometry
 - $\mathcal{C} = \text{SmMan}$;
 - $\mathcal{C} = \text{Schemes}$;
 - ...
- ...

Balmer Support and the Classification Theorem

Classification Goals: A mentsh trakht un Got lakht

Goal: classify the objects in a category up to isomorphism:

Balmer Support and the Classification Theorem

Classification Goals: A mentsh trakht un Got lakht

Goal: classify the objects in a category up to isomorphism:

- Classify finite dimensional vector spaces;

Balmer Support and the Classification Theorem

Classification Goals: A mentsh trakht un Got lakht

Goal: classify the objects in a category up to isomorphism:

- Classify finite dimensional vector spaces;
- Classify all finite dimensional representations of a group G ;

Classification Goals: A mentsh trakht un Got lakht

Goal: classify the objects in a category up to isomorphism:

- Classify finite dimensional vector spaces;
- Classify all finite dimensional representations of a group G ;
- Classify all low dimensional CW- complexes;
- ...

Balmer Support and the Classification Theorem

Classification Goals: A mentsh trakht un Got lakht

Goal: classify the objects in a category up to isomorphism:

- Classify finite dimensional vector spaces;
- Classify all finite dimensional representations of a group G ;
- Classify all low dimensional CW- complexes;
- ...

Unfortunately, we are confronted with "wild classification problems"

- Can't classify all finite dim representations of group G in positive characteristic case;

Balmer Support and the Classification Theorem

Classification Goals: A mentsh trakht un Got lakht

Goal: classify the objects in a category up to isomorphism:

- Classify finite dimensional vector spaces;
- Classify all finite dimensional representations of a group G ;
- Classify all low dimensional CW- complexes;
- ...

Unfortunately, we are confronted with "wild classification problems"

- Can't classify all finite dim representations of group G in positive characteristic case;
- Can't classify finite CW complexes up to homotopy equivalence;
- No more hope for classifying all complexes of sheaves on an algebraic variety V ;
- ...

Balmer Support and the Classification Theorem

Change in Paradigm

Important development in last few decades: change the category, and work "stably," in some "stable" category:

Balmer Support and the Classification Theorem

Change in Paradigm

Important development in last few decades: change the category, and work "stably," in some "stable" category:

- $\text{rep}(G) \rightsquigarrow \text{stab}(kG)$

Balmer Support and the Classification Theorem

Change in Paradigm

Important development in last few decades: change the category, and work "stably," in some "stable" category:

- $\text{rep}(G) \rightsquigarrow \text{stab}(kG)$
- $\text{HTop} \rightsquigarrow \text{SH}^{\text{fin}}$

Balmer Support and the Classification Theorem

Change in Paradigm

Important development in last few decades: change the category, and work "stably," in some "stable" category:

- $\text{rep}(G) \rightsquigarrow \text{stab}(kG)$
- $\text{HTop} \rightsquigarrow \text{SH}^{\text{fin}}$
- $\text{R-Mod} \rightsquigarrow \text{D}^{\text{perf}}(\text{R})$
- ...

Balmer Support and the Classification Theorem

Change in Paradigm

Important development in last few decades: change the category, and work "stably," in some "stable" category:

- $\text{rep}(G) \rightsquigarrow \text{stab}(kG)$
- $\text{HTop} \rightsquigarrow \text{SH}^{\text{fin}}$
- $\text{R-Mod} \rightsquigarrow \text{D}^{\text{perf}}(\text{R})$
- ...

These "stable" categories all have a tensor-triangulated category structure in common.

Balmer Support and the Classification Theorem

Change in Paradigm

Important development in last few decades: change the category, and work "stably," in some "stable" category:

- $\text{rep}(G) \rightsquigarrow \text{stab}(kG)$
- $\text{HTop} \rightsquigarrow \text{SH}^{\text{fin}}$
- $\text{R-Mod} \rightsquigarrow \text{D}^{\text{perf}}(\text{R})$
- ...

These "stable" categories all have a tensor-triangulated category structure in common.

- Instead of classifying objects "on the nose," classify objects "up to the tensor-triangulated structure"

Balmer Support and the Classification Theorem

Change in Paradigm

Important development in last few decades: change the category, and work "stably," in some "stable" category:

- $\text{rep}(G) \rightsquigarrow \text{stab}(kG)$
- $\text{HTop} \rightsquigarrow \text{SH}^{\text{fin}}$
- $\text{R-Mod} \rightsquigarrow \text{D}^{\text{perf}}(\text{R})$
- ...

These "stable" categories all have a tensor-triangulated category structure in common.

- Instead of classifying objects "on the nose," classify objects "up to the tensor-triangulated structure"
- Regard objects as being equivalent if they can build each other using the tt-structure

Balmer Support and the Classification Theorem

Change in Paradigm

Important development in last few decades: change the category, and work "stably," in some "stable" category:

- $\text{rep}(G) \rightsquigarrow \text{stab}(kG)$
- $\text{HTop} \rightsquigarrow \text{SH}^{\text{fin}}$
- $\text{R-Mod} \rightsquigarrow \text{D}^{\text{perf}}(R)$
- ...

These "stable" categories all have a tensor-triangulated category structure in common.

- Instead of classifying objects "on the nose," classify objects "up to the tensor-triangulated structure"
- Regard objects as being equivalent if they can build each other using the tt-structure
- Technically after a classification the thick tensor ideals

Balmer Support and the Classification Theorem

Some Historical Examples

Theorem (Benson-Carlson-Rickard 1997)

Let k be a field of char p dividing the order of a group G . There is a bijection between

Balmer Support and the Classification Theorem

Some Historical Examples

Theorem (Benson-Carlson-Rickard 1997)

Let k be a field of char p dividing the order of a group G . There is a bijection between

$$\{\text{Thick } \otimes\text{-ideals of } \text{stab}(kG)\} \xrightarrow{\sim} \{\text{Specialization Closed subsets of } \text{Proj}(H^\bullet(G, k))\}$$

$$\mathcal{C} \rightarrow \bigcup_{x \in \mathcal{C}} V_G(x)$$

Balmer Support and the Classification Theorem

Some Historical Examples

Theorem (Benson-Carlson-Rickard 1997)

Let k be a field of char p dividing the order of a group G . There is a bijection between

$$\{\text{Thick } \otimes\text{-ideals of } \text{stab}(kG)\} \xrightarrow{\sim} \{\text{Specialization Closed subsets of } \text{Proj}(H^\bullet(G, k))\}$$

$$\mathcal{C} \rightarrow \bigcup_{x \in \mathcal{C}} V_G(x)$$

Theorem (Neeman-Thomason)

Let R be a commutative ring. There is a bijection

Balmer Support and the Classification Theorem

Some Historical Examples

Theorem (Benson-Carlson-Rickard 1997)

Let k be a field of char p dividing the order of a group G . There is a bijection between

$$\{\text{Thick } \otimes\text{-ideals of } \text{stab}(kG)\} \xrightarrow{\sim} \{\text{Specialization Closed subsets of } \text{Proj}(H^\bullet(G, k))\}$$
$$\mathcal{C} \rightarrow \bigcup_{x \in \mathcal{C}} V_G(x)$$

Theorem (Neeman-Thomason)

Let R be a commutative ring. There is a bijection

$$\{\text{Thick } \otimes\text{-ideals of } D^{\text{perf}}(R)\} \xrightarrow{\sim} \{\text{Thomason Subsets of } \text{Spec}(R)\}$$
$$\mathcal{C} \rightarrow \bigcup_{x \in \mathcal{C}} \text{supph}(x)$$

Balmer Support and the Classification Theorem

Support Theories

- These classifications were accomplished using various notions of "support"
- Greatly clarified and unified via the work of Paul Balmer (2005)

Support Theories

- These classifications were accomplished using various notions of "support"
- Greatly clarified and unified via the work of Paul Balmer (2005)
 - Given essentially small tt-category \mathcal{K} get topological space

$$\mathrm{Spc}(\mathcal{K}) = \{\mathcal{P} \subseteq \mathcal{K} : \mathcal{P} \text{ is a prime ideal}\}$$

Support Theories

- These classifications were accomplished using various notions of "support"
- Greatly clarified and unified via the work of Paul Balmer (2005)
 - Given essentially small tt-category \mathcal{K} get topological space

$$\mathrm{Spc}(\mathcal{K}) = \{\mathcal{P} \subseteq \mathcal{K} : \mathcal{P} \text{ is a prime ideal}\}$$

- Equipped with a closed subset $\mathrm{supp}(x) \subseteq \mathrm{Spc}(\mathcal{K})$ for all $x \in \mathcal{K}$

Balmer Support and the Classification Theorem

Support Theories

- These classifications were accomplished using various notions of "support"
- Greatly clarified and unified via the work of Paul Balmer (2005)
 - Given essentially small tt-category \mathcal{K} get topological space

$$\mathrm{Spc}(\mathcal{K}) = \{\mathcal{P} \subseteq \mathcal{K} : \mathcal{P} \text{ is a prime ideal}\}$$

- Equipped with a closed subset $\mathrm{supp}(x) \subseteq \mathrm{Spc}(\mathcal{K})$ for all $x \in \mathcal{K}$
- The pair $(\mathrm{Spc}(\mathcal{K}), \mathrm{supp})$ is the universal space with well-behaved notion of support

Classification Theorem and Examples

For mild assumptions on \mathcal{K} there is a bijection

$$\{\text{Thick } \otimes\text{-ideals of } \mathcal{K}\} \xrightarrow{\sim} \{\text{Thomason Subsets of } \text{Spc}(\mathcal{K})\}$$

$$\mathcal{C} \rightarrow \bigcup_{x \in \mathcal{C}} \text{supp}(x)$$

Classification Theorem and Examples

For mild assumptions on \mathcal{K} there is a bijection

$$\{\text{Thick } \otimes\text{-ideals of } \mathcal{K}\} \xrightarrow{\sim} \{\text{Thomason Subsets of } \text{Spc}(\mathcal{K})\}$$

$$C \rightarrow \bigcup_{x \in C} \text{supp}(x)$$

- Thomason subset = union of closed subsets with quasi-compact complement.

Classification Theorem and Examples

For mild assumptions on \mathcal{K} there is a bijection

$$\{\text{Thick } \otimes\text{-ideals of } \mathcal{K}\} \xrightarrow{\sim} \{\text{Thomason Subsets of } \text{Spc}(\mathcal{K})\}$$

$$C \rightarrow \bigcup_{x \in C} \text{supp}(x)$$

- Thomason subset = union of closed subsets with quasi-compact complement. (In noetherian context, Thomason = specialization closed)

Classification Theorem and Examples

For mild assumptions on \mathcal{K} there is a bijection

$$\begin{aligned} \{\text{Thick } \otimes\text{-ideals of } \mathcal{K}\} &\xrightarrow{\sim} \{\text{Thomason Subsets of } \text{Spc}(\mathcal{K})\} \\ C &\rightarrow \bigcup_{x \in C} \text{supp}(x) \end{aligned}$$

- Thomason subset = union of closed subsets with quasi-compact complement. (In noetherian context, Thomason = specialization closed)
- $\langle x \rangle = \langle y \rangle \iff \text{supp}(x) = \text{supp}(y)$

Balmer Support and the Classification Theorem

Classification Theorem and Examples

For mild assumptions on \mathcal{K} there is a bijection

$$\begin{aligned} \{\text{Thick } \otimes\text{-ideals of } \mathcal{K}\} &\xrightarrow{\sim} \{\text{Thomason Subsets of } \text{Spc}(\mathcal{K})\} \\ \mathcal{C} &\rightarrow \bigcup_{x \in \mathcal{C}} \text{supp}(x) \end{aligned}$$

- Thomason subset = union of closed subsets with quasi-compact complement. (In noetherian context, Thomason = specialization closed)
- $\langle x \rangle = \langle y \rangle \iff \text{supp}(x) = \text{supp}(y)$

Goal: Determine the Balmer spectrum for various tt-categories- and therefore deduce the classification in that context:

Classification Theorem and Examples

For mild assumptions on \mathcal{K} there is a bijection

$$\{\text{Thick } \otimes\text{-ideals of } \mathcal{K}\} \xrightarrow{\sim} \{\text{Thomason Subsets of } \text{Spc}(\mathcal{K})\}$$
$$C \rightarrow \bigcup_{x \in C} \text{supp}(x)$$

- Thomason subset = union of closed subsets with quasi-compact complement. (In noetherian context, Thomason = specialization closed)
- $\langle x \rangle = \langle y \rangle \iff \text{supp}(x) = \text{supp}(y)$

Goal: Determine the Balmer spectrum for various tt-categories- and therefore deduce the classification in that context:

- $\text{Spc}(\text{stab}(kG)) \cong \text{Proj}(H^\bullet(G, k))$

Balmer Support and the Classification Theorem

Classification Theorem and Examples

For mild assumptions on \mathcal{K} there is a bijection

$$\{\text{Thick } \otimes\text{-ideals of } \mathcal{K}\} \xrightarrow{\sim} \{\text{Thomason Subsets of } \text{Spc}(\mathcal{K})\}$$
$$C \rightarrow \bigcup_{x \in C} \text{supp}(x)$$

- Thomason subset = union of closed subsets with quasi-compact complement. (In noetherian context, Thomason = specialization closed)
- $\langle x \rangle = \langle y \rangle \iff \text{supp}(x) = \text{supp}(y)$

Goal: Determine the Balmer spectrum for various tt-categories- and therefore deduce the classification in that context:

- $\text{Spc}(\text{stab}(kG)) \cong \text{Proj}(H^\bullet(G, k))$
- $\text{Spc}(D^{\text{perf}}(R)) \cong \text{Spec}(R)$
- ...

Limiting factor of Balmer Support

- Many interesting tt-categories are really some "finite" piece of some "larger" tt-category

Limiting factor of Balmer Support

- Many interesting tt-categories are really some "finite" piece of some "larger" tt-category
 - $\text{stab}(kG) \subset \text{Stab}(kG)$
 - $\text{SH}^{\text{fin}} \subset \text{SH}$
 - $D^{\text{perf}}(R) \subset D(R)$

Limiting factor of Balmer Support

- Many interesting tt-categories are really some "finite" piece of some "larger" tt-category
 - $\text{stab}(kG) \subset \text{Stab}(kG)$
 - $\text{SH}^{\text{fin}} \subset \text{SH}$
 - $D^{\text{perf}}(R) \subset D(R)$
- This "finite" idea can be made precise via the concept of "compact objects"

Limiting factor of Balmer Support

- Many interesting tt-categories are really some "finite" piece of some "larger" tt-category
 - $\text{stab}(kG) \subset \text{Stab}(kG)$
 - $\text{SH}^{\text{fin}} \subset \text{SH}$
 - $D^{\text{perf}}(R) \subset D(R)$
- This "finite" idea can be made precise via the concept of "compact objects"
- The large, non-compact objects, however, are very interesting!
 - Objects representing cohomology theories are not compact in SH

Limiting factor of Balmer Support

- Many interesting tt-categories are really some "finite" piece of some "larger" tt-category
 - $\text{stab}(kG) \subset \text{Stab}(kG)$
 - $\text{SH}^{\text{fin}} \subset \text{SH}$
 - $D^{\text{perf}}(R) \subset D(R)$
- This "finite" idea can be made precise via the concept of "compact objects"
- The large, non-compact objects, however, are very interesting!
 - Objects representing cohomology theories are not compact in SH
 - There are major open questions about the structure of the larger objects (e.g. The telescope conjecture)

How to Classify the Big Objects

- Goal = understand the structure of a "large" tt-category \mathcal{T}

How to Classify the Big Objects

- Goal = understand the structure of a "large" tt-category \mathcal{T}
- Similar to the compact case, after a classification of the "localizing tensor ideals"

How to Classify the Big Objects

- Goal = understand the structure of a "large" tt-category \mathcal{T}
- Similar to the compact case, after a classification of the "localizing tensor ideals"
- In this case, there is no "Balmer spectrum" for \mathcal{T} - less clear how to define a "universal" notion of support for non-compact objects

How to Classify the Big Objects

- Goal = understand the structure of a "large" tt-category \mathcal{T}
- Similar to the compact case, after a classification of the "localizing tensor ideals"
- In this case, there is no "Balmer spectrum" for \mathcal{T} - less clear how to define a "universal" notion of support for non-compact objects
 - e.g.) For x compact, $\text{supp}(x)$ will be closed, but this is not expected for non-compact objects

How to Classify the Big Objects

- Goal = understand the structure of a "large" tt-category \mathcal{T}
- Similar to the compact case, after a classification of the "localizing tensor ideals"
- In this case, there is no "Balmer spectrum" for \mathcal{T} - less clear how to define a "universal" notion of support for non-compact objects
 - e.g.) For x compact, $\text{supp}(x)$ will be closed, but this is not expected for non-compact objects
- Nonetheless, been some success using novel theories of support to develop such classifications in noetherian contexts (BIK, Neemen, ect)

How to Classify the Big Objects

- Goal = understand the structure of a "large" tt-category \mathcal{T}
- Similar to the compact case, after a classification of the "localizing tensor ideals"
- In this case, there is no "Balmer spectrum" for \mathcal{T} - less clear how to define a "universal" notion of support for non-compact objects
 - e.g.) For x compact, $\text{supp}(x)$ will be closed, but this is not expected for non-compact objects
- Nonetheless, been some success using novel theories of support to develop such classifications in noetherian contexts (BIK, Neemen, ect)
- In recent work Barthel, Heard and Sanders (BHS, 2021) developed a support theory for noetherian large tt-categories \mathcal{T}

How to Classify the Big Objects

- Goal = understand the structure of a "large" tt-category \mathcal{T}
- Similar to the compact case, after a classification of the "localizing tensor ideals"
- In this case, there is no "Balmer spectrum" for \mathcal{T} - less clear how to define a "universal" notion of support for non-compact objects
 - e.g.) For x compact, $\text{supp}(x)$ will be closed, but this is not expected for non-compact objects
- Nonetheless, been some success using novel theories of support to develop such classifications in noetherian contexts (BIK, Neemen, ect)
- In recent work Barthel, Heard and Sanders (BHS, 2021) developed a support theory for noetherian large tt-categories \mathcal{T}
 - There is no " $\text{Spc}(\mathcal{T})$ " but can consider $\text{Spc}(\mathcal{T}^c)$
 - The support for arbitrary objects will be a subset of $\text{Spc}(\mathcal{T}^c)$

Stratification

Sanders, et al use this support to show that for many noetherian tt-categories there is a bijection

Stratification

Sanders, et al use this support to show that for many noetherian tt-categories there is a bijection

$$\begin{aligned} \{\text{Localizing } \otimes\text{-ideals of } \mathcal{T}\} &\xrightarrow{\sim} \{\text{Arbitrary subsets of } \mathrm{Spc}(\mathcal{T}^c)\} \\ \mathcal{L} &\rightarrow \bigcup_{t \in \mathcal{L}} \mathrm{Supp}(t) \end{aligned}$$

When such a category admits this bijection, they say the category is *stratified*.

Stratification

Sanders, et al use this support to show that for many noetherian tt-categories there is a bijection

$$\{\text{Localizing } \otimes\text{-ideals of } \mathcal{T}\} \xrightarrow{\sim} \{\text{Arbitrary subsets of } \text{Spc}(\mathcal{T}^c)\}$$
$$\mathcal{L} \rightarrow \bigcup_{t \in \mathcal{L}} \text{Supp}(t)$$

When such a category admits this bijection, they say the category is *stratified*.

- Develop a framework for deducing such a bijection

Stratification

Sanders, et al use this support to show that for many noetherian tt-categories there is a bijection

$$\{\text{Localizing } \otimes\text{-ideals of } \mathcal{T}\} \xrightarrow{\sim} \{\text{Arbitrary subsets of } \text{Spc}(\mathcal{T}^c)\}$$
$$\mathcal{L} \rightarrow \bigcup_{t \in \mathcal{L}} \text{Supp}(t)$$

When such a category admits this bijection, they say the category is *stratified*.

- Develop a framework for deducing such a bijection
- Involves showing a certain minimality condition at each prime \mathcal{P}

Stratification

Sanders, et al use this support to show that for many noetherian tt-categories there is a bijection

$$\{\text{Localizing } \otimes\text{-ideals of } \mathcal{T}\} \xrightarrow{\sim} \{\text{Arbitrary subsets of } \text{Spc}(\mathcal{T}^c)\}$$
$$\mathcal{L} \rightarrow \bigcup_{t \in \mathcal{L}} \text{Supp}(t)$$

When such a category admits this bijection, they say the category is *stratified*.

- Develop a framework for deducing such a bijection
- Involves showing a certain minimality condition at each prime \mathcal{P}
- **Important point:** Can work "locally" at each prime \mathcal{P}

Many noetherian categories are stratified in this sense:

Stratification

Sanders, et al use this support to show that for many noetherian tt-categories there is a bijection

$$\{\text{Localizing } \otimes\text{-ideals of } \mathcal{T}\} \xrightarrow{\sim} \{\text{Arbitrary subsets of } \text{Spc}(\mathcal{T}^c)\}$$
$$\mathcal{L} \rightarrow \bigcup_{t \in \mathcal{L}} \text{Supp}(t)$$

When such a category admits this bijection, they say the category is *stratified*.

- Develop a framework for deducing such a bijection
- Involves showing a certain minimality condition at each prime \mathcal{P}
- **Important point:** Can work "locally" at each prime \mathcal{P}

Many noetherian categories are stratified in this sense:

- $\text{Stab}(kG)$ the large stable module category;

Stratification

Sanders, et al use this support to show that for many noetherian tt-categories there is a bijection

$$\{\text{Localizing } \otimes\text{-ideals of } \mathcal{T}\} \xrightarrow{\sim} \{\text{Arbitrary subsets of } \text{Spc}(\mathcal{T}^c)\}$$
$$\mathcal{L} \rightarrow \bigcup_{t \in \mathcal{L}} \text{Supp}(t)$$

When such a category admits this bijection, they say the category is *stratified*.

- Develop a framework for deducing such a bijection
- Involves showing a certain minimality condition at each prime \mathcal{P}
- **Important point:** Can work "locally" at each prime \mathcal{P}

Many noetherian categories are stratified in this sense:

- $\text{Stab}(kG)$ the large stable module category;
- $D(R)$ the unbounded derived category for R noetherian ring (fails spectacularly for R not noetherian);

Stratification

Sanders, et al use this support to show that for many noetherian tt-categories there is a bijection

$$\{\text{Localizing } \otimes\text{-ideals of } \mathcal{T}\} \xrightarrow{\sim} \{\text{Arbitrary subsets of } \text{Spc}(\mathcal{T}^c)\}$$
$$\mathcal{L} \rightarrow \bigcup_{t \in \mathcal{L}} \text{Supp}(t)$$

When such a category admits this bijection, they say the category is *stratified*.

- Develop a framework for deducing such a bijection
- Involves showing a certain minimality condition at each prime \mathcal{P}
- **Important point:** Can work "locally" at each prime \mathcal{P}

Many noetherian categories are stratified in this sense:

- $\text{Stab}(\text{kG})$ the large stable module category;
- $\text{D}(\text{R})$ the unbounded derived category for R noetherian ring (fails spectacularly for R not noetherian);
- $\text{D}(\text{X})$ for X noetherian scheme
- ...

Some Historical Comments

- "Motives" were originated by Grothendieck and his students in the 1960s in their study of algebraic cycles and cohomology theories in algebraic geometry

Some Historical Comments

- "Motives" were originated by Grothendieck and his students in the 1960s in their study of algebraic cycles and cohomology theories in algebraic geometry
- Idea: should be a "universal cohomology theory" (motivic cohomology theory) in algebraic geometry

Some Historical Comments

- "Motives" were originated by Grothendieck and his students in the 1960s in their study of algebraic cycles and cohomology theories in algebraic geometry
- Idea: should be a "universal cohomology theory" (motivic cohomology theory) in algebraic geometry
- Not much success in constructing this abelian category of motives

Motives and their Derived Categories

Some Historical Comments

- "Motives" were originated by Grothendieck and his students in the 1960s in their study of algebraic cycles and cohomology theories in algebraic geometry
- Idea: should be a "universal cohomology theory" (motivic cohomology theory) in algebraic geometry
- Not much success in constructing this abelian category of motives
- Instead Voevodsky was able to construct a "derived category" of motives, denoted $DM(\mathbb{F})$

Motives and their Derived Categories

Some Historical Comments

- "Motives" were originated by Grothendieck and his students in the 1960s in their study of algebraic cycles and cohomology theories in algebraic geometry
- Idea: should be a "universal cohomology theory" (motivic cohomology theory) in algebraic geometry
- Not much success in constructing this abelian category of motives
- Instead Voevodsky was able to construct a "derived category" of motives, denoted $DM(\mathbb{F})$
- This theory greatly increased our understanding of motivic cohomology.

Motives and their Derived Categories

Some Historical Comments

- "Motives" were originated by Grothendieck and his students in the 1960s in their study of algebraic cycles and cohomology theories in algebraic geometry
- Idea: should be a "universal cohomology theory" (motivic cohomology theory) in algebraic geometry
- Not much success in constructing this abelian category of motives
- Instead Voevodsky was able to construct a "derived category" of motives, denoted $DM(\mathbb{F})$
- This theory greatly increased our understanding of motivic cohomology.
- Prove some longstanding fundamental conjectures in algebraic geometry (e.g. The Milnor conjecture and the Bloch-Kato conjecture).

Necessary Facts about $DM(\mathbb{F}, R)$

- $DM(\mathbb{F}, R)$ is a "large" tensor triangulated category and there is an associated "motive functor"

Necessary Facts about $DM(\mathbb{F}, R)$

- $DM(\mathbb{F}, R)$ is a "large" tensor triangulated category and there is an associated "motive functor"

$$R : S_{\mathbb{m}}/\mathbb{F} \rightarrow DM(\mathbb{F}, R)$$

- There are invertible objects $R(n) \in DM(\mathbb{F}, R)$ for each $n \in \mathbb{Z}$, such that $R(n) \otimes R(m) = R(n + m)$ called the Tate Twists.

Necessary Facts about $DM(\mathbb{F}, \mathbb{R})$

- $DM(\mathbb{F}, \mathbb{R})$ is a "large" tensor triangulated category and there is an associated "motive functor"

$$R : \text{Sm}/\mathbb{F} \rightarrow DM(\mathbb{F}, \mathbb{R})$$

- There are invertible objects $R(n) \in DM(\mathbb{F}, \mathbb{R})$ for each $n \in \mathbb{Z}$, such that $R(n) \otimes R(m) = R(n + m)$ called the Tate Twists.
- The motivic cohomology groups for a variety X are then defined as

$$H^{m,n}(X) = \text{hom}_{DM(\mathbb{F}, \mathbb{R})}(R(X), R(n)[m])$$

Necessary Facts about $DM(\mathbb{F}, \mathbb{R})$

- $DM(\mathbb{F}, \mathbb{R})$ is a "large" tensor triangulated category and there is an associated "motive functor"

$$R : \text{Sm}/\mathbb{F} \rightarrow DM(\mathbb{F}, \mathbb{R})$$

- There are invertible objects $R(n) \in DM(\mathbb{F}, \mathbb{R})$ for each $n \in \mathbb{Z}$, such that $R(n) \otimes R(m) = R(n + m)$ called the Tate Twists.
- The motivic cohomology groups for a variety X are then defined as

$$H^{m,n}(X) = \text{hom}_{DM(\mathbb{F}, \mathbb{R})}(R(X), R(n)[m])$$

The category $DM(\mathbb{F}, \mathbb{R})$ is extremely complex

Motives and their Derived Categories

Necessary Facts about $DM(\mathbb{F}, \mathbb{R})$

- $DM(\mathbb{F}, \mathbb{R})$ is a "large" tensor triangulated category and there is an associated "motive functor"

$$R : \text{Sm}/\mathbb{F} \rightarrow DM(\mathbb{F}, \mathbb{R})$$

- There are invertible objects $R(n) \in DM(\mathbb{F}, \mathbb{R})$ for each $n \in \mathbb{Z}$, such that $R(n) \otimes R(m) = R(n + m)$ called the Tate Twists.
- The motivic cohomology groups for a variety X are then defined as

$$H^{m,n}(X) = \text{hom}_{DM(\mathbb{F}, \mathbb{R})}(R(X), R(n)[m])$$

The category $DM(\mathbb{F}, \mathbb{R})$ is extremely complex

- Idea: study first a "piece" of the category; the Tate motives
- The localizing subcategory generated by the Tate twists is the (large) category of Tate motives, denoted by $DTM(\mathbb{F}, \mathbb{R})$.

Étale Motives

There is a similar story to tell with so called étale motives.

- There is a (large) derived category of étale motives, denoted $DM^{\text{ét}}(\mathbb{F}, \mathbb{R})$, again with an étale motive functor:

Étale Motives

There is a similar story to tell with so called étale motives.

- There is a (large) derived category of étale motives, denoted $DM^{\text{ét}}(\mathbb{F}, \mathbb{R})$, again with an étale motive functor:

$$R^{\text{ét}} : \mathbf{Sm}/\mathbb{F} \rightarrow DM^{\text{ét}}(\mathbb{F}, \mathbb{R})$$

- As before, there are invertible objects $R^{\text{ét}}(n)$ satisfying the same tensor product formula

Étale Motives

There is a similar story to tell with so called étale motives.

- There is a (large) derived category of étale motives, denoted $DM^{\text{ét}}(\mathbb{F}, \mathbb{R})$, again with an étale motive functor:

$$R^{\text{ét}} : \mathbf{Sm}/\mathbb{F} \rightarrow DM^{\text{ét}}(\mathbb{F}, \mathbb{R})$$

- As before, there are invertible objects $R^{\text{ét}}(n)$ satisfying the same tensor product formula
- Again first study $DTM^{\text{ét}}(\mathbb{F}, \mathbb{R})$, the localizing subcategory generated by étale Tate twists

Étale Motives

There is a similar story to tell with so called étale motives.

- There is a (large) derived category of étale motives, denoted $DM^{\text{ét}}(\mathbb{F}, \mathbb{R})$, again with an étale motive functor:

$$R^{\text{ét}} : \mathbf{Sm}/\mathbb{F} \rightarrow DM^{\text{ét}}(\mathbb{F}, \mathbb{R})$$

- As before, there are invertible objects $R^{\text{ét}}(n)$ satisfying the same tensor product formula
- Again first study $DTM^{\text{ét}}(\mathbb{F}, \mathbb{R})$, the localizing subcategory generated by étale Tate twists
- There are analogous étale motivic cohomology groups for a variety X

Étale Motives

There is a similar story to tell with so called étale motives.

- There is a (large) derived category of étale motives, denoted $DM^{\text{ét}}(\mathbb{F}, \mathbb{R})$, again with an étale motive functor:

$$R^{\text{ét}} : \text{Sm}/\mathbb{F} \rightarrow DM^{\text{ét}}(\mathbb{F}, \mathbb{R})$$

- As before, there are invertible objects $R^{\text{ét}}(n)$ satisfying the same tensor product formula
- Again first study $DTM^{\text{ét}}(\mathbb{F}, \mathbb{R})$, the localizing subcategory generated by étale Tate twists
- There are analogous étale motivic cohomology groups for a variety X

These two constructions are very similar: there is an "étale sheafification" functor

$$\alpha_{\text{ét}} : DM(\mathbb{F}, \mathbb{R}) \rightarrow DM^{\text{ét}}(\mathbb{F}, \mathbb{R})$$

Moreover,

Étale Motives

There is a similar story to tell with so called étale motives.

- There is a (large) derived category of étale motives, denoted $DM^{\text{ét}}(\mathbb{F}, \mathbb{R})$, again with an étale motive functor:

$$R^{\text{ét}} : \text{Sm}/\mathbb{F} \rightarrow DM^{\text{ét}}(\mathbb{F}, \mathbb{R})$$

- As before, there are invertible objects $R^{\text{ét}}(n)$ satisfying the same tensor product formula
- Again first study $DTM^{\text{ét}}(\mathbb{F}, \mathbb{R})$, the localizing subcategory generated by étale Tate twists
- There are analogous étale motivic cohomology groups for a variety X

These two constructions are very similar: there is an "étale sheafification" functor

$$\alpha_{\text{ét}} : DM(\mathbb{F}, \mathbb{R}) \rightarrow DM^{\text{ét}}(\mathbb{F}, \mathbb{R})$$

Moreover,

- Whenever $\mathbb{Q} \subset \mathbb{R}$, $\alpha_{\text{ét}}$ is an equivalence of categories.

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

The Balmer spectrum of $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c$

Theorem (Gallauer 2019)


(1) *The Balmer spectrum of $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c$ is the following picture:*

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

The Balmer spectrum of $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c$

Theorem (Gallauer 2019)

(1) *The Balmer spectrum of $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c$ is the following picture:*



In this picture the specialization relations are pointing upwards.

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

The Balmer spectrum of $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c$

Theorem (Gallauer 2019)

(1) *The Balmer spectrum of $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c$ is the following picture:*

$$\mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

In this picture the specialization relations are pointing upwards.

(2) $\mathrm{Spc}(\mathrm{DTM}^{\mathrm{\acute{e}t}}(\overline{\mathbb{Q}}, \mathbb{Z})^c) \cong \mathrm{Spec}(\mathbb{Z})$

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

The Balmer spectrum of $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c$

Theorem (Gallauer 2019)

- (1) *The Balmer spectrum of $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c$ is the following picture:*

$$\mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

In this picture the specialization relations are pointing upwards.

- (2) $\mathrm{Spc}(\mathrm{DTM}^{\mathrm{\acute{e}t}}(\overline{\mathbb{Q}}, \mathbb{Z})^c) \cong \mathrm{Spec}(\mathbb{Z})$
- (3) *The étale sheafification map induces a map $\mathrm{Spec}(\mathbb{Z}) \xrightarrow{\mathrm{Spc}(\alpha_{\acute{e}t})} \mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c)$ which is a homeomorphism onto the subspace $\{m_o, e_p\}$*

My Problem: Stratification for $\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

Remarks on These Computations

$$\text{Spc}(\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) =$$

```
graph TD
    m2[m2] --- e2[e2]
    m3[m3] --- e3[e3]
    mp[mp] --- ep[ep]
    e2 --> m0[m0]
    e3 --> m0
    mp --> m0
```

Let us explain what these m_p, e_p, m_0 are:

My Problem: Stratification for $\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

Remarks on These Computations

$$\text{Spc}(\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

Let us explain what these m_p, e_p, m_0 are:

(m_p) : kernel of map $\gamma_p^* : \text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c \rightarrow \text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c$

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

Remarks on These Computations

$$\mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccccc} m_2 & m_3 & & \cdots & m_p & \cdots \\ | & | & & & | & \\ e_2 & e_3 & & \cdots & e_p & \cdots \\ & \searrow & & & \nearrow & \\ & m_0 & & & & \end{array}$$

Let us explain what these m_p, e_p, m_0 are:

(m_p) : kernel of map $\gamma_p^* : \mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c \rightarrow \mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c$. A result of Gallauer is that this kernel coincides with those motives whose mod- p motivic cohomology vanishes.

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

Remarks on These Computations

$$\mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) =$$

Let us explain what these m_p, e_p, m_0 are:

(m_p) : kernel of map $\gamma_p^* : \mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c \rightarrow \mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c$. A result of Gallauer is that this kernel coincides with those motives whose mod- p motivic cohomology vanishes.

(e_p) : kernels of the composite

$$\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c \xrightarrow{\gamma^*} \mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c \xrightarrow{\alpha_{\text{ét}}} \mathrm{DTM}^{\text{ét}}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c$$

Again, this kernel coincides with those motives whose mod- p étale cohomology vanishes.

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

Remarks on These Computations

$$\mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

Let us explain what these m_p, e_p, m_0 are:

(m_p) : kernel of map $\gamma_p^* : \mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c \rightarrow \mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c$. A result of Gallauer is that this kernel coincides with those motives whose mod- p motivic cohomology vanishes.

(e_p) : kernels of the composite

$$\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c \xrightarrow{\gamma^*} \mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c \xrightarrow{\alpha_{\text{ét}}} \mathrm{DTM}^{\text{ét}}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c$$

Again, this kernel coincides with those motives whose mod- p étale cohomology vanishes.

(m_0) : kernel of the rationalization map $\gamma^* : \mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c \rightarrow \mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Q})^c$, which coincides with those motives whose rational motivic cohomology vanishes.

My Problem: Stratification for $\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

Remarks on These Computations

$$\text{Spc}(\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

- Morally there are only 3 flavors of primes in $\text{Spc}(\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c)$

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

Remarks on These Computations

$$\mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

- Morally there are only 3 flavors of primes in $\mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c)$
- Reduce the problem to each "vertical slice" in the spectrum and just consider the 3 primes in each slice

My Problem: Stratification for $\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

How To Establish Minimality at the Primes

$$\text{Spc}(\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{cccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

(m_p) : Suffices to pass to the "residue field" $\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})$.

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

How To Establish Minimality at the Primes

$$\mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

(m_p) : Suffices to pass to the "residue field" $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})$.

- Gallauer proves that $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c \cong D_{\mathrm{fil}}^b(\mathbb{Z}/p\mathbb{Z})$ is equivalent to the filtered bounded derived category of $\mathbb{Z}/p\mathbb{Z}$

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

How To Establish Minimality at the Primes

$$\mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

(m_p) : Suffices to pass to the "residue field" $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})$.

- Gallauer proves that $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c \cong D_{\mathrm{fil}}^b(\mathbb{Z}/p\mathbb{Z})$ is equivalent to the filtered bounded derived category of $\mathbb{Z}/p\mathbb{Z}$
- Does this lift to the large categories?

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

How To Establish Minimality at the Primes

$$\mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

(m_p) : Suffices to pass to the "residue field" $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})$.

- Gallauer proves that $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c \cong D_{\mathrm{fil}}^b(\mathbb{Z}/p\mathbb{Z})$ is equivalent to the filtered bounded derived category of $\mathbb{Z}/p\mathbb{Z}$
- Does this lift to the large categories?

(e_p) : Have successfully shown minimality for these

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

How To Establish Minimality at the Primes

$$\mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

(m_p) : Suffices to pass to the "residue field" $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})$.

- Gallauer proves that $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c \cong D_{\mathrm{fil}}^b(\mathbb{Z}/p\mathbb{Z})$ is equivalent to the filtered bounded derived category of $\mathbb{Z}/p\mathbb{Z}$
- Does this lift to the large categories?

(e_p) : Have successfully shown minimality for these

- Gallauer showed the local categories at the primes e_p are just $\mathrm{DTM}^{\mathrm{ét}}(\overline{\mathbb{Q}}, \mathbb{Z}_p)$

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

How To Establish Minimality at the Primes

$$\mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

(m_p) : Suffices to pass to the "residue field" $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})$.

- Gallauer proves that $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c \cong D_{\mathrm{fil}}^b(\mathbb{Z}/p\mathbb{Z})$ is equivalent to the filtered bounded derived category of $\mathbb{Z}/p\mathbb{Z}$
- Does this lift to the large categories?

(e_p) : Have successfully shown minimality for these

- Gallauer showed the local categories at the primes e_p are just $\mathrm{DTM}^{\mathrm{\acute{e}t}}(\overline{\mathbb{Q}}, \mathbb{Z}_p)$
- Again pass to the residue field $\mathrm{DTM}^{\mathrm{\acute{e}t}}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})$. The Rigidity Theorem establishes stratification for this category.

My Problem: Stratification for $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

How To Establish Minimality at the Primes

$$\mathrm{Spc}(\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

(m_p) : Suffices to pass to the "residue field" $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})$.

- Gallauer proves that $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})^c \cong D_{\mathrm{fil}}^b(\mathbb{Z}/p\mathbb{Z})$ is equivalent to the filtered bounded derived category of $\mathbb{Z}/p\mathbb{Z}$
- Does this lift to the large categories?

(e_p) : Have successfully shown minimality for these

- Gallauer showed the local categories at the primes e_p are just $\mathrm{DTM}^{\mathrm{\acute{e}t}}(\overline{\mathbb{Q}}, \mathbb{Z}_p)$
- Again pass to the residue field $\mathrm{DTM}^{\mathrm{\acute{e}t}}(\overline{\mathbb{Q}}, \mathbb{Z}/p\mathbb{Z})$. The Rigidity Theorem establishes stratification for this category.

(m_0) : Reduces to showing minimality in $\mathrm{DTM}(\overline{\mathbb{Q}}, \mathbb{Q})$.

My Problem: Stratification for $\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

Final Comments and Summery

$$\text{Spc}(\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccccc} m_2 & m_3 & & \cdots & m_p & \cdots \\ | & | & & & | & \\ e_2 & e_3 & & \cdots & e_p & \cdots \\ & \searrow & \swarrow & & \swarrow & \\ & & m_0 & & & \end{array}$$

- In summary, we want to get a classification for the localizing tensor ideals for $\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$.

My Problem: Stratification for $\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

Final Comments and Summery

$$\text{Spc}(\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccccc} m_2 & m_3 & & \cdots & m_p & \cdots \\ | & | & & & | & \\ e_2 & e_3 & & \cdots & e_p & \cdots \\ & \searrow & \swarrow & & \swarrow & \\ & & m_0 & & & \end{array}$$

- In summary, we want to get a classification for the localizing tensor ideals for $\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$.
- Using the results of Sanders, et al we are tasked with checking a certain minimality condition at every prime.

My Problem: Stratification for $\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$

Final Comments and Summery

$$\text{Spc}(\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})^c) = \begin{array}{ccccccc} m_2 & m_3 & \cdots & m_p & \cdots \\ | & | & & | & \\ e_2 & e_3 & \cdots & e_p & \cdots \\ & \searrow & & \nearrow & \\ & m_0 & & & \end{array}$$

- In summary, we want to get a classification for the localizing tensor ideals for $\text{DTM}(\overline{\mathbb{Q}}, \mathbb{Z})$.
- Using the results of Sanders, et al we are tasked with checking a certain minimality condition at every prime.
- In this case, we can first take vertical slices of the spectrum, and then check minimality at local categories for mod p and rational coefficients