

An Introduction into Building Theory

(Using Abstract Simplicial Complexes and Coxeter Groups)

Malachi Alexander

UC Santa Cruz

May 20, 2021

Table of Contents

- 1 Motivation
- 2 Basics
 - Finite Reflection Groups
 - Coxeter Complex
- 3 Examples

Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field.

Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).

Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).
- (2) A collection of chambers together with a Weyl-group valued distance function.

Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).
- (2) A collection of chambers together with a Weyl-group valued distance function.
- (3) The apartments in the first perspective are viewed as metric spaces rather than simplicial complexes with specific geometric properties.

Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).
- (2) A collection of chambers together with a Weyl-group valued distance function.
- (3) The apartments in the first perspective are viewed as metric spaces rather than simplicial complexes with specific geometric properties.

The perspectives differ in how one views the chambers:

Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).
- (2) A collection of chambers together with a Weyl-group valued distance function.
- (3) The apartments in the first perspective are viewed as metric spaces rather than simplicial complexes with specific geometric properties.

The perspectives differ in how one views the chambers:

- (1) Chambers are maximal simplices.

Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).
- (2) A collection of chambers together with a Weyl-group valued distance function.
- (3) The apartments in the first perspective are viewed as metric spaces rather than simplicial complexes with specific geometric properties.

The perspectives differ in how one views the chambers:

- (1) Chambers are maximal simplices.
- (2) Chambers are elements of an abstract set.

Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).
- (2) A collection of chambers together with a Weyl-group valued distance function.
- (3) The apartments in the first perspective are viewed as metric spaces rather than simplicial complexes with specific geometric properties.

The perspectives differ in how one views the chambers:

- (1) Chambers are maximal simplices.
- (2) Chambers are elements of an abstract set.
- (3) Chambers are metric spaces.

Motivation and Perspectives

In each perspective, the notion of Coxeter groups are required.

Motivation and Perspectives

In each perspective, the notion of Coxeter groups are required.

- We will use finite reflection groups (which characterize finite Coxeter groups).

Motivation and Perspectives

In each perspective, the notion of Coxeter groups are required.

- We will use finite reflection groups (which characterize finite Coxeter groups).
- We will present from the simplicial complex perspective.

Motivation and Perspectives

In each perspective, the notion of Coxeter groups are required.

- We will use finite reflection groups (which characterize finite Coxeter groups).
- We will present from the simplicial complex perspective.

Definition of a Building

A (*weak*) *building* is a simplicial complex Δ that can be expressed as the union of subcomplexes Σ (called *apartments*) satisfying axioms:

- (B0) Each apartment Σ is a Coxeter complex.
- (B1) For any two simplices $A, B \in \Delta$, there is an apartment Σ containing both of them.
- (B2) If Σ and Σ' are two apartments containing A and B , then there is an isomorphism $\Sigma \rightarrow \Sigma'$ fixing A and B pointwise.


Finite Reflection Groups

Let V be a Euclidean vector space, i.e. a finite-dimensional \mathbb{R} -vector space with an inner product and let V have dimension n .

Finite Reflection Groups

Let V be a Euclidean vector space, i.e. a finite-dimensional \mathbb{R} -vector space with an inner product and let V have dimension n .

Example: Let $V = \mathbb{R}^3$ equipped with the dot product.



Inner Product

$$\langle \mathbf{u}, \mathbf{v} \rangle = u_1 v_1 + u_2 v_2 + u_3 v_3$$


Finite Reflection Groups

Definition: A hyperplane H in V is a subspace of codimension 1 (we are assuming $\dim_{\mathbb{R}}(V) = n$, thus $\dim_{\mathbb{R}}(H) = n - 1$).

Finite Reflection Groups

Definition: A hyperplane H in V is a subspace of codimension 1 (we are assuming $\dim_{\mathbb{R}}(V) = n$, thus $\dim_{\mathbb{R}}(H) = n - 1$).

Example: Let $V = \mathbb{R}^3$ and $H_1 = \{\mathbf{u} \in \mathbb{R}^3 : \mathbf{u} = (u_1, 0, u_3)\}$.





Finite Reflection Groups

Definition: A reflection with respect to H is a linear transformation $s_H : V \rightarrow V$ such that s_H is the identity on H and multiplication by -1 along the orthogonal complement.

Finite Reflection Groups

Definition: A reflection with respect to H is a linear transformation $s_H : V \rightarrow V$ such that s_H is the identity on H and multiplication by -1 along the orthogonal complement.

Example: Let $u_1 = (1, 0, 1)$ and $u_2 = (1, 1, 1)$.



We obtain $s_{H_1}(1, 0, 1) = (1, 0, 1)$ and $s_{H_1}(1, 1, 1) = (1, -1, 1)$.

Finite Reflection Groups




Definition: Let \mathcal{H} be a set of hyperplanes in V , we call this a hyperplane arrangement. A (finite) reflection group W is a (finite) group generated by reflections s_H for $H \in \mathcal{H}$.

Finite Reflection Groups




Definition: Let \mathcal{H} be a set of hyperplanes in V , we call this a hyperplane arrangement. A (finite) reflection group W is a (finite) group generated by reflections s_H for $H \in \mathcal{H}$.

Example: Let $V = \mathbb{R}^3$, and


- $H_1 = \{\mathbf{u} \in \mathbb{R}^3 : \mathbf{u} = (u_1, 0, u_2)\}$
- $H_2 = \{\mathbf{u} \in \mathbb{R}^3 : \mathbf{u} = (u_1, \sqrt{3}u_2, u_2)\}$
- $H_3 = \{\mathbf{u} \in \mathbb{R}^3 : \mathbf{u} = (u_1, \sqrt{3}u_2, -u_2)\}$



Finite Reflection Groups




Finite Reflection Groups



We obtain the group: $W = \langle s_{H_1}, s_{H_2}, s_{H_3} \rangle$.


Finite Reflection Groups



We obtain the group: $W = \langle s_{H_1}, s_{H_2}, s_{H_3} \rangle$.

(1) For all i , $s_{H_i}^2 = \text{id}_V$


Finite Reflection Groups



We obtain the group: $W = \langle s_{H_1}, s_{H_2}, s_{H_3} \rangle$.

- (1) For all i , $s_{H_i}^2 = \text{id}_V$
- (2) For $i \neq j$, $(s_{H_i} \circ s_{H_j})^3 = 1$


Finite Reflection Groups



We obtain the group: $W = \langle s_{H_1}, s_{H_2}, s_{H_3} \rangle$.

- (1) For all i , $s_{H_i}^2 = \text{id}_V$
- (2) For $i \neq j$, $(s_{H_i} \circ s_{H_j})^3 = 1$
- (3) This group can be generated by two elements ($s_{H_3} = s_{H_1} \circ s_{H_2} \circ s_{H_1}$).

Finite Reflection Groups



We obtain the group: $W = \langle s_{H_1}, s_{H_2}, s_{H_3} \rangle$.

- (1) For all i , $s_{H_i}^2 = \text{id}_V$
- (2) For $i \neq j$, $(s_{H_i} \circ s_{H_j})^3 = 1$
- (3) This group can be generated by two elements ($s_{H_3} = s_{H_1} \circ s_{H_2} \circ s_{H_1}$).
- (4) This group is isomorphic to S_3 .

Finite Reflection Groups

Remarks:

Finite Reflection Groups

Remarks:

- (1) In general, each s_H has order 2 in W .

Finite Reflection Groups

Remarks:


- (1) In general, each s_H has order 2 in W .
- (2) Each hyperplane H is codimension 1, and thus H^\perp is spanned by some nonzero vector α .

Finite Reflection Groups

Remarks:

- (1) In general, each s_H has order 2 in W .
- (2) Each hyperplane H is codimension 1, and thus H^\perp is spanned by some nonzero vector α .

Example: For the hyperplane H_1 , we can choose $\alpha_1 = (0, 1.25, 0)$.



Finite Reflection Groups

Remarks:

Finite Reflection Groups

Remarks:

- (3) For each $H \in \mathcal{H}$, associate some $0 \neq \alpha \in H^\perp$, the collection of such vectors Φ is the (generalized) root system associated to the (Weyl) group $W =: W_\Phi$.

Finite Reflection Groups

Remarks:

- (3) For each $H \in \mathcal{H}$, associate some $0 \neq \alpha \in H^\perp$, the collection of such vectors Φ is the (generalized) root system associated to the (Weyl) group $W =: W_\Phi$.
- (4) For each $H \in \mathcal{H}$, we can explicitly write down s_H in terms of α (and thus we denote s_H by s_α):

$$s_\alpha(x) = x - 2 \frac{\langle \alpha, x \rangle}{\langle \alpha, \alpha \rangle} \alpha$$

Finite Reflection Groups

Remarks:

- (3) For each $H \in \mathcal{H}$, associate some $0 \neq \alpha \in H^\perp$, the collection of such vectors Φ is the (generalized) root system associated to the (Weyl) group $W =: W_\Phi$.
- (4) For each $H \in \mathcal{H}$, we can explicitly write down s_H in terms of α (and thus we denote s_H by s_α):

$$s_\alpha(x) = x - 2 \frac{\langle \alpha, x \rangle}{\langle \alpha, \alpha \rangle} \alpha$$

Example: Let $0 \neq \alpha \in H_1^\perp$, then

$$s_{\alpha_1}(\mathbf{u}) = (u_1, -u_2, u_3)$$

The function s_H is well-defined for any choice of $0 \neq \alpha \in H^\perp$.

Finite Reflection Groups

Remarks:

(5) Let $\alpha^\vee := \frac{2\alpha}{\langle \alpha, \alpha \rangle}$, then $s_\alpha(x) = x - \langle \alpha^\vee, x \rangle \alpha$.

Finite Reflection Groups

Remarks:

- (5) Let $\alpha^\vee := \frac{2\alpha}{\langle \alpha, \alpha \rangle}$, then $s_\alpha(x) = x - \langle \alpha^\vee, x \rangle \alpha$. This has a few properties:

Finite Reflection Groups

Remarks:

(5) Let $\alpha^\vee := \frac{2\alpha}{\langle \alpha, \alpha \rangle}$, then $s_\alpha(x) = x - \langle \alpha^\vee, x \rangle \alpha$. This has a few properties:

$$- \langle \alpha^\vee, \alpha \rangle = \left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \alpha \right\rangle = \frac{2}{\langle \alpha, \alpha \rangle} \langle \alpha, \alpha \rangle = 2$$

Finite Reflection Groups

Remarks:

(5) Let $\alpha^\vee := \frac{2\alpha}{\langle \alpha, \alpha \rangle}$, then $s_\alpha(x) = x - \langle \alpha^\vee, x \rangle \alpha$. This has a few properties:

$$\begin{aligned} - \langle \alpha^\vee, \alpha \rangle &= \left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \alpha \right\rangle = \frac{2}{\langle \alpha, \alpha \rangle} \langle \alpha, \alpha \rangle = 2 \\ - (\alpha^\vee)^\vee &= \frac{2\left(\frac{2\alpha}{\langle \alpha, \alpha \rangle}\right)}{\left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \frac{2\alpha}{\langle \alpha, \alpha \rangle} \right\rangle} = \frac{\frac{4}{\langle \alpha, \alpha \rangle} \alpha}{\frac{4}{\langle \alpha, \alpha \rangle^2} \langle \alpha, \alpha \rangle} = \alpha. \end{aligned}$$

Finite Reflection Groups

Remarks:

(5) Let $\alpha^\vee := \frac{2\alpha}{\langle \alpha, \alpha \rangle}$, then $s_\alpha(x) = x - \langle \alpha^\vee, x \rangle \alpha$. This has a few properties:

$$- \langle \alpha^\vee, \alpha \rangle = \left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \alpha \right\rangle = \frac{2}{\langle \alpha, \alpha \rangle} \langle \alpha, \alpha \rangle = 2$$

$$- (\alpha^\vee)^\vee = \frac{2\left(\frac{2\alpha}{\langle \alpha, \alpha \rangle}\right)}{\left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \frac{2\alpha}{\langle \alpha, \alpha \rangle} \right\rangle} = \frac{\frac{4}{\langle \alpha, \alpha \rangle} \alpha}{\frac{4}{\langle \alpha, \alpha \rangle^2} \langle \alpha, \alpha \rangle} = \alpha.$$

A crystallographic root system is a (generalized) root system if Φ satisfies:

$$\langle \alpha^\vee, \beta \rangle \in \mathbb{Z}$$

for all $\alpha, \beta \in \Phi$. In Lie algebra theory, this condition arises naturally.

Finite Reflection Groups

Remarks:

(5) Let $\alpha^\vee := \frac{2\alpha}{\langle \alpha, \alpha \rangle}$, then $s_\alpha(x) = x - \langle \alpha^\vee, x \rangle \alpha$. This has a few properties:

$$\begin{aligned} - \langle \alpha^\vee, \alpha \rangle &= \left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \alpha \right\rangle = \frac{2}{\langle \alpha, \alpha \rangle} \langle \alpha, \alpha \rangle = 2 \\ - (\alpha^\vee)^\vee &= \frac{2\left(\frac{2\alpha}{\langle \alpha, \alpha \rangle}\right)}{\left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \frac{2\alpha}{\langle \alpha, \alpha \rangle} \right\rangle} = \frac{\frac{4}{\langle \alpha, \alpha \rangle} \alpha}{\frac{4}{\langle \alpha, \alpha \rangle^2} \langle \alpha, \alpha \rangle} = \alpha. \end{aligned}$$

A crystallographic root system is a (generalized) root system if Φ satisfies:

$$\langle \alpha^\vee, \beta \rangle \in \mathbb{Z}$$

for all $\alpha, \beta \in \Phi$. In Lie algebra theory, this condition arises naturally. We define $\Phi^\vee := \{\alpha^\vee : \alpha \in \Phi\}$ to be the coroot system of W , in particular, a Weyl group can have multiple root systems (the choice of Φ is important to the structure).

Finite Reflection Groups

We have an orthogonal decomposition of V via the action (W, Φ) .

Finite Reflection Groups

We have an orthogonal decomposition of V via the action (W, Φ) .

$$V_0 := \bigcap_{\alpha \in \Phi} \alpha^\perp$$

Finite Reflection Groups

We have an orthogonal decomposition of V via the action (W, Φ) .

$$V_0 := \bigcap_{\alpha \in \Phi} \alpha^\perp$$

This is the fixed-point set $V^W = \{v \in V : wv = v \text{ for all } w \in W\}$. We call this the *inessential part* of V .

Finite Reflection Groups

We have an orthogonal decomposition of V via the action (W, Φ) .

$$V_0 := \bigcap_{\alpha \in \Phi} \alpha^\perp$$

This is the fixed-point set $V^W = \{v \in V : wv = v \text{ for all } w \in W\}$. We call this the *inessential part* of V .

$$V_1 := \text{span}_{\mathbb{R}}(\Phi)$$

Finite Reflection Groups

We have an orthogonal decomposition of V via the action (W, Φ) .

$$V_0 := \bigcap_{\alpha \in \Phi} \alpha^\perp$$

This is the fixed-point set $V^W = \{v \in V : wv = v \text{ for all } w \in W\}$. We call this the *inessential part* of V .

$$V_1 := \text{span}_{\mathbb{R}}(\Phi)$$

The action of W is completely determined by V_1 . We call this the *essential part* of V .

Finite Reflection Groups

We have an orthogonal decomposition of V via the action (W, Φ) .

$$V_0 := \bigcap_{\alpha \in \Phi} \alpha^\perp$$

This is the fixed-point set $V^W = \{v \in V : wv = v \text{ for all } w \in W\}$. We call this the *inessential part* of V .




$$V_1 := \text{span}_{\mathbb{R}}(\Phi)$$

The action of W is completely determined by V_1 . We call this the *essential part* of V .

$$V = V_0 \oplus V_1$$


Finite Reflection Groups

Example: Let $V = \mathbb{R}^3$, and H_1 , H_2 and H_3 as before and choose orthogonal vectors $\alpha_1 = (0, 1, 0)$, $\alpha_2 = (0, -1, \sqrt{3})$, and $\alpha_3 = (0, 1, \sqrt{3})$, respectively.



Finite Reflection Groups


Example: Let $V = \mathbb{R}^3$, and H_1 , H_2 and H_3 as before and choose orthogonal vectors $\alpha_1 = (0, 1, 0)$, $\alpha_2 = (0, -1, \sqrt{3})$, and $\alpha_3 = (0, 1, \sqrt{3})$, respectively.




The x -axis is contained in each hyperplane and thus is orthogonal to each α_i chosen, or equivalently, is fixed by the reflections across each H_i :

$$V_0 := (0, 1, 0)^\perp \cap (0, -1, \sqrt{3})^\perp \cap (0, 1, \sqrt{3})^\perp$$


Finite Reflection Groups



Finite Reflection Groups



By contracting V_0 to a point, we see the essential part of (W, Φ) :



Finite Reflection Groups

Remarks:

- (6) One can restrict themselves to studying essential reflection groups, i.e. for $V = V_0 \oplus V_1$ as above, we require $V_0 = 0$.

Finite Reflection Groups

Remarks:

- (6) One can restrict themselves to studying essential reflection groups, i.e. for $V = V_0 \oplus V_1$ as above, we require $V_0 = 0$.
- (7) We can define a product $(W', V') \times (W'', V'') := (W' \times W'', V' \oplus V'')$. If (W, V) cannot be expressed as a product, then we say that (W, V) is irreducible. One can restrict themselves to studying irreducible reflection groups.

Finite Reflection Groups

Remarks:

- (6) One can restrict themselves to studying essential reflection groups, i.e. for $V = V_0 \oplus V_1$ as above, we require $V_0 = 0$.
- (7) We can define a product $(W', V') \times (W'', V'') := (W' \times W'', V' \oplus V'')$. If (W, V) cannot be expressed as a product, then we say that (W, V) is irreducible. One can restrict themselves to studying irreducible reflection groups.
- (8) There is a classification of non-trivial, essential, irreducible, finite reflection groups.

Finite Reflection Groups

Remarks:

- (6) One can restrict themselves to studying essential reflection groups, i.e. for $V = V_0 \oplus V_1$ as above, we require $V_0 = 0$.
- (7) We can define a product $(W', V') \times (W'', V'') := (W' \times W'', V' \oplus V'')$. If (W, V) cannot be expressed as a product, then we say that (W, V) is irreducible. One can restrict themselves to studying irreducible reflection groups.
- (8) There is a classification of non-trivial, essential, irreducible, finite reflection groups.

Among these are: group of permutations on n letters, group of signed permutations on n letters, dihedral groups, symmetries of regular solids (see Section 1.3 in Buildings - Theory and Applications, Abramenko and Brown, 2008).

Finite Reflection Groups

We have introduced quite a bit of notation:

- V is a Euclidean vector space.
- n is the dimension of V .
- H is a hyperplane, there exists $0 \neq \alpha \in H^\perp$ called a root.
- \mathcal{H} is a hyperplane arrangement, Φ is a generalized root system.
- s_H is the reflection across H , also denoted s_α , where α is a root (determined by H).
- W is the Weyl group.
- $V_1 = \text{span}_{\mathbb{R}}(\Phi)$ is the essential part of V .
- $V_0 = V^W$ is the unessential part of V .

Coxeter Complex

Definition: A (finite) abstract simplicial complex is a (finite) set A together with a collection Δ of finite subsets of A such that if $X \in \Delta$ and $Y \subseteq X$, then $Y \in \Delta$.

Coxeter Complex

Definition: A (finite) abstract simplicial complex is a (finite) set A together with a collection Δ of finite subsets of A such that if $X \in \Delta$ and $Y \subseteq X$, then $Y \in \Delta$. An abstract simplicial complex is denoted by the collection Δ .

Coxeter Complex

Definition: A (finite) abstract simplicial complex is a (finite) set A together with a collection Δ of finite subsets of A such that if $X \in \Delta$ and $Y \subseteq X$, then $Y \in \Delta$. An abstract simplicial complex is denoted by the collection Δ .

(1) A vertex is a $v \in A$ such that $\{v\} \in \Delta$.

Coxeter Complex

Definition: A (finite) abstract simplicial complex is a (finite) set A together with a collection Δ of finite subsets of A such that if $X \in \Delta$ and $Y \subseteq X$, then $Y \in \Delta$. An abstract simplicial complex is denoted by the collection Δ .

- (1) A vertex is a $v \in A$ such that $\{v\} \in \Delta$.
- (2) The set of vertices is denoted $V(\Delta)$.

Coxeter Complex

Definition: A (finite) abstract simplicial complex is a (finite) set A together with a collection Δ of finite subsets of A such that if $X \in \Delta$ and $Y \subseteq X$, then $Y \in \Delta$. An abstract simplicial complex is denoted by the collection Δ .

- (1) A vertex is a $v \in A$ such that $\{v\} \in \Delta$.
- (2) The set of vertices is denoted $V(\Delta)$.
- (3) The elements of Δ are called simplices.

Coxeter Complex

Definition: A (finite) abstract simplicial complex is a (finite) set A together with a collection Δ of finite subsets of A such that if $X \in \Delta$ and $Y \subseteq X$, then $Y \in \Delta$. An abstract simplicial complex is denoted by the collection Δ .

- (1) A vertex is a $v \in A$ such that $\{v\} \in \Delta$.
- (2) The set of vertices is denoted $V(\Delta)$.
- (3) The elements of Δ are called simplices.
- (4) The dimension of a simplex $\delta \in \Delta$ is $|\delta| - 1$, i.e. vertices are zero dimensional, edges are one-dimensional, etc.

Coxeter Complex

Definition: A (finite) abstract simplicial complex is a (finite) set A together with a collection Δ of finite subsets of A such that if $X \in \Delta$ and $Y \subseteq X$, then $Y \in \Delta$. An abstract simplicial complex is denoted by the collection Δ .

- (1) A vertex is a $v \in A$ such that $\{v\} \in \Delta$.
- (2) The set of vertices is denoted $V(\Delta)$.
- (3) The elements of Δ are called simplices.
- (4) The dimension of a simplex $\delta \in \Delta$ is $|\delta| - 1$, i.e. vertices are zero dimensional, edges are one-dimensional, etc.



To simplify notation with subsets, I will denote subsets $X = \{a_1, a_2, \dots, a_k\}$ by $X = a_1 a_2 \dots a_k$.

Coxeter Complex

Example: Let $A = \{1, 2, 3, 4\}$ and consider
 $\Delta_1 = \{\emptyset, 1, 2, 3, 12, 13, 23, 123\}$ and $\Delta_2 = \{\emptyset, 1, 2, 4, 12, 14\}$.

Coxeter Complex



Example: Let $A = \{1, 2, 3, 4\}$ and consider $\Delta_1 = \{\emptyset, 1, 2, 3, 12, 13, 23, 123\}$ and $\Delta_2 = \{\emptyset, 1, 2, 4, 12, 14\}$.




Coxeter Complex

Example: Let $A = \{1, 2, 3, 4\}$ and consider

$\Delta_1 = \{\emptyset, 1, 2, 3, 12, 13, 23, 123\}$ and $\Delta_2 = \{\emptyset, 1, 2, 4, 12, 14\}$.



However, $\Delta_3 = \{\emptyset, 1, 2, 23, 123\}$ is not an abstract simplicial complex.



Coxeter Complex

Definition: Let Δ_1 and Δ_2 be two abstract simplicial complexes. A simplicial map is a function $f : V(\Delta_1) \rightarrow V(\Delta_2)$ such that for all $\delta \in \Delta_1$, $f(\delta) \in \Delta_2$.

Coxeter Complex


Definition: Let Δ_1 and Δ_2 be two abstract simplicial complexes. A simplicial map is a function $f : V(\Delta_1) \rightarrow V(\Delta_2)$ such that for all $\delta \in \Delta_1$, $f(\delta) \in \Delta_2$.

Example: Let $f : V(\Delta_1) \rightarrow V(\Delta_2)$ be defined by $f(1) = 1$ and $f(2) = 2$ and $f(3) = 1$.

Coxeter Complex

Definition: Let Δ_1 and Δ_2 be two abstract simplicial complexes. A simplicial map is a function $f : V(\Delta_1) \rightarrow V(\Delta_2)$ such that for all $\delta \in \Delta_1$, $f(\delta) \in \Delta_2$.


Example: Let $f : V(\Delta_1) \rightarrow V(\Delta_2)$ be defined by $f(1) = 1$ and $f(2) = 2$ and $f(3) = 1$.



Coxeter Complex

Definition: Let Δ_1 and Δ_2 be two abstract simplicial complexes. A simplicial map is a function $f : V(\Delta_1) \rightarrow V(\Delta_2)$ such that for all $\delta \in \Delta_1$, $f(\delta) \in \Delta_2$.


Example: Let $f : V(\Delta_1) \rightarrow V(\Delta_2)$ be defined by $f(1) = 1$ and $f(2) = 2$ and $f(3) = 1$.



Coxeter Complex

Definition: Let Δ_1 and Δ_2 be two abstract simplicial complexes. A simplicial map is a function $f : V(\Delta_1) \rightarrow V(\Delta_2)$ such that for all $\delta \in \Delta_1$, $f(\delta) \in \Delta_2$.


Example: Let $f : V(\Delta_1) \rightarrow V(\Delta_2)$ be defined by $f(1) = 1$ and $f(2) = 2$ and $f(3) = 1$.



Coxeter Complex

Definition: Let Δ_1 and Δ_2 be two abstract simplicial complexes. A simplicial map is a function $f : V(\Delta_1) \rightarrow V(\Delta_2)$ such that for all $\delta \in \Delta_1$, $f(\delta) \in \Delta_2$.


Example: Let $f : V(\Delta_1) \rightarrow V(\Delta_2)$ be defined by $f(1) = 1$ and $f(2) = 2$ and $f(3) = 1$.



Coxeter Complex

Definition: Let Δ_1 and Δ_2 be two abstract simplicial complexes. A simplicial map is a function $f : V(\Delta_1) \rightarrow V(\Delta_2)$ such that for all $\delta \in \Delta_1$, $f(\delta) \in \Delta_2$.

Example: Let $f : V(\Delta_1) \rightarrow V(\Delta_2)$ be defined by $f(1) = 1$ and $f(2) = 2$ and $f(3) = 1$.



Coxeter Complex

(1) The identity map $\text{id}_\Delta : \Delta \rightarrow \Delta$ is a simplicial map.

Coxeter Complex

- (1) The identity map $\text{id}_\Delta : \Delta \rightarrow \Delta$ is a simplicial map.
- (2) The composition of simplicial maps is simplicial.


Coxeter Complex

- (1) The identity map $\text{id}_\Delta : \Delta \rightarrow \Delta$ is a simplicial map.
- (2) The composition of simplicial maps is simplicial.
- (3) An isomorphism of simplicial complexes Δ_1 and Δ_2 is a bijective function $f : V(\Delta_1) \rightarrow V(\Delta_2)$ such that f^{-1} is simplicial.

Coxeter Complex

- (1) The identity map $\text{id}_\Delta : \Delta \rightarrow \Delta$ is a simplicial map.
- (2) The composition of simplicial maps is simplicial.
- (3) An isomorphism of simplicial complexes Δ_1 and Δ_2 is a bijective function $f : V(\Delta_1) \rightarrow V(\Delta_2)$ such that f^{-1} is simplicial.


Example: Let $f : V(\Delta_2) \rightarrow V(\Delta_1)$ be defined by $f(1) = 1$, $f(2) = 2$, and $f(4) = 3$.



Coxeter Complex

- (1) The identity map $\text{id}_\Delta : \Delta \rightarrow \Delta$ is a simplicial map.
- (2) The composition of simplicial maps is simplicial.
- (3) An isomorphism of simplicial complexes Δ_1 and Δ_2 is a bijective function $f : V(\Delta_1) \rightarrow V(\Delta_2)$ such that f^{-1} is simplicial.


Example: Let $f : V(\Delta_2) \rightarrow V(\Delta_1)$ be defined by $f(1) = 1$, $f(2) = 2$, and $f(4) = 3$.



Coxeter Complex

- (1) The identity map $\text{id}_\Delta : \Delta \rightarrow \Delta$ is a simplicial map.
- (2) The composition of simplicial maps is simplicial.
- (3) An isomorphism of simplicial complexes Δ_1 and Δ_2 is a bijective function $f : V(\Delta_1) \rightarrow V(\Delta_2)$ such that f^{-1} is simplicial.


Example: Let $f : V(\Delta_2) \rightarrow V(\Delta_1)$ be defined by $f(1) = 1$, $f(2) = 2$, and $f(4) = 3$.



Coxeter Complex

- (1) The identity map $\text{id}_\Delta : \Delta \rightarrow \Delta$ is a simplicial map.
- (2) The composition of simplicial maps is simplicial.
- (3) An isomorphism of simplicial complexes Δ_1 and Δ_2 is a bijective function $f : V(\Delta_1) \rightarrow V(\Delta_2)$ such that f^{-1} is simplicial.


Example: Let $f : V(\Delta_2) \rightarrow V(\Delta_1)$ be defined by $f(1) = 1$, $f(2) = 2$, and $f(4) = 3$.



Coxeter Complex

- (1) The identity map $\text{id}_\Delta : \Delta \rightarrow \Delta$ is a simplicial map.
- (2) The composition of simplicial maps is simplicial.
- (3) An isomorphism of simplicial complexes Δ_1 and Δ_2 is a bijective function $f : V(\Delta_1) \rightarrow V(\Delta_2)$ such that f^{-1} is simplicial.

Example: Let $f : V(\Delta_2) \rightarrow V(\Delta_1)$ be defined by $f(1) = 1$, $f(2) = 2$, and $f(4) = 3$.




Coxeter Complex

Our next goal is to construct to a chamber complex associated to an essential finite reflection group (W, V) , which we will denote by $\Sigma := \Sigma(W, S)$.

Coxeter Complex

Our next goal is to construct a chamber complex associated to an essential finite reflection group (W, V) , which we will denote by $\Sigma := \Sigma(W, S)$. Let's return to our running example:



where $V = \mathbb{R}^3$ with the usual inner product. We can reformulate this example in terms of the essential part of the action.

Coxeter Complex

Let $V = \mathbb{R}^2$, with the usual inner product. Let $\mathcal{H} = \{H_1, H_2, H_3\}$.

- $H_1 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (0, u_2)\}$
- $H_2 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (\sqrt{3}u_2, u_2)\}$
- $H_3 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (\sqrt{3}u_2, -u_2)\}$

Coxeter Complex

Let $V = \mathbb{R}^2$, with the usual inner product. Let $\mathcal{H} = \{H_1, H_2, H_3\}$.

- $H_1 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (0, u_2)\}$
- $H_2 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (\sqrt{3}u_2, u_2)\}$
- $H_3 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (\sqrt{3}u_2, -u_2)\}$

Set $s := s_{H_1}$ and $t := s_{H_2}$, then $s_{H_3} = s \circ t \circ s$. We will denote the *set of fundamental reflections* by $S = \{s, t\}$.

Coxeter Complex

Let $V = \mathbb{R}^2$, with the usual inner product. Let $\mathcal{H} = \{H_1, H_2, H_3\}$.

- $H_1 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (0, u_2)\}$
- $H_2 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (\sqrt{3}u_2, u_2)\}$
- $H_3 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (\sqrt{3}u_2, -u_2)\}$

Set $s := s_{H_1}$ and $t := s_{H_2}$, then $s_{H_3} = s \circ t \circ s$. We will denote the *set of fundamental reflections* by $S = \{s, t\}$. To each hyperplane, we will associate a linear function $f_i : V \rightarrow \mathbb{R}$ such that $f_i|_{H_i} = 0$ and $f_i \neq 0$.

- $f_1(x, y) = x$
- $f_2(x, y) = -x + \sqrt{3}y$
- $f_3(x, y) = x + \sqrt{3}y$

I will refer to the pair (W, S) as a Coxeter group.

Coxeter Complex

Definition: A *cell* in V with respect to $\mathcal{H} = \{H_i\}_{i \in I}$ is a nonempty set A obtained by choosing for each i a sign $\sigma_i \in \{+, -, 0\}$ and specifying $f_i = \sigma_i$.

Coxeter Complex

Definition: A *cell* in V with respect to $\mathcal{H} = \{H_i\}_{i \in I}$ is a nonempty set A obtained by choosing for each i a sign $\sigma_i \in \{+, -, 0\}$ and specifying $f_i = \sigma_i$. In other words, A is defined by a set of linear equalities and strict inequalities, one for each hyperplane in \mathcal{H} .

$$A = \bigcap_{i \in I} U_i$$

for U_i is either H_i or an open half-space with H_i as a boundary. We denote the set of cells $\Sigma(\mathcal{H})$.

Coxeter Complex

Definition: A *cell* in V with respect to $\mathcal{H} = \{H_i\}_{i \in I}$ is a nonempty set A obtained by choosing for each i a sign $\sigma_i \in \{+, -, 0\}$ and specifying $f_i = \sigma_i$. In other words, A is defined by a set of linear equalities and strict inequalities, one for each hyperplane in \mathcal{H} .

$$A = \bigcap_{i \in I} U_i$$

for U_i is either H_i or an open half-space with H_i as a boundary. We denote the set of cells $\Sigma(\mathcal{H})$.

Definition: If each $\sigma_i \neq 0$, then we call the cell a *chamber*. We denote the set of chamber by $\mathcal{C}(\mathcal{H})$.

Coxeter Complex

Definition: A *cell* in V with respect to $\mathcal{H} = \{H_i\}_{i \in I}$ is a nonempty set A obtained by choosing for each i a sign $\sigma_i \in \{+, -, 0\}$ and specifying $f_i = \sigma_i$. In other words, A is defined by a set of linear equalities and strict inequalities, one for each hyperplane in \mathcal{H} .








$$A = \bigcap_{i \in I} U_i$$

for U_i is either H_i or an open half-space with H_i as a boundary. We denote the set of cells $\Sigma(\mathcal{H})$.


Definition: If each $\sigma_i \neq 0$, then we call the cell a *chamber*. We denote the set of chamber by $\mathcal{C}(\mathcal{H})$.

Definition: If $\sigma_i = 0$ for exactly one $i \in I$, then we call the cell a *panel*.


Coxeter Complex




Coxeter Complex




$$f_1 = 0$$




$$f_1 = +$$




$$f_1 = -$$




$$f_2 = 0$$



$$f_2 = +$$



$$f_2 = -$$



$$f_3 = 0$$




$$f_3 = +$$











$$f_3 = -$$


Coxeter Complex



Coxeter Complex



Coxeter Complex




Coxeter Complex

Definition: Given cells $A, B \in \Sigma(\mathcal{H})$, B is a face of A , denoted $B \leq A$, if for each $i \in I$, either $\sigma_i(B) = 0$ or $\sigma_i(B) = \sigma_i(A)$.

Coxeter Complex

Definition: Given cells $A, B \in \Sigma(\mathcal{H})$, B is a face of A , denoted $B \leq A$, if for each $i \in I$, either $\sigma_i(B) = 0$ or $\sigma_i(B) = \sigma_i(A)$.


Example: In our previous examples of cells, $B \leq A$.



We say that B is a panel of A , and the hyperplane H_1 containing B is the *wall* of A .

Coxeter Complex

In this example, there are 13 cells: 6 chambers, 6 open rays and the origin.



Coxeter Complex


We distinguish an arbitrary chamber C in the cell decomposition.

Coxeter Complex

We distinguish an arbitrary chamber C in the cell decomposition. Recall $s = s_{H_1}$ and $t = s_{H_2}$.


Coxeter Complex

We distinguish an arbitrary chamber C in the cell decomposition. Recall $s = s_{H_1}$ and $t = s_{H_2}$.



Coxeter Complex

We distinguish an arbitrary chamber C in the cell decomposition. Recall $s = s_{H_1}$ and $t = s_{H_2}$.



Note $w_0 = sts = tst$. By this definition, it is clear W acts transitively on the chambers and $|\mathcal{C}(\mathcal{H})| = |W|$.

Coxeter Complex

Let $\Sigma_{\leq C}$ be the subcomplex of faces of C . For $A \leq C$, let W_A be the stabilizer of A .

Coxeter Complex

Let $\Sigma_{\leq C}$ be the subcomplex of faces of C . For $A \leq C$, let W_A be the stabilizer of A .


Example: Let C be the fundamental chamber, C , R_1^+ , R_2^+ , and 0 are the faces of C .

$$W_C = \{e\}$$

$$W_{R_1^+} = \{e, s\}$$

$$W_{R_2^+} = \{e, t\}$$

$$W_0 = W$$



Coxeter Complex

Note that $W_{R_1^+} = \langle s \rangle$, $W_{R_2^+} = \langle t \rangle$ and $W_0 = \langle s, t \rangle$.

Coxeter Complex

Note that $W_{R_1^+} = \langle s \rangle$, $W_{R_2^+} = \langle t \rangle$ and $W_0 = \langle s, t \rangle$.


The stabilizers are generated by subsets of the set of fundamental reflections S , which are called *parabolic subgroups* of (W, S) .

Coxeter Complex

Note that $W_{R_1^+} = \langle s \rangle$, $W_{R_2^+} = \langle t \rangle$ and $W_0 = \langle s, t \rangle$.

The stabilizers are generated by subsets of the set of fundamental reflections S , which are called *parabolic subgroups* of (W, S) . There is an order-reversing isomorphism of posets:

$$\phi : \Sigma_{\leq C} \xrightarrow{\sim} (\text{parabolic subgroups})^{\text{op}}$$




Coxeter Complex

Note that $W_{R_1^+} = \langle s \rangle$, $W_{R_2^+} = \langle t \rangle$ and $W_0 = \langle s, t \rangle$.


The stabilizers are generated by subsets of the set of fundamental reflections S , which are called *parabolic subgroups* of (W, S) . There is an order-reversing isomorphism of posets:

$$\phi : \Sigma_{\leq C} \xrightarrow{\sim} (\text{parabolic subgroups})^{\text{op}}$$



We can extend this isomorphism (of posets) to the whole poset Σ by including cosets of parabolic subgroups $\hat{\phi} : \Sigma \xrightarrow{\sim} (\text{parabolic cosets})^{\text{op}}$.

Coxeter Complex



Coxeter Complex

Therefore, using the isomorphism of posets $\hat{\phi}$, we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on $\Sigma(W, S)$. Via this isomorphism, W is the underlying set of Σ .

Coxeter Complex

Therefore, using the isomorphism of posets $\hat{\phi}$, we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on $\Sigma(W, S)$. Via this isomorphism, W is the underlying set of Σ .

- The vertices of Σ are $\{w\}$, which are the chambers.

Coxeter Complex

Therefore, using the isomorphism of posets $\hat{\phi}$, we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on $\Sigma(W, S)$. Via this isomorphism, W is the underlying set of Σ .

- The vertices of Σ are $\{w\}$, which are the chambers.
- The edges of Σ are $w\langle u \rangle = \{w, wu\}$ which is a panel of $\{w\}$ (the parabolic cosets of order 2). We say w and wu are u -adjacent.

Coxeter Complex

Therefore, using the isomorphism of posets $\hat{\phi}$, we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on $\Sigma(W, S)$. Via this isomorphism, W is the underlying set of Σ .

- The vertices of Σ are $\{w\}$, which are the chambers.
- The edges of Σ are $w\langle u \rangle = \{w, wu\}$ which is a panel of $\{w\}$ (the parabolic cosets of order 2). We say w and wu are u -adjacent.

Remarks:

Coxeter Complex

Therefore, using the isomorphism of posets $\hat{\phi}$, we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on $\Sigma(W, S)$. Via this isomorphism, W is the underlying set of Σ .

- The vertices of Σ are $\{w\}$, which are the chambers.
- The edges of Σ are $w\langle u \rangle = \{w, wu\}$ which is a panel of $\{w\}$ (the parabolic cosets of order 2). We say w and wu are u -adjacent.

Remarks:

- (1) Σ is *thin*, meaning each panel is the face of exactly two chambers.

Coxeter Complex

Therefore, using the isomorphism of posets $\hat{\phi}$, we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on $\Sigma(W, S)$. Via this isomorphism, W is the underlying set of Σ .

- The vertices of Σ are $\{w\}$, which are the chambers.
- The edges of Σ are $w\langle u \rangle = \{w, wu\}$ which is a panel of $\{w\}$ (the parabolic cosets of order 2). We say w and wu are u -adjacent.

Remarks:

- (1) Σ is *thin*, meaning each panel is the face of exactly two chambers.
- (2) Σ is *colorable*, meaning we can assign a labeling which distinguishes types of vertices.

Coxeter Complex

Therefore, using the isomorphism of posets $\hat{\phi}$, we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on $\Sigma(W, S)$. Via this isomorphism, W is the underlying set of Σ .

- The vertices of Σ are $\{w\}$, which are the chambers.
- The edges of Σ are $w\langle u \rangle = \{w, wu\}$ which is a panel of $\{w\}$ (the parabolic cosets of order 2). We say w and wu are u -adjacent.

Remarks:

- (1) Σ is *thin*, meaning each panel is the face of exactly two chambers.
- (2) Σ is *colorable*, meaning we can assign a labeling which distinguishes types of vertices.
- (3) The action of W on Σ is *type-preserving*, meaning that the coloring is preserved under the action.

Coxeter Complex

Definition: Any simplicial complex Δ isomorphic to $\Sigma(W, S)$ for some (W, S) Coxeter group (i.e. a finite reflection group) is a Coxeter complex.

Coxeter Complex

Definition: Any simplicial complex Δ isomorphic to $\Sigma(W, S)$ for some (W, S) Coxeter group (i.e. a finite reflection group) is a Coxeter complex.

Definition of a Building

A (*weak*) *building* is a simplicial complex Δ that can be expressed as the union of subcomplexes Σ (called *apartments*) satisfying axioms:

- (B0) Each apartment Σ is a Coxeter complex.
- (B1) For any two simplices $A, B \in \Delta$, there is an apartment Σ containing both of them.
- (B2) If Σ and Σ' are two apartments containing A and B , then there is an isomorphism $\Sigma \rightarrow \Sigma'$ fixing A and B pointwise.

Coxeter Complex

Definition: Any simplicial complex Δ isomorphic to $\Sigma(W, S)$ for some (W, S) Coxeter group (i.e. a finite reflection group) is a Coxeter complex.

Definition of a Building

A (weak) *building* is a simplicial complex Δ that can be expressed as the union of subcomplexes Σ (called *apartments*) satisfying axioms:

- (B0) Each apartment Σ is a Coxeter complex.
- (B1) For any two simplices $A, B \in \Delta$, there is an apartment Σ containing both of them.
- (B2) If Σ and Σ' are two apartments containing A and B , then there is an isomorphism $\Sigma \rightarrow \Sigma'$ fixing A and B pointwise.

Remark: Coxeter complexes characterize thin buildings with a single apartment.

Buildings

For a building Δ , let \mathcal{A} be the collection of apartments satisfying the axioms of a building (this is called a *system of apartments* for Δ).

Buildings

For a building Δ , let \mathcal{A} be the collection of apartments satisfying the axioms of a building (this is called a *system of apartments* for Δ).

Remarks:

- (1) We do not require a specific system of apartments; however, every building admits a canonical system of apartments.

Buildings

For a building Δ , let \mathcal{A} be the collection of apartments satisfying the axioms of a building (this is called a *system of apartments* for Δ).

Remarks:

- (1) We do not require a specific system of apartments; however, every building admits a canonical system of apartments.
- (2) Furthermore, when the apartments are finite Coxeter complexes, there is a unique system of apartments.

Buildings

For a building Δ , let \mathcal{A} be the collection of apartments satisfying the axioms of a building (this is called a *system of apartments* for Δ).

Remarks:

- (1) We do not require a specific system of apartments; however, every building admits a canonical system of apartments.
- (2) Furthermore, when the apartments are finite Coxeter complexes, there is a unique system of apartments.
- (3) A building is colorable, and isomorphisms between buildings can be taken to be type-preserving.

Buildings

For a building Δ , let \mathcal{A} be the collection of apartments satisfying the axioms of a building (this is called a *system of apartments* for Δ).

Remarks:

- (1) We do not require a specific system of apartments; however, every building admits a canonical system of apartments.
- (2) Furthermore, when the apartments are finite Coxeter complexes, there is a unique system of apartments.
- (3) A building is colorable, and isomorphisms between buildings can be taken to be type-preserving.
- (4) All apartments are Coxeter complexes for the same Coxeter group.

Buildings

Definition: Let P be a set with a binary relation \mathcal{I} called “incidence” such that \mathcal{I} is reflexive and symmetric. A *flag* in P is a set of pairwise incident elements of P .

Buildings

Definition: Let P be a set with a binary relation \mathcal{I} called “incidence” such that \mathcal{I} is reflexive and symmetric. A *flag* in P is a set of pairwise incident elements of P .

Note: If P is a poset, then a flag is a linearly ordered subset of P .

We will assume that P is partitioned into nonempty subsets P_0, P_1, \dots, P_{n-1} , where $p \in P_i$ is said to have type i .

Buildings

Example: Let V be a finite dimension F -vector space (dimension $n \geq 2$).

Buildings

Example: Let V be a finite dimension F -vector space (dimension $n \geq 2$). Let P be the collection of nonzero proper subspaces of V and let two subspaces be “incident” if one is contained in the other.

Buildings

Example: Let V be a finite dimension F -vector space (dimension $n \geq 2$). Let P be the collection of nonzero proper subspaces of V and let two subspaces be “incident” if one is contained in the other. Let $\Delta(V)$ be the flag complex consisting of simplices as chains of subspaces

$$V_1 < V_2 < \cdots < V_k$$

where $\dim_F(V_i) = i$. *Chambers* are chains:

$$V_1 < V_2 < \cdots < V_{n-1}$$

Buildings

Example: Let V be a finite dimension F -vector space (dimension $n \geq 2$). Let P be the collection of nonzero proper subspaces of V and let two subspaces be “incident” if one is contained in the other. Let $\Delta(V)$ be the flag complex consisting of simplices as chains of subspaces

$$V_1 < V_2 < \cdots < V_k$$

where $\dim_F(V_i) = i$. *Chambers* are chains:

$$V_1 < V_2 < \cdots < V_{n-1}$$

Let $\mathcal{F} = \{L_1, \dots, L_n\}$ be a collection of 1-dimensional subspaces which span V . The subcomplex $\Sigma(\mathcal{F})$ of flags is an *apartment*.

Buildings

Example: Let V be a finite dimension F -vector space (dimension $n \geq 2$). Let P be the collection of nonzero proper subspaces of V and let two subspaces be “incident” if one is contained in the other. Let $\Delta(V)$ be the flag complex consisting of simplices as chains of subspaces

$$V_1 < V_2 < \cdots < V_k$$

where $\dim_F(V_i) = i$. *Chambers* are chains:

$$V_1 < V_2 < \cdots < V_{n-1}$$

Let $\mathcal{F} = \{L_1, \dots, L_n\}$ be a collection of 1-dimensional subspaces which span V . The subcomplex $\Sigma(\mathcal{F})$ of flags is an *apartment*.

Proof: S_n acts on each $\Sigma(\mathcal{F})$, each chain is contained in a composition series, and apply Jordan-Hölder theorem to obtain canonical isomorphisms and projections.

Buildings

Consider $V = \mathbb{F}_2^3$.

Buildings

Consider $V = \mathbb{F}_2^3$. Set

$$\begin{array}{llll} L_0 := \mathbb{F}_2(0, 0, 0) & L_1 := \mathbb{F}_2(1, 0, 0) & L_2 := \mathbb{F}_2(0, 1, 0) & L_3 := \mathbb{F}_2(0, 0, 1) \\ L_4 := \mathbb{F}_2(1, 1, 0) & L_5 := \mathbb{F}_2(1, 0, 1) & L_6 := \mathbb{F}_2(0, 1, 1) & L_7 := \mathbb{F}_2(1, 1, 1) \end{array}$$

Buildings

Consider $V = \mathbb{F}_2^3$. Set

$$\begin{aligned} L_0 &:= \mathbb{F}_2(0, 0, 0) & L_1 &:= \mathbb{F}_2(1, 0, 0) & L_2 &:= \mathbb{F}_2(0, 1, 0) & L_3 &:= \mathbb{F}_2(0, 0, 1) \\ L_4 &:= \mathbb{F}_2(1, 1, 0) & L_5 &:= \mathbb{F}_2(1, 0, 1) & L_6 &:= \mathbb{F}_2(0, 1, 1) & L_7 &:= \mathbb{F}_2(1, 1, 1) \end{aligned}$$

Note that L_0 is not a 1-dimensional subspace. Therefore, we have seven 1-dimensional subspaces, which we will call “type 1” vertices.

Buildings

Consider $V = \mathbb{F}_2^3$. Set

$$\begin{aligned} L_0 &:= \mathbb{F}_2(0, 0, 0) & L_1 &:= \mathbb{F}_2(1, 0, 0) & L_2 &:= \mathbb{F}_2(0, 1, 0) & L_3 &:= \mathbb{F}_2(0, 0, 1) \\ L_4 &:= \mathbb{F}_2(1, 1, 0) & L_5 &:= \mathbb{F}_2(1, 0, 1) & L_6 &:= \mathbb{F}_2(0, 1, 1) & L_7 &:= \mathbb{F}_2(1, 1, 1) \end{aligned}$$

Note that L_0 is not a 1-dimensional subspace. Therefore, we have seven 1-dimensional subspaces, which we will call “type 1” vertices. Set

$$\begin{aligned} V_1 &:= L_1 + L_2 & V_2 &:= L_1 + L_3 & V_3 &:= L_1 + L_4 & V_4 &:= L_2 + L_3 \\ V_5 &:= L_2 + L_5 & V_6 &:= L_3 + L_4 & V_7 &:= L_4 + L_5 \end{aligned}$$

Therefore, we have seven 2-dimensional subspaces, which we will call “type 2” vertices.

Buildings

Consider $V = \mathbb{F}_2^3$. Set

$$\begin{aligned} L_0 &:= \mathbb{F}_2(0, 0, 0) & L_1 &:= \mathbb{F}_2(1, 0, 0) & L_2 &:= \mathbb{F}_2(0, 1, 0) & L_3 &:= \mathbb{F}_2(0, 0, 1) \\ L_4 &:= \mathbb{F}_2(1, 1, 0) & L_5 &:= \mathbb{F}_2(1, 0, 1) & L_6 &:= \mathbb{F}_2(0, 1, 1) & L_7 &:= \mathbb{F}_2(1, 1, 1) \end{aligned}$$

Note that L_0 is not a 1-dimensional subspace. Therefore, we have seven 1-dimensional subspaces, which we will call “type 1” vertices. Set


$$\begin{aligned} V_1 &:= L_1 + L_2 & V_2 &:= L_1 + L_3 & V_3 &:= L_1 + L_4 & V_4 &:= L_2 + L_3 \\ V_5 &:= L_2 + L_5 & V_6 &:= L_3 + L_4 & V_7 &:= L_4 + L_5 \end{aligned}$$

Therefore, we have seven 2-dimensional subspaces, which we will call “type 2” vertices. Note that each V_i contains exactly three subspaces.

$$V_1 = L_1 + L_2 = L_1 + L_4 = L_2 + L_4$$


Buildings

For each $\Sigma(\{L_i, L_j, L_k\})$ such that $V = L_i \oplus L_j \oplus L_k$, we obtain an apartment:



Buildings

For each $\Sigma(\{L_i, L_j, L_k\})$ such that $V = L_i \oplus L_j \oplus L_k$, we obtain an apartment:




Chambers are edges with a type 1 and type 2 vertex, S_3 acts by permuting the labels $\{i, j, k\}$.

Buildings

The axioms **B1** and **B2** of a building give us criteria of how to glue.


Buildings

The axioms **B1** and **B2** of a building give us criteria of how to glue.



Buildings

The axioms **B1** and **B2** of a building give us criteria of how to glue.



The apartment $\Sigma(\{L_1, L_2, L_3\})$ is shown above.

Important Question:

Buildings

Important Question: What exactly was Jacques Tits trying to construct?

Buildings

Important Question: What exactly was Jacques Tits trying to construct? Consider the group $\mathrm{SL}(2, \mathbb{C})$, it has a maximal compact subgroup $\mathrm{SU}(2)$.

Important Question: What exactly was Jacques Tits trying to construct? Consider the group $SL(2, \mathbb{C})$, it has a maximal compact subgroup $SU(2)$. Note that $SL(2, \mathbb{C})/SU(2) \cong \mathbb{H} = \{z \in \mathbb{C} : \text{Im}(z) > 0\}$ via the map

$$SL(2, \mathbb{C})/SU(2) \rightarrow \mathbb{H}$$
$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \frac{ai + b}{ci + d}$$

Important Question: What exactly was Jacques Tits trying to construct? Consider the group $SL(2, \mathbb{C})$, it has a maximal compact subgroup $SU(2)$. Note that $SL(2, \mathbb{C})/SU(2) \cong \mathbb{H} = \{z \in \mathbb{C} : \text{Im}(z) > 0\}$ via the map

$$SL(2, \mathbb{C})/SU(2) \rightarrow \mathbb{H}$$
$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \frac{ai + b}{ci + d}$$

And in general, if we take a semisimple group G over \mathbb{R} and quotient by its maximal compact subgroup K , G/K is a nice geometric space.

Buildings

Important Question: What exactly was Jacques Tits trying to construct? Consider the group $\mathrm{SL}(2, \mathbb{C})$, it has a maximal compact subgroup $\mathrm{SU}(2)$. Note that $\mathrm{SL}(2, \mathbb{C})/\mathrm{SU}(2) \cong \mathbb{H} = \{z \in \mathbb{C} : \mathrm{Im}(z) > 0\}$ via the map

$$\mathrm{SL}(2, \mathbb{C})/\mathrm{SU}(2) \rightarrow \mathbb{H}$$
$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \frac{ai + b}{ci + d}$$

And in general, if we take a semisimple group G over \mathbb{R} and quotient by its maximal compact subgroup K , G/K is a nice geometric space.

However, if we consider a local field $F = \mathbb{Q}_p$ and the algebraic group $\mathrm{SL}(2, F)$, its maximal compact subgroup is $\mathrm{SL}(2, \mathcal{O}_F)$ and $\mathrm{SL}(2, F)/\mathrm{SL}(2, \mathcal{O}_F)$, it is a totally disconnected topological space.

Buildings

Important Question: What exactly was Jacques Tits trying to construct? Consider the group $\mathrm{SL}(2, \mathbb{C})$, it has a maximal compact subgroup $\mathrm{SU}(2)$. Note that $\mathrm{SL}(2, \mathbb{C})/\mathrm{SU}(2) \cong \mathbb{H} = \{z \in \mathbb{C} : \mathrm{Im}(z) > 0\}$ via the map

$$\mathrm{SL}(2, \mathbb{C})/\mathrm{SU}(2) \rightarrow \mathbb{H}$$
$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \frac{ai + b}{ci + d}$$

And in general, if we take a semisimple group G over \mathbb{R} and quotient by its maximal compact subgroup K , G/K is a nice geometric space.

However, if we consider a local field $F = \mathbb{Q}_p$ and the algebraic group $\mathrm{SL}(2, F)$, its maximal compact subgroup is $\mathrm{SL}(2, \mathcal{O}_F)$ and $\mathrm{SL}(2, F)/\mathrm{SL}(2, \mathcal{O}_F)$, it is a totally disconnected topological space. It is a *pretty lame space*.

Buildings

This occurs as a property of non-archimedean fields, i.e. the natural numbers are bounded under the absolute value on the field.

Buildings

This occurs as a property of non-archimedean fields, i.e. the natural numbers are bounded under the absolute value on the field.

In the case of $SL(2, \mathbb{C})$, we obtain a Riemannian symmetric space and we want an analogue to this when the field is non-archimedean.

Buildings

This occurs as a property of non-archimedean fields, i.e. the natural numbers are bounded under the absolute value on the field.

In the case of $SL(2, \mathbb{C})$, we obtain a Riemannian symmetric space and we want an analogue to this when the field is non-archimedean. A Bruhat-Tits building is the correct analogue. Historically, the following building appeared as a way to prove that every torsion-free subgroup of $SL(2, \mathbb{Q}_p)$ is a free group (Ihara, 1966).

Buildings

This occurs as a property of non-archimedean fields, i.e. the natural numbers are bounded under the absolute value on the field.

In the case of $SL(2, \mathbb{C})$, we obtain a Riemannian symmetric space and we want an analogue to this when the field is non-archimedean. A Bruhat-Tits building is the correct analogue. Historically, the following building appeared as a way to prove that every torsion-free subgroup of $SL(2, \mathbb{Q}_p)$ is a free group (Ihara, 1966). To make the construction concrete, we'll use $SL(2, \mathbb{Q}_p)$ as our algebraic group.

Buildings

This occurs as a property of non-archimedean fields, i.e. the natural numbers are bounded under the absolute value on the field.

In the case of $SL(2, \mathbb{C})$, we obtain a Riemannian symmetric space and we want an analogue to this when the field is non-archimedean. A Bruhat-Tits building is the correct analogue. Historically, the following building appeared as a way to prove that every torsion-free subgroup of $SL(2, \mathbb{Q}_p)$ is a free group (Ihara, 1966). To make the construction concrete, we'll use $SL(2, \mathbb{Q}_p)$ as our algebraic group.

Note: Any element of \mathbb{Q}_p is of the form:

$$x = \sum_{i=k}^{\infty} a_i p^i$$

for $0 \leq a_i < p$. The valuation of x , is $\nu(x) = k$, where k is the smallest index such that $a_k \neq 0$.

Definition: A non-archimedean absolute value on a field F is a map

$$| - | : F \rightarrow \mathbb{R}$$

such that

- (1) $|x| \geq 0$ for all $x \in F$, and $|x| = 0$ if and only if $x = 0$.
- (2) $|xy| = |x||y|$ for all $x, y \in F$
- (3) $|x + y| \leq \max\{|x|, |y|\}$ for all $x, y \in F$.

Definition: A non-archimedean absolute value on a field F is a map

$$| - | : F \rightarrow \mathbb{R}$$

such that

- (1) $|x| \geq 0$ for all $x \in F$, and $|x| = 0$ if and only if $x = 0$.
- (2) $|xy| = |x||y|$ for all $x, y \in F$
- (3) $|x + y| \leq \max\{|x|, |y|\}$ for all $x, y \in F$.

Example: For \mathbb{Q}_p , the absolute value is defined by $|x|_p = p^{-\nu(x)}$.

Definition: A non-archimedean absolute value on a field F is a map

$$| - | : F \rightarrow \mathbb{R}$$

such that

- (1) $|x| \geq 0$ for all $x \in F$, and $|x| = 0$ if and only if $x = 0$.
- (2) $|xy| = |x||y|$ for all $x, y \in F$
- (3) $|x + y| \leq \max\{|x|, |y|\}$ for all $x, y \in F$.

Example: For \mathbb{Q}_p , the absolute value is defined by $|x|_p = p^{-\nu(x)}$.

Important Note: The image of $\nu(x)$ is an integer and thus $\nu(\mathbb{Q}_p^\times) = \mathbb{Z}$ is discrete. Or, in particular importance for us, $\log(|\mathbb{Q}_p^\times|_p)$ is discrete. And, $\mathcal{O} = \{x \in \mathbb{Q}_p : |x|_p \leq 1\}$.

Buildings

Instead of choosing the maximal compact subgroup, we look at tori. The torus will be

$$T = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} : a \in \mathbb{Q}_p^\times \right\}$$

Buildings


Instead of choosing the maximal compact subgroup, we look at tori. The torus will be

$$T = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} : a \in \mathbb{Q}_p^\times \right\}$$

Let $A(T)$ be the apartment corresponding to T , it is an \mathbb{R} affine space with simplicial structure defined by the map

$$T \rightarrow \mathbb{R}$$

$$\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \mapsto \log(|a|_p)$$




Buildings

Instead of choosing the maximal compact subgroup, we look at tori. The torus will be

$$T = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} : a \in \mathbb{Q}_p^\times \right\}$$

Let $A(T)$ be the apartment corresponding to T , it is an \mathbb{R} affine space with simplicial structure defined by the map

$$\begin{aligned} T &\rightarrow \mathbb{R} \\ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} &\mapsto \log(|a|_p) \end{aligned}$$


Our collection of vertices are of the form: $n \log(p)$ for some $n \in \mathbb{Z}$.

Buildings

We then take the collection of tori $\{gTg^{-1} : g \in \mathrm{SL}_2(\mathbb{Q}_p)\}$ and glue together the corresponding system of apartments

$$\mathcal{A} = \{A(gTg^{-1}) : g \in \mathrm{SL}_2(\mathbb{Q}_p)\}$$

Buildings

We then take the collection of tori $\{gTg^{-1} : g \in \mathrm{SL}_2(\mathbb{Q}_p)\}$ and glue together the corresponding system of apartments

$$\mathcal{A} = \{A(gTg^{-1}) : g \in \mathrm{SL}_2(\mathbb{Q}_p)\}$$


The stabilizers of each vertex is of the form

$$\mathrm{Stab}(\log(|a|_p)) = \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \cdot \mathrm{SL}_2(\mathcal{O}) \cdot \begin{pmatrix} a^{-1} & 0 \\ 0 & a \end{pmatrix}$$

and for any two apartments $A(T)$ and $g \cdot A(T) = A(gTg^{-1})$, we glue at $n \log(p)$ if and only if $g \in \mathrm{Stab}(n \log(p))$.


Buildings

Example: Let $g = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. For $a = up^{-n}$, note that $g \in \text{Stab}(n \log(p))$ if and only if $|a|_p \geq 1$ if and only if $n \log(p) \geq 0$.




Buildings

Example: Let $g = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. For $a = up^{-n}$, note that $g \in \text{Stab}(n \log(p))$ if and only if $|a|_p \geq 1$ if and only if $n \log(p) \geq 0$.




Buildings

Assume that $p = 2$, then the building is given by:



Buildings


Assume that $p = 2$, then the building is given by:



An apartment $A(gTg^{-1})$ is given by an infinite path along the tree.


Buildings

Similarly for $p = 3$, then the building is given by:



Buildings

Similarly for $p = 3$, then the building is given by:



Again, an apartment $A(gTg^{-1})$ is given by an infinite path along the tree.

Bibliography



Peter Abramenko, Kenneth S. Brown. *Buildings - Theory and Applications*. Springer Science+Business Media, LLC, 2008.



Jean-Pierre Serre. *Trees*. Springer-Verlag Berlin Heidelberg, 1980.
(Translated by John Stillwell)



Annette Werner. *Buildings and tropical geometry [Video]*. Mathematical Sciences Research Institute, 2009.



Annette Werner. *A tropical view on Bruhat-Tits buildings and their compactifications*. Central European Journal of Mathematics, 2010.



Bertrand Rémy, Amaury Thuillier, Annette Werner. *Bruhat-Tits Buildings and Analytic Geometry* Springer International Publishing Switzerland, 2015.



Federico Ardila. *Coxeter groups [Online Lecture Series]*. San Francisco State University, 2008.



Dmitry Kozlov. *Combinatorial Algebraic Topology*. Springer-Verlag Berlin Heidelberg, 2008.