

# 1.5 Centre and commutator subspaces

Reading Seminar, Linckelmann Chapter 1

---

Deewang Bhamidipati

15th July 2021

## Theorem 1.5.1

If  $G$  is a finite group the centre of the group algebra  $kG$  admits a canonical  $k$ -basis, namely its conjugacy class sums.

## Theorem 1.5.1

If  $G$  is a finite group the centre of the group algebra  $kG$  admits a canonical  $k$ -basis, namely its conjugacy class sums.

### Theorem 1.5.1

Let  $G$  be a finite group. Let  $\mathcal{K}$  be the set of conjugacy classes in  $G$ , and for any conjugacy class  $C$  in  $\mathcal{K}$  denote by  $\bar{C}$  the conjugacy class sum  $\bar{C} = \sum_{y \in C} y$  in  $kG$ .

## Theorem 1.5.1

If  $G$  is a finite group the centre of the group algebra  $kG$  admits a canonical  $k$ -basis, namely its conjugacy class sums.

### Theorem 1.5.1

Let  $G$  be a finite group. Let  $\mathcal{K}$  be the set of conjugacy classes in  $G$ , and for any conjugacy class  $C$  in  $\mathcal{K}$  denote by  $\bar{C}$  the conjugacy class sum  $\bar{C} = \sum_{y \in C} y$  in  $kG$ . The set  $\{\bar{C} : C \in \mathcal{K}\}$  is a  $k$ -basis of  $Z(kG)$ ,

## Theorem 1.5.1

If  $G$  is a finite group the centre of the group algebra  $kG$  admits a canonical  $k$ -basis, namely its conjugacy class sums.

### Theorem 1.5.1

Let  $G$  be a finite group. Let  $\mathcal{K}$  be the set of conjugacy classes in  $G$ , and for any conjugacy class  $C$  in  $\mathcal{K}$  denote by  $\overline{C}$  the conjugacy class sum  $\overline{C} = \sum_{y \in C} y$  in  $kG$ . The set  $\{\overline{C} : C \in \mathcal{K}\}$  is a  $k$ -basis of  $Z(kG)$ , and for any two  $C, C' \in \mathcal{K}$  there are non-negative integers  $a_D$  for all  $D \in \mathcal{K}$  such that  $\overline{C} \cdot \overline{C'} = \sum_{D \in \mathcal{K}} a_D \overline{D}$ .

## Theorem 1.5.1

If  $G$  is a finite group the centre of the group algebra  $kG$  admits a canonical  $k$ -basis, namely its conjugacy class sums.

### Theorem 1.5.1

Let  $G$  be a finite group. Let  $\mathcal{K}$  be the set of conjugacy classes in  $G$ , and for any conjugacy class  $C$  in  $\mathcal{K}$  denote by  $\overline{C}$  the conjugacy class sum  $\overline{C} = \sum_{y \in C} y$  in  $kG$ . The set  $\{\overline{C} : C \in \mathcal{K}\}$  is a  $k$ -basis of  $Z(kG)$ , and for any two  $C, C' \in \mathcal{K}$  there are non-negative integers  $a_D$  for all  $D \in \mathcal{K}$  such that  $\overline{C} \cdot \overline{C'} = \sum_{D \in \mathcal{K}} a_D \overline{D}$ .

In particular,  $Z(kG)$  is free as a  $k$ -module with rank equal to the number of conjugacy classes in  $G$ .

## Proposition 1.5.2

## Proposition 1.5.2

### Proposition 1.5.2

Let  $G, H$  be finite groups. There is a unique algebra isomorphism

$$Z(k(G \times H)) \cong Z(kG) \otimes_k Z(kH)$$



## Proposition 1.5.2

$$C_{x,y} = \sum_{z \sim \text{conjugate to } (x,y)} z \mapsto C_x \otimes C_y$$

### Proposition 1.5.2

Let  $G, H$  be finite groups. There is a unique algebra isomorphism

$$Z(k(G \times H)) \cong Z(kG) \otimes_k Z(kH)$$

sending the conjugacy class sum of  $(x, y) \in G \times H$  in  $k(G \times H)$  to  $C_x \otimes C_y$ , where  $C_x, C_y$  are the conjugacy class sums of  $x$  and  $y$  in  $kG$  and  $kH$ , respectively.

## Proposition 1.5.2

$$\begin{aligned} k(G \times H) &\longrightarrow kG \otimes_k kH \\ (x, y) &\longmapsto x \otimes y \end{aligned}$$

### Proposition 1.5.2

Let  $G, H$  be finite groups. There is a unique algebra isomorphism

$$Z(k(G \times H)) \cong Z(kG) \otimes_k Z(kH)$$



sending the conjugacy class sum of  $(x, y) \in G \times H$  in  $k(G \times H)$  to  $C_x \otimes C_y$ , where  $C_x, C_y$  are the conjugacy class sums of  $x$  and  $y$  in  $kG$  and  $kH$ , respectively.

### *Proof*

The given map is an algebra homomorphism, induced by the algebra isomorphism in 1.1.4, restricted to the centre of  $k(G \times H)$ . The explicit description of the centre of a finite group algebra in terms of conjugacy class sums 1.5.1 implies that this induces an isomorphism as stated.

## Theorem 1.5.3

The fact that taking centres commutes with taking tensor products of algebras as in the previous proposition holds in a bit more generality.

## Theorem 1.5.3

The fact that taking centres commutes with taking tensor products of algebras as in the previous proposition holds in a bit more generality.

### Theorem 1.5.1

Let  $A, B$  be  $k$ -algebras.

## Theorem 1.5.3

The fact that taking centres commutes with taking tensor products of algebras as in the previous proposition holds in a bit more generality.

### Theorem 1.5.1

Let  $A, B$  be  $k$ -algebras. Suppose that  $A$  and  $Z(B)$  are free as  $k$ -modules

## Theorem 1.5.3

The fact that taking centres commutes with taking tensor products of algebras as in the previous proposition holds in a bit more generality.

### Theorem 1.5.1

Let  $A, B$  be  $k$ -algebras. Suppose that  $A$  and  $Z(B)$  are free as  $k$ -modules and that  $Z(B)$  is a direct summand of  $B$  as a  $k$ -module.

$$kG = Z(kG) \oplus \dots$$

Alg 2  
- June 2021

## Theorem 1.5.3

The fact that taking centres commutes with taking tensor products of algebras as in the previous proposition holds in a bit more generality.

### Theorem 1.5.1

Let  $A, B$  be  $k$ -algebras. Suppose that  $A$  and  $Z(B)$  are free as  $k$ -modules and that  $Z(B)$  is a direct summand of  $B$  as a  $k$ -module. Then the centraliser of the subalgebra  $1 \otimes B$  of  $A \otimes_k B$  is  $A \otimes_k Z(B)$ ,

## Theorem 1.5.3

The fact that taking centres commutes with taking tensor products of algebras as in the previous proposition holds in a bit more generality.

### Theorem 1.5.1

Let  $A, B$  be  $k$ -algebras. Suppose that  $A$  and  $Z(B)$  are free as  $k$ -modules and that  $Z(B)$  is a direct summand of  $B$  as a  $k$ -module. Then the centraliser of the subalgebra  $1 \otimes B$  of  $A \otimes_k B$  is  $A \otimes_k Z(B)$ , and we have  $Z(A \otimes_k B) = Z(A) \otimes_k Z(B)$ .



## Theorem 1.5.3

The fact that taking centres commutes with taking tensor products of algebras as in the previous proposition holds in a bit more generality.

### Theorem 1.5.1

Let  $A, B$  be  $k$ -algebras. Suppose that  $A$  and  $Z(B)$  are free as  $k$ -modules and that  $Z(B)$  is a direct summand of  $B$  as a  $k$ -module. Then the centraliser of the subalgebra  $1 \otimes B$  of  $A \otimes_k B$  is  $A \otimes_k Z(B)$ , and we have  $Z(A \otimes_k B) = Z(A) \otimes_k Z(B)$ .

#### *Proof*

By assumption  $Z(B)$  is a direct summand of  $B$ , as  $k$ -modules, so we can identify  $A \otimes_k Z(B)$  as a direct summand of  $A \otimes_k B$ . Similarly, we identify  $Z(A) \otimes_k Z(B)$  with its image in  $A \otimes_k B$ .

$$B = Z(B) \oplus C$$

$$A \otimes B = A \otimes Z(B) \oplus A \otimes C$$

## Theorem 1.5.3

The fact that taking centres commutes with taking tensor products of algebras as in the previous proposition holds in a bit more generality.

### Theorem 1.5.1

Let  $A, B$  be  $k$ -algebras. Suppose that  $A$  and  $Z(B)$  are free as  $k$ -modules and that  $Z(B)$  is a direct summand of  $B$  as a  $k$ -module. Then the centraliser of the subalgebra  $1 \otimes B$  of  $A \otimes_k B$  is  $A \otimes_k Z(B)$ , and we have  $Z(A \otimes_k B) = Z(A) \otimes_k Z(B)$ .

#### *Proof*

By assumption  $Z(B)$  is a direct summand of  $B$ , as  $k$ -modules, so we can identify  $A \otimes_k Z(B)$  as a direct summand of  $A \otimes_k B$ . Similarly, we identify  $Z(A) \otimes_k Z(B)$  with its image in  $A \otimes_k B$ .

Now, naturally  $A \otimes_k Z(B)$  centralises  $1 \otimes B$  and  $Z(A) \otimes_k Z(B)$  is contained in  $Z(A \otimes_k B)$ .

$$\begin{aligned} (a \otimes z)(1 \otimes b) &= (1 \otimes b)(a \otimes z) \\ \underline{a} \otimes \underline{z}b &= \underline{a} \otimes \underline{bz} \end{aligned} \quad A \otimes Z(B) \subseteq \text{Cent}(1 \otimes B)$$

## Theorem 1.5.3

The fact that taking centres commutes with taking tensor products of algebras as in the previous proposition holds in a bit more generality.

### Theorem 1.5.1

Let  $A, B$  be  $k$ -algebras. Suppose that  $A$  and  $Z(B)$  are free as  $k$ -modules and that  $Z(B)$  is a direct summand of  $B$  as a  $k$ -module. Then the centraliser of the subalgebra  $1 \otimes B$  of  $A \otimes_k B$  is  $A \otimes_k Z(B)$ , and we have  $Z(A \otimes_k B) = Z(A) \otimes_k Z(B)$ .

#### *Proof*

By assumption  $Z(B)$  is a direct summand of  $B$ , as  $k$ -modules, so we can identify  $A \otimes_k Z(B)$  as a direct summand of  $A \otimes_k B$ . Similarly, we identify  $Z(A) \otimes_k Z(B)$  with its image in  $A \otimes_k B$ .

Now, naturally  $A \otimes_k Z(B)$  centralises  $1 \otimes B$  and  $Z(A) \otimes_k Z(B)$  is contained in  $Z(A \otimes_k B)$ .

To show the reverse inclusions, we employ a  $k$ -basis of our free modules.

## Theorem 1.5.3

The fact that taking centres commutes with taking tensor products of algebras as in the previous proposition holds in a bit more generality.

### Theorem 1.5.1

Let  $A, B$  be  $k$ -algebras. Suppose that  $A$  and  $Z(B)$  are free as  $k$ -modules and that  $Z(B)$  is a direct summand of  $B$  as a  $k$ -module. Then the centraliser of the subalgebra  $1 \otimes B$  of  $A \otimes_k B$  is  $A \otimes_k Z(B)$ , and we have  $Z(A \otimes_k B) = Z(A) \otimes_k Z(B)$ .

#### *Proof*

By assumption  $Z(B)$  is a direct summand of  $B$ , as  $k$ -modules, so we can identify  $A \otimes_k Z(B)$  as a direct summand of  $A \otimes_k B$ . Similarly, we identify  $Z(A) \otimes_k Z(B)$  with its image in  $A \otimes_k B$ .

Now, naturally  $A \otimes_k Z(B)$  centralises  $1 \otimes B$  and  $Z(A) \otimes_k Z(B)$  is contained in  $Z(A \otimes_k B)$ .

To show the reverse inclusions, we employ a  $k$ -basis of our free modules. Let  $\{a_i\}_{i \in I}$  be a  $k$ -basis of  $A$  and let  $\{z_j\}_{j \in J}$  be a  $k$ -basis of  $Z(B)$ .

## Theorem 1.5.3

The fact that taking centres commutes with taking tensor products of algebras as in the previous proposition holds in a bit more generality.

### Theorem 1.5.1

Let  $A, B$  be  $k$ -algebras. Suppose that  $A$  and  $Z(B)$  are free as  $k$ -modules and that  $Z(B)$  is a direct summand of  $B$  as a  $k$ -module. Then the centraliser of the subalgebra  $1 \otimes B$  of  $A \otimes_k B$  is  $A \otimes_k Z(B)$ , and we have  $Z(A \otimes_k B) = Z(A) \otimes_k Z(B)$ .

#### *Proof*

By assumption  $Z(B)$  is a direct summand of  $B$ , as  $k$ -modules, so we can identify  $A \otimes_k Z(B)$  as a direct summand of  $A \otimes_k B$ . Similarly, we identify  $Z(A) \otimes_k Z(B)$  with its image in  $A \otimes_k B$ .

Now, naturally  $A \otimes_k Z(B)$  centralises  $1 \otimes B$  and  $Z(A) \otimes_k Z(B)$  is contained in  $Z(A \otimes_k B)$ .

To show the reverse inclusions, we employ a  $k$ -basis of our free modules. Let  $\{a_i\}_{i \in I}$  be a  $k$ -basis of  $A$  and let  $\{z_j\}_{j \in J}$  be a  $k$ -basis of  $Z(B)$ . Then the set  $\{a_i \otimes 1_B\}_{i \in I}$  is a  $B$ -basis of  $A \otimes_k B$  as a right  $B$ -module,

## Theorem 1.5.3

The fact that taking centres commutes with taking tensor products of algebras as in the previous proposition holds in a bit more generality.

### Theorem 1.5.1

Let  $A, B$  be  $k$ -algebras. Suppose that  $A$  and  $Z(B)$  are free as  $k$ -modules and that  $Z(B)$  is a direct summand of  $B$  as a  $k$ -module. Then the centraliser of the subalgebra  $1 \otimes B$  of  $A \otimes_k B$  is  $A \otimes_k Z(B)$ , and we have  $Z(A \otimes_k B) = Z(A) \otimes_k Z(B)$ .

#### *Proof*

By assumption  $Z(B)$  is a direct summand of  $B$ , as  $k$ -modules, so we can identify  $A \otimes_k Z(B)$  as a direct summand of  $A \otimes_k B$ . Similarly, we identify  $Z(A) \otimes_k Z(B)$  with its image in  $A \otimes_k B$ .

Now, naturally  $A \otimes_k Z(B)$  centralises  $1 \otimes B$  and  $Z(A) \otimes_k Z(B)$  is contained in  $Z(A \otimes_k B)$ .

To show the reverse inclusions, we employ a  $k$ -basis of our free modules. Let  $\{a_i\}_{i \in I}$  be a  $k$ -basis of  $A$  and let  $\{z_j\}_{j \in J}$  be a  $k$ -basis of  $Z(B)$ . Then the set  $\{a_i \otimes 1_B\}_{i \in I}$  is a  $B$ -basis of  $A \otimes_k B$  as a right  $B$ -module, and the set  $\{1_A \otimes z_j\}_{j \in J}$  is an  $A$ -basis of  $A \otimes_k Z(B)$  as a left  $A$ -module.

## Theorem 1.5.3

$$A = \bigoplus k a_i \quad A \otimes B = \bigoplus k a_i \otimes b_j$$

(contd.)

Let  $z \in A \otimes_k B$ .

## Theorem 1.5.3

(contd.)

Let  $z \in A \otimes_k B$ . By the above,  $z$  can be written in the form

$$z = \sum_{i \in I} a_i \otimes b_i = \sum (a_i \otimes 1) b_i$$

with uniquely determined elements  $b_i \in B$  of which only finitely many are nonzero.



## Theorem 1.5.3

(contd.)

Let  $z \in A \otimes_k B$ . By the above,  $z$  can be written in the form

$$z = \sum_{i \in I} a_i \otimes b_i$$

with uniquely determined elements  $b_i \in B$  of which only finitely many are nonzero. Suppose that  $z$  centralises  $1 \otimes B$ . This is equivalent to  $(1_A \otimes b)z = z(1_A \otimes b)$  for all  $b \in B$ ,

## Theorem 1.5.3

(contd.)

Let  $z \in A \otimes_k B$ . By the above,  $z$  can be written in the form

$$z = \sum_{i \in I} a_i \otimes b_i$$

with uniquely determined elements  $b_i \in B$  of which only finitely many are nonzero. Suppose that  $z$  centralises  $1 \otimes B$ . This is equivalent to  $(1_A \otimes b)z = z(1_A \otimes b)$  for all  $b \in B$ , which is equivalent to  $\sum_{i \in I} a_i \otimes bb_i = \sum_{i \in I} a_i \otimes b_i b$  for all  $b \in B$ .

## Theorem 1.5.3

(contd.)

Let  $z \in A \otimes_k B$ . By the above,  $z$  can be written in the form

$$z = \sum_{i \in I} a_i \otimes b_i$$

with uniquely determined elements  $b_i \in B$  of which only finitely many are nonzero. Suppose that  $z$  centralises  $1 \otimes B$ . This is equivalent to  $(1_A \otimes b)z = z(1_A \otimes b)$  for all  $b \in B$ , which is equivalent to  $\sum_{i \in I} a_i \otimes bb_i = \sum_{i \in I} a_i \otimes b_i b$  for all  $b \in B$ . The uniqueness of expressions of this form implies that this is equivalent to  $bb_i = b_i b$  for all  $i \in I$  and all  $b \in B$ .

## Theorem 1.5.3

(contd.)

Let  $z \in A \otimes_k B$ . By the above,  $z$  can be written in the form

$$z = \sum_{i \in I} a_i \otimes b_i$$

with uniquely determined elements  $b_i \in B$  of which only finitely many are nonzero. Suppose that  $z$  centralises  $1 \otimes B$ . This is equivalent to  $(1_A \otimes b)z = z(1_A \otimes b)$  for all  $b \in B$ , which is equivalent to  $\sum_{i \in I} a_i \otimes bb_i = \sum_{i \in I} a_i \otimes b_i b$  for all  $b \in B$ . The uniqueness of expressions of this form implies that this is equivalent to  $bb_i = b_i b$  for all  $i \in I$  and all  $b \in B$ . Thus  $b_i \in Z(B)$  for  $i \in I$ , and hence  $z \in A \otimes_k Z(B)$ . This shows the first statement.

## Theorem 1.5.3

(contd.)

Let  $z \in A \otimes_k B$ . By the above,  $z$  can be written in the form

$$z = \sum_{i \in I} a_i \otimes b_i$$

with uniquely determined elements  $b_i \in B$  of which only finitely many are nonzero. Suppose that  $z$  centralises  $1 \otimes B$ . This is equivalent to  $(1_A \otimes b)z = z(1_A \otimes b)$  for all  $b \in B$ , which is equivalent to  $\sum_{i \in I} a_i \otimes bb_i = \sum_{i \in I} a_i \otimes b_i b$  for all  $b \in B$ . The uniqueness of expressions of this form implies that this is equivalent to  $bb_i = b_i b$  for all  $i \in I$  and all  $b \in B$ . Thus  $b_i \in Z(B)$  for  $i \in I$ , and hence  $z \in A \otimes_k Z(B)$ . This shows the first statement.

Suppose that  $z \in Z(A \otimes_k B)$ .

## Theorem 1.5.3

(contd.)

Let  $z \in A \otimes_k B$ . By the above,  $z$  can be written in the form

$$z = \sum_{i \in I} a_i \otimes b_i$$

with uniquely determined elements  $b_i \in B$  of which only finitely many are nonzero. Suppose that  $z$  centralises  $1 \otimes B$ . This is equivalent to  $(1_A \otimes b)z = z(1_A \otimes b)$  for all  $b \in B$ , which is equivalent to  $\sum_{i \in I} a_i \otimes bb_i = \sum_{i \in I} a_i \otimes b_i b$  for all  $b \in B$ . The uniqueness of expressions of this form implies that this is equivalent to  $bb_i = b_i b$  for all  $i \in I$  and all  $b \in B$ . Thus  $b_i \in Z(B)$  for  $i \in I$ , and hence  $z \in A \otimes_k Z(B)$ . This shows the first statement.

Suppose that  $z \in Z(A \otimes_k B)$ . By the first statement, we have  $z \in A \otimes_k Z(B)$ .

## Theorem 1.5.3

(contd.)

Let  $z \in A \otimes_k B$ . By the above,  $z$  can be written in the form

$$z = \sum_{i \in I} a_i \otimes b_i$$

with uniquely determined elements  $b_i \in B$  of which only finitely many are nonzero. Suppose that  $z$  centralises  $1 \otimes B$ . This is equivalent to  $(1_A \otimes b)z = z(1_A \otimes b)$  for all  $b \in B$ , which is equivalent to  $\sum_{i \in I} a_i \otimes bb_i = \sum_{i \in I} a_i \otimes b_i b$  for all  $b \in B$ . The uniqueness of expressions of this form implies that this is equivalent to  $bb_i = b_i b$  for all  $i \in I$  and all  $b \in B$ . Thus  $b_i \in Z(B)$  for  $i \in I$ , and hence  $z \in A \otimes_k Z(B)$ . This shows the first statement.

Suppose that  $z \in Z(A \otimes_k B)$ . By the first statement, we have  $z \in A \otimes_k Z(B)$ . Thus  $z$  can be written in the form

$$z = \sum_{j \in J} c_j \otimes z_j = \sum c_j (1 \otimes z_j)$$

with uniquely determined elements  $c_j \in A$  of which at most finitely many are nonzero.

## Theorem 1.5.3

(contd.)

Let  $z \in A \otimes_k B$ . By the above,  $z$  can be written in the form

$$z = \sum_{i \in I} a_i \otimes b_i$$

with uniquely determined elements  $b_i \in B$  of which only finitely many are nonzero. Suppose that  $z$  centralises  $1 \otimes B$ . This is equivalent to  $(1_A \otimes b)z = z(1_A \otimes b)$  for all  $b \in B$ , which is equivalent to  $\sum_{i \in I} a_i \otimes bb_i = \sum_{i \in I} a_i \otimes b_i b$  for all  $b \in B$ . The uniqueness of expressions of this form implies that this is equivalent to  $bb_i = b_i b$  for all  $i \in I$  and all  $b \in B$ . Thus  $b_i \in Z(B)$  for  $i \in I$ , and hence  $z \in A \otimes_k Z(B)$ . This shows the first statement.

Suppose that  $z \in Z(A \otimes_k B)$ . By the first statement, we have  $z \in A \otimes_k Z(B)$ . Thus  $z$  can be written in the form

$$z = \sum_{j \in J} c_j \otimes z_j$$

with uniquely determined elements  $c_j \in A$  of which at most finitely many are nonzero. Then the equality  $(a \otimes 1_B)z = z(a \otimes 1_B)$  for all  $a \in A$



## Theorem 1.5.3

(contd.)

Let  $z \in A \otimes_k B$ . By the above,  $z$  can be written in the form

$$z = \sum_{i \in I} a_i \otimes b_i$$

with uniquely determined elements  $b_i \in B$  of which only finitely many are nonzero. Suppose that  $z$  centralises  $1 \otimes B$ . This is equivalent to  $(1_A \otimes b)z = z(1_A \otimes b)$  for all  $b \in B$ , which is equivalent to  $\sum_{i \in I} a_i \otimes bb_i = \sum_{i \in I} a_i \otimes b_i b$  for all  $b \in B$ . The uniqueness of expressions of this form implies that this is equivalent to  $bb_i = b_i b$  for all  $i \in I$  and all  $b \in B$ . Thus  $b_i \in Z(B)$  for  $i \in I$ , and hence  $z \in A \otimes_k Z(B)$ . This shows the first statement.

Suppose that  $z \in Z(A \otimes_k B)$ . By the first statement, we have  $z \in A \otimes_k Z(B)$ . Thus  $z$  can be written in the form

$$z = \sum_{j \in J} c_j \otimes z_j$$

with uniquely determined elements  $c_j \in A$  of which at most finitely many are nonzero. Then the equality  $(a \otimes 1_B)z = z(a \otimes 1_B)$  for all  $a \in A$  is equivalent to  $ac_j = c_j a$  for all  $a \in A$ ,

## Theorem 1.5.3

(contd.)

Let  $z \in A \otimes_k B$ . By the above,  $z$  can be written in the form

$$z = \sum_{i \in I} a_i \otimes b_i$$

with uniquely determined elements  $b_i \in B$  of which only finitely many are nonzero. Suppose that  $z$  centralises  $1 \otimes B$ . This is equivalent to  $(1_A \otimes b)z = z(1_A \otimes b)$  for all  $b \in B$ , which is equivalent to  $\sum_{i \in I} a_i \otimes bb_i = \sum_{i \in I} a_i \otimes b_i b$  for all  $b \in B$ . The uniqueness of expressions of this form implies that this is equivalent to  $bb_i = b_i b$  for all  $i \in I$  and all  $b \in B$ . Thus  $b_i \in Z(B)$  for  $i \in I$ , and hence  $z \in A \otimes_k Z(B)$ . This shows the first statement.

Suppose that  $z \in Z(A \otimes_k B)$ . By the first statement, we have  $z \in A \otimes_k Z(B)$ . Thus  $z$  can be written in the form

$$z = \sum_{j \in J} c_j \otimes z_j$$

with uniquely determined elements  $c_j \in A$  of which at most finitely many are nonzero. Then the equality  $(a \otimes 1_B)z = z(a \otimes 1_B)$  for all  $a \in A$  is equivalent to  $ac_j = c_j a$  for all  $a \in A$ , hence  $c_j \in Z(A)$  for all  $j \in J$ , which proves that  $z \in Z(A) \otimes_k Z(B)$ . The result follows.

## Remarks A.

The centre of a twisted group algebra  $k_\alpha G$  does not in general admit a basis as in 1.5.1,

## Remarks A.

The centre of a twisted group algebra  $k_\alpha G$  does not in general admit a basis as in 1.5.1, essentially because the multiplicative inverse in  $k_\alpha G$  of a group element  $x$  need not be equal to the inverse  $x^{-1}$  of  $x$  as an element in the group  $G$ .

$x \in G$   $x^{-1} \in G$  is NOT  $\hookrightarrow \alpha(x) x \dots$  the inverse of  $x$  in  $k_\alpha G$ .

## Remarks A.

The centre of a twisted group algebra  $k_\alpha G$  does not in general admit a basis as in 1.5.1, essentially because the multiplicative inverse in  $k_\alpha G$  of a group element  $x$  need not be equal to the inverse  $x^{-1}$  of  $x$  as an element in the group  $G$ .

Another measure for how far a  $k$ -algebra  $A$  is from being commutative is the *commutator space*  $[A, A]$ .

## Remarks A.

The centre of a twisted group algebra  $k_\alpha G$  does not in general admit a basis as in 1.5.1, essentially because the multiplicative inverse in  $k_\alpha G$  of a group element  $x$  need not be equal to the inverse  $x^{-1}$  of  $x$  as an element in the group  $G$ .

Another measure for how far a  $k$ -algebra  $A$  is from being commutative is the *commutator space*  $[A, A]$ .

- This is the  $k$ -submodule of  $A$  generated by all *additive commutators*  $[a, b] = ab - ba$ , where  $a, b \in A$ .

## Remarks A.

The centre of a twisted group algebra  $k_\alpha G$  does not in general admit a basis as in 1.5.1, essentially because the multiplicative inverse in  $k_\alpha G$  of a group element  $x$  need not be equal to the inverse  $x^{-1}$  of  $x$  as an element in the group  $G$ .

Another measure for how far a  $k$ -algebra  $A$  is from being commutative is the *commutator space*  $[A, A]$ .

- This is the  $k$ -submodule of  $A$  generated by all *additive commutators*  $[a, b] = ab - ba$ , where  $a, b \in A$ .
- We have  $[A, A] = \{0\}$  if and only if  $A$  is commutative.

## Remarks A.

The centre of a twisted group algebra  $k_{\alpha}G$  does not in general admit a basis as in 1.5.1, essentially because the multiplicative inverse in  $k_{\alpha}G$  of a group element  $x$  need not be equal to the inverse  $x^{-1}$  of  $x$  as an element in the group  $G$ .

Another measure for how far a  $k$ -algebra  $A$  is from being commutative is the *commutator space*  $[A, A]$ .

- This is the  $k$ -submodule of  $A$  generated by all *additive commutators*  $[a, b] = ab - ba$ , where  $a, b \in A$ .
- We have  $[A, A] = \{0\}$  if and only if  $A$  is commutative.
- Note that for  $a, b \in A$  and  $z \in Z(A)$  we have

$$z[a, b] = z(ab - ba) = (za)b - b(za) = [za, b].$$



## Remarks A.

The centre of a twisted group algebra  $k_{\alpha}G$  does not in general admit a basis as in 1.5.1, essentially because the multiplicative inverse in  $k_{\alpha}G$  of a group element  $x$  need not be equal to the inverse  $x^{-1}$  of  $x$  as an element in the group  $G$ .

Another measure for how far a  $k$ -algebra  $A$  is from being commutative is the *commutator space*  $[A, A]$ .

- This is the  $k$ -submodule of  $A$  generated by all *additive commutators*  $[a, b] = ab - ba$ , where  $a, b \in A$ .
- We have  $[A, A] = \{0\}$  if and only if  $A$  is commutative.
- Note that for  $a, b \in A$  and  $z \in Z(A)$  we have

$$z[a, b] = z(ab - ba) = (za)b - b(za) = [za, b].$$

Therefore  $[A, A]$  is a  $Z(A)$ -submodule of  $A$ , and hence  $A/[A, A]$  is a  $Z(A)$ -module.

## Proposition 1.5.4

Commutator subspaces of group algebras admit the following description.

## Proposition 1.5.4

Commutator subspaces of group algebras admit the following description.

### Proposition 1.5.4

Let  $G$  be a finite group.

## Proposition 1.5.4

Commutator subspaces of group algebras admit the following description.

### Proposition 1.5.4

Let  $G$  be a finite group.

- (i) The commutator space  $[kG, kG]$  is spanned, as a  $k$ -module, by the set of commutators  $[x, y] = xy - yx$ , with  $x, y \in G$ .

## Proposition 1.5.4

Commutator subspaces of group algebras admit the following description.

### Proposition 1.5.4

Let  $G$  be a finite group.

- (i) The commutator space  $[kG, kG]$  is spanned, as a  $k$ -module, by the set of commutators  $[x, y] = xy - yx$ , with  $x, y \in G$ . *[not the same as  $G'$ ]*
- (ii) If  $x, x' \in G$  are conjugate, then  $x - x' \in [kG, kG]$ .

$$x' = g x g^{-1}$$

$$x - x' = x - g x g^{-1} = (x g^{-1}) g - g (x g^{-1}) = [x g^{-1}, g]$$

## Proposition 1.5.4

$$\sum_{x \in C} \alpha_x x \text{ st } \sum_{y \in C} \alpha_y y = 0 \quad \forall C$$

Commutator subspaces of group algebras admit the following description.

### Proposition 1.5.4

Let  $G$  be a finite group.

- (i) The commutator space  $[kG, kG]$  is spanned, as a  $k$ -module, by the set of commutators  $[x, y] = xy - yx$ , with  $x, y \in G$ . ✓
- (ii) If  $x, x' \in G$  are conjugate, then  $x - x' \in [kG, kG]$ .
- (iii) An element  $\sum_{x \in G} \alpha_x x$  of  $kG$  belongs to the commutator space  $[kG, kG]$  if and only if  $\sum_{x \in C} \alpha_x = 0$  for any conjugacy class  $C$  in  $G$ .

$$(\Rightarrow) z \in [kG, kG] ; \quad \underline{xy - yx} \quad \checkmark \quad yx = y(xy)y^{-1}$$

$$[kG, kG] = \text{span}([x, y])$$

$$\underbrace{\sum_{\lambda \in G} \alpha_\lambda \lambda \in \mathfrak{kg}}_{\text{at}} \quad \sum_{\lambda \in C} \alpha_\lambda = 0$$

$$\sum_{\lambda \in G} \alpha_\lambda \lambda = \sum_C \sum_{\lambda \in C} \alpha_\lambda \lambda$$

$$= \sum_C \underbrace{\sum_{\lambda \in C} \alpha_\lambda (\lambda - \lambda_C)}_{\text{at}}, \quad \lambda_C \in C$$

$$\in [\mathfrak{kg}, \mathfrak{kg}]$$

## Proposition 1.5.4

Commutator subspaces of group algebras admit the following description.

### Proposition 1.5.4

Let  $G$  be a finite group.

- (i) The commutator space  $[kG, kG]$  is spanned, as a  $k$ -module, by the set of commutators  $[x, y] = xy - yx$ , with  $x, y \in G$ .
- (ii) If  $x, x' \in G$  are conjugate, then  $x - x' \in [kG, kG]$ .
- (iii) An element  $\sum_{x \in G} \alpha_x x$  of  $kG$  belongs to the commutator space  $[kG, kG]$  if and only if  $\sum_{x \in C} \alpha_x = 0$  for any conjugacy class  $C$  in  $G$ .
- (iv) Let  $\mathcal{K}$  be the set of conjugacy classes, and for  $C \in \mathcal{K}$ , let  $x_C \in C$ . The  $k$ -module  $\sum_{C \in \mathcal{K}} k x_C$  is a complement of  $[kG, kG]$  in  $kG$ .

$$kG = [kG, kG] \oplus \left( \bigoplus_C k x_C \right)$$



## Remarks B.

For a group algebra  $kG$  of a finite group  $G$ , we have a close connection between the  $k$ -dual of the centre and the commutator subspace.

For a group algebra  $kG$  of a finite group  $G$ , we have a close connection between the  $k$ -dual of the centre and the commutator subspace.

Since the centre  $Z(kG)$  has as canonical  $k$ -basis the set of conjugacy class sums, the  $k$ -dual  $Z(kG)^* = \text{Hom}_k(Z(kG), k)$  therefore also has a canonical basis indexed by the set  $\mathcal{K}$  of conjugacy classes of  $G$ , namely the dual basis  $\{\sigma_C : C \in \mathcal{K}\}$ , where  $\sigma_C$  sends  $C$  to 1 and  $C' \neq C$  to 0, for  $C, C' \in \mathcal{K}$ .

For a group algebra  $kG$  of a finite group  $G$ , we have a close connection between the  $k$ -dual of the centre and the commutator subspace.

Since the centre  $Z(kG)$  has as canonical  $k$ -basis the set of conjugacy class sums, the  $k$ -dual  $Z(kG)^* = \text{Hom}_k(Z(kG), k)$  therefore also has a canonical basis indexed by the set  $\mathcal{K}$  of conjugacy classes of  $G$ , namely the dual basis  $\{\sigma_C : C \in \mathcal{K}\}$ , where  $\sigma_C$  sends  $C$  to 1 and  $C' \neq C$  to 0, for  $C, C' \in \mathcal{K}$ .

## Proposition 1.5.5

The dual of the centre and the commutator space of a finite group algebra are related as follows

## Proposition 1.5.5

The dual of the centre and the commutator space of a finite group algebra are related as follows

### Proposition 1.5.4

Let  $G$  be a finite group.

## Proposition 1.5.5

The dual of the centre and the commutator space of a finite group algebra are related as follows

### Proposition 1.5.4

Let  $G$  be a finite group.

- (i) We have an isomorphism of  $k$ -modules  $kG/[kG, kG] \cong Z(kG)^*$

## Proposition 1.5.5

The dual of the centre and the commutator space of a finite group algebra are related as follows

### Proposition 1.5.4

Let  $G$  be a finite group.

- (i) We have an isomorphism of  $k$ -modules  $kG/[kG, kG] \cong Z(kG)^*$  sending  $x + [kG, kG]$  to the unique element  $\sigma \in Z(kG)^*$  which maps the conjugacy class sum of  $x^{-1}$  to 1 and any other conjugacy class sum to zero.

$$x + [kG, kG] \mapsto \sigma \begin{cases} C_{x^{-1}} \mapsto 1 \\ \text{otherwise} \mapsto 0 \end{cases}$$

## Proposition 1.5.5

The dual of the centre and the commutator space of a finite group algebra are related as follows

### Proposition 1.5.4

Let  $G$  be a finite group.

- (i) We have an isomorphism of  $k$ -modules  $kG/[kG, kG] \cong Z(kG)^*$  sending  $x + [kG, kG]$  to the unique element  $\sigma \in Z(kG)^*$  which maps the conjugacy class sum of  $x^{-1}$  to 1 and any other conjugacy class sum to zero.

In particular, the  $k$ -module  $kG/[kG, kG]$  is free of rank equal to the number of conjugacy classes of  $G$



## Proposition 1.5.5

The dual of the centre and the commutator space of a finite group algebra are related as follows

### Proposition 1.5.4

Let  $G$  be a finite group.

- (i) We have an isomorphism of  $k$ -modules  $kG/[kG, kG] \cong Z(kG)^*$  sending  $x + [kG, kG]$  to the unique element  $\sigma \in Z(kG)^*$  which maps the conjugacy class sum of  $x^{-1}$  to 1 and any other conjugacy class sum to zero.

In particular, the  $k$ -module  $kG/[kG, kG]$  is free of rank equal to the number of conjugacy classes of  $G$  and the image in  $kG/[kG, kG]$  of a system of representatives of the conjugacy classes of  $G$  is a  $k$ -basis of  $kG/[kG, kG]$ .

## Proposition 1.5.5

The dual of the centre and the commutator space of a finite group algebra are related as follows

### Proposition 1.5.4

Let  $G$  be a finite group.

- (i) We have an isomorphism of  $k$ -modules  $kG/[kG, kG] \cong Z(kG)^*$  sending  $x + [kG, kG]$  to the unique element  $\sigma \in Z(kG)^*$  which maps the conjugacy class sum of  $x^{-1}$  to 1 and any other conjugacy class sum to zero.

In particular, the  $k$ -module  $kG/[kG, kG]$  is free of rank equal to the number of conjugacy classes of  $G$  and the image in  $kG/[kG, kG]$  of a system of representatives of the conjugacy classes of  $G$  is a  $k$ -basis of  $kG/[kG, kG]$ .

- (ii) We have an isomorphism of  $k$ -modules  $(kG/[kG, kG])^* \cong Z(kG)$

## Proposition 1.5.5

The dual of the centre and the commutator space of a finite group algebra are related as follows

### Proposition 1.5.4

Let  $G$  be a finite group.

- (i) We have an isomorphism of  $k$ -modules  $kG/[kG, kG] \cong Z(kG)^*$  sending  $x + [kG, kG]$  to the unique element  $\sigma \in Z(kG)^*$  which maps the conjugacy class sum of  $x^{-1}$  to 1 and any other conjugacy class sum to zero.

In particular, the  $k$ -module  $kG/[kG, kG]$  is free of rank equal to the number of conjugacy classes of  $G$  and the image in  $kG/[kG, kG]$  of a system of representatives of the conjugacy classes of  $G$  is a  $k$ -basis of  $kG/[kG, kG]$ .

- (ii) We have an isomorphism of  $k$ -modules  $(kG/[kG, kG])^* \cong Z(kG)$  sending

$$\phi \mapsto \sum_{x \in G} \phi(x^{-1})x,$$

## Proposition 1.5.5

The dual of the centre and the commutator space of a finite group algebra are related as follows

### Proposition 1.5.4

Let  $G$  be a finite group.

- (i) We have an isomorphism of  $k$ -modules  $kG/[kG, kG] \cong Z(kG)^*$  sending  $x + [kG, kG]$  to the unique element  $\sigma \in Z(kG)^*$  which maps the conjugacy class sum of  $x^{-1}$  to 1 and any other conjugacy class sum to zero.

In particular, the  $k$ -module  $kG/[kG, kG]$  is free of rank equal to the number of conjugacy classes of  $G$  and the image in  $kG/[kG, kG]$  of a system of representatives of the conjugacy classes of  $G$  is a  $k$ -basis of  $kG/[kG, kG]$ .

- (ii) We have an isomorphism of  $k$ -modules  $(kG/[kG, kG])^* \cong Z(kG)$  sending

$$\phi \mapsto \sum_{x \in G} \phi(x^{-1})x,$$

where we regard  $\phi$  as a map  $kG \rightarrow k$  via the canonical surjection  $kG \rightarrow kG/[kG, kG]$ .

## Proposition 1.5.5, 1.5.6

### *Proof*

It follows from 1.5.4 that  $kG/[kG, kG]$  is free of rank equal to the number of conjugacy classes of  $G$ ,

## Proposition 1.5.5, 1.5.6

### *Proof*

It follows from 1.5.4 that  $kG/[kG, kG]$  is free of rank equal to the number of conjugacy classes of  $G$ , and that the image in  $kG/[kG, kG]$  of a set  $\mathcal{R}$  of representatives of the conjugacy classes is a basis that is independent of the choice of  $\mathcal{R}$  because conjugate group elements have the same image in  $kG/[kG, kG]$ .

## Proposition 1.5.5, 1.5.6

### *Proof*

It follows from 1.5.4 that  $kG/[kG, kG]$  is free of rank equal to the number of conjugacy classes of  $G$ , and that the image in  $kG/[kG, kG]$  of a set  $\mathcal{R}$  of representatives of the conjugacy classes is a basis that is independent of the choice of  $\mathcal{R}$  because conjugate group elements have the same image in  $kG/[kG, kG]$ . Since both  $Z(kG)$  and its  $k$ -dual have bases indexed by the conjugacy classes of  $G$ , statement (i) follows.

## Proposition 1.5.5, 1.5.6

### *Proof*

It follows from 1.5.4 that  $kG/[kG, kG]$  is free of rank equal to the number of conjugacy classes of  $G$ , and that the image in  $kG/[kG, kG]$  of a set  $\mathcal{R}$  of representatives of the conjugacy classes is a basis that is independent of the choice of  $\mathcal{R}$  because conjugate group elements have the same image in  $kG/[kG, kG]$ . Since both  $Z(kG)$  and its  $k$ -dual have bases indexed by the conjugacy classes of  $G$ , statement (i) follows.

One can prove (ii) by observing that this is the dual map of the isomorphism in (i).



## Proposition 1.5.5, 1.5.6

### *Proof*

It follows from 1.5.4 that  $kG/[kG, kG]$  is free of rank equal to the number of conjugacy classes of  $G$ , and that the image in  $kG/[kG, kG]$  of a set  $\mathcal{R}$  of representatives of the conjugacy classes is a basis that is independent of the choice of  $\mathcal{R}$  because conjugate group elements have the same image in  $kG/[kG, kG]$ . Since both  $Z(kG)$  and its  $k$ -dual have bases indexed by the conjugacy classes of  $G$ , statement (i) follows.

One can prove (ii) by observing that this is the dual map of the isomorphism in (i).

### Proposition 1.5.6

Let  $A, B$  be  $k$ -algebras.

## Proposition 1.5.5, 1.5.6

### *Proof*

It follows from 1.5.4 that  $kG/[kG, kG]$  is free of rank equal to the number of conjugacy classes of  $G$ , and that the image in  $kG/[kG, kG]$  of a set  $\mathcal{R}$  of representatives of the conjugacy classes is a basis that is independent of the choice of  $\mathcal{R}$  because conjugate group elements have the same image in  $kG/[kG, kG]$ . Since both  $Z(kG)$  and its  $k$ -dual have bases indexed by the conjugacy classes of  $G$ , statement (i) follows.

One can prove (ii) by observing that this is the dual map of the isomorphism in (i).

### Proposition 1.5.6

Let  $A, B$  be  $k$ -algebras. We have

$$[A \otimes_k B, A \otimes_k B] = A \otimes_k [B, B] + [A, A] \otimes_k B$$

as  $k$ -submodules of  $A \otimes_k B$ .

$$\text{So, } [k(G \times H), k(G \times H)] = kG \otimes_k [kH, kH] + [kG, kG] \otimes_k kH$$

## Proposition 1.5.7

Given a positive integer  $n$ , the trace of a square matrix  $M = (m_{ij})_{1 \leq i, j \leq n}$  in  $M_n(k)$  is the sum of its diagonal elements  $\text{tr}(M) = \sum_{1 \leq i \leq n} m_{ii}$ .

## Proposition 1.5.7

Given a positive integer  $n$ , the trace of a square matrix  $M = (m_{ij})_{1 \leq i, j \leq n}$  in  $M_n(k)$  is the sum of its diagonal elements  $\text{tr}(M) = \sum_{1 \leq i \leq n} m_{ii}$ .

Taking traces defines a  $k$ -linear map  $\text{tr} : M_n(k) \rightarrow k$ . This map is surjective as it maps a matrix with exactly one diagonal entry 1 (and all other entries 0) to 1. An elementary verification shows that the trace is symmetric; that is,  $\text{tr}(MN) = \text{tr}(NM)$  for any two matrices  $M, N \in M_n(k)$ .

## Proposition 1.5.7

Given a positive integer  $n$ , the trace of a square matrix  $M = (m_{ij})_{1 \leq i, j \leq n}$  in  $M_n(k)$  is the sum of its diagonal elements  $\text{tr}(M) = \sum_{1 \leq i \leq n} m_{ii}$ .

Taking traces defines a  $k$ -linear map  $\text{tr} : M_n(k) \rightarrow k$ . This map is surjective as it maps a matrix with exactly one diagonal entry 1 (and all other entries 0) to 1. An elementary verification shows that the trace is symmetric; that is,  $\text{tr}(MN) = \text{tr}(NM)$  for any two matrices  $M, N \in M_n(k)$ .

The commutator space of a matrix algebra consists of all matrices whose trace is zero.

## Proposition 1.5.7

Given a positive integer  $n$ , the trace of a square matrix  $M = (m_{ij})_{1 \leq i, j \leq n}$  in  $M_n(k)$  is the sum of its diagonal elements  $\text{tr}(M) = \sum_{1 \leq i \leq n} m_{ii}$ .

Taking traces defines a  $k$ -linear map  $\text{tr} : M_n(k) \rightarrow k$ . This map is surjective as it maps a matrix with exactly one diagonal entry 1 (and all other entries 0) to 1. An elementary verification shows that the trace is symmetric; that is,  $\text{tr}(MN) = \text{tr}(NM)$  for any two matrices  $M, N \in M_n(k)$ .

The commutator space of a matrix algebra consists of all matrices whose trace is zero.

### Proposition 1.5.7

Let  $n$  be a positive integer. We have  $[M_n(k), M_n(k)] = \ker(\text{tr})$ , as a  $k$ -module,  $\ker(\text{tr})$  is free of rank  $n^2 - 1$ ,

## Proposition 1.5.7

Given a positive integer  $n$ , the trace of a square matrix  $M = (m_{ij})_{1 \leq i, j \leq n}$  in  $M_n(k)$  is the sum of its diagonal elements  $\text{tr}(M) = \sum_{1 \leq i \leq n} m_{ii}$ .

Taking traces defines a  $k$ -linear map  $\text{tr} : M_n(k) \rightarrow k$ . This map is surjective as it maps a matrix with exactly one diagonal entry 1 (and all other entries 0) to 1. An elementary verification shows that the trace is symmetric; that is,  $\text{tr}(MN) = \text{tr}(NM)$  for any two matrices  $M, N \in M_n(k)$ .

The commutator space of a matrix algebra consists of all matrices whose trace is zero.

### Proposition 1.5.7

Let  $n$  be a positive integer. We have  $[M_n(k), M_n(k)] = \ker(\text{tr})$ , as a  $k$ -module,  $\ker(\text{tr})$  is free of rank  $n^2 - 1$ , and the  $k$ -module  $M_n(k)/[M_n(k), M_n(k)]$  is free of rank 1.

## Proposition 1.5.7

Given a positive integer  $n$ , the trace of a square matrix  $M = (m_{ij})_{1 \leq i, j \leq n}$  in  $M_n(k)$  is the sum of its diagonal elements  $\text{tr}(M) = \sum_{1 \leq i \leq n} m_{ii}$ .

Taking traces defines a  $k$ -linear map  $\text{tr} : M_n(k) \rightarrow k$ . This map is surjective as it maps a matrix with exactly one diagonal entry 1 (and all other entries 0) to 1. An elementary verification shows that the trace is symmetric; that is,  $\text{tr}(MN) = \text{tr}(NM)$  for any two matrices  $M, N \in M_n(k)$ .

The commutator space of a matrix algebra consists of all matrices whose trace is zero.

### Proposition 1.5.7

Let  $n$  be a positive integer. We have  $[M_n(k), M_n(k)] = \ker(\text{tr})$ , as a  $k$ -module,  $\ker(\text{tr})$  is free of rank  $n^2 - 1$ , and the  $k$ -module  $M_n(k)/[M_n(k), M_n(k)]$  is free of rank 1. Moreover,  $M_n(k)$  is equal to the  $k$ -linear span of the group  $\text{GL}_n(k) = M_n(k)^\times$  of invertible elements in  $M_n(k)$ .



## Proposition 1.5.7

### *Proof*

Since  $\text{tr}(MN) = \text{tr}(NM)$  it follows that for any  $M, N \in M_n(k)$  we have  $[M_n(k), M_n(k)] \subseteq \ker(\text{tr})$ .

## Proposition 1.5.7

### *Proof*

Since  $\text{tr}(MN) = \text{tr}(NM)$  it follows that for any  $M, N \in M_n(k)$  we have  $[M_n(k), M_n(k)] \subseteq \ker(\text{tr})$ .

For  $1 \leq i, j \leq n$  denote by  $E_{ij}$  the matrix with entry  $(i, j)$  equal to 1 and all other entries zero. Elementary verifications show that  $E_{ij}E_{rs} = E_{is}$  if  $j = r$  and zero otherwise.

## Proposition 1.5.7

### *Proof*

Since  $\text{tr}(MN) = \text{tr}(NM)$  it follows that for any  $M, N \in M_n(k)$  we have  $[M_n(k), M_n(k)] \subseteq \ker(\text{tr})$ .

For  $1 \leq i, j \leq n$  denote by  $E_{ij}$  the matrix with entry  $(i, j)$  equal to 1 and all other entries zero. Elementary verifications show that  $E_{ij}E_{rs} = E_{is}$  if  $j = r$  and zero otherwise.

If  $i \neq j$ , then  $\text{tr}(E_{ij}) = 0$  and  $E_{ij} = [E_{ii}, E_{ij}] \in [M_n(k), M_n(k)]$ .

## Proposition 1.5.7

### *Proof*

Since  $\text{tr}(MN) = \text{tr}(NM)$  it follows that for any  $M, N \in M_n(k)$  we have  $[M_n(k), M_n(k)] \subseteq \ker(\text{tr})$ .

For  $1 \leq i, j \leq n$  denote by  $E_{ij}$  the matrix with entry  $(i, j)$  equal to 1 and all other entries zero. Elementary verifications show that  $E_{ij}E_{rs} = E_{is}$  if  $j = r$  and zero otherwise.

If  $i \neq j$ , then  $\text{tr}(E_{ij}) = 0$  and  $E_{ij} = [E_{ii}, E_{ij}] \in [M_n(k), M_n(k)]$ .

For  $2 \leq j \leq n$  denote by  $D_j$  the matrix with entry  $(1, 1)$  equal to 1, entry  $(j, j)$  equal to  $-1$  and zero everywhere else; that is,  $D_j = E_{11} - E_{jj}$ . Again  $\text{tr}(D_j) = 0$  and  $D_j = [E_{1j}, E_{j1}] \in [M_n(k), M_n(k)]$ .

## Proposition 1.5.7

### *Proof*

Since  $\text{tr}(MN) = \text{tr}(NM)$  it follows that for any  $M, N \in M_n(k)$  we have  $[M_n(k), M_n(k)] \subseteq \ker(\text{tr})$ .

For  $1 \leq i, j \leq n$  denote by  $E_{ij}$  the matrix with entry  $(i, j)$  equal to 1 and all other entries zero. Elementary verifications show that  $E_{ij}E_{rs} = E_{is}$  if  $j = r$  and zero otherwise.

If  $i \neq j$ , then  $\text{tr}(E_{ij}) = 0$  and  $E_{ij} = [E_{ii}, E_{ij}] \in [M_n(k), M_n(k)]$ .

For  $2 \leq j \leq n$  denote by  $D_j$  the matrix with entry  $(1, 1)$  equal to 1, entry  $(j, j)$  equal to  $-1$  and zero everywhere else; that is,  $D_j = E_{11} - E_{jj}$ . Again  $\text{tr}(D_j) = 0$  and  $D_j = [E_{1j}, E_{j1}] \in [M_n(k), M_n(k)]$ .

The set of matrices  $E_{ij}$  with  $i \neq j$  and  $D_j$  with  $2 \leq j \leq n$  is linearly independent, contained in  $\ker(\text{tr})$  and has  $n^2 - 1$  elements.

## Proposition 1.5.7

$$M_n(k) = \ker(\text{tr}) \oplus kE_n$$

### *Proof*

Since  $\text{tr}(MN) = \text{tr}(NM)$  it follows that for any  $M, N \in M_n(k)$  we have  $[M_n(k), M_n(k)] \subseteq \ker(\text{tr})$ .

For  $1 \leq i, j \leq n$  denote by  $E_{ij}$  the matrix with entry  $(i, j)$  equal to 1 and all other entries zero. Elementary verifications show that  $E_{ij}E_{rs} = E_{is}$  if  $j = r$  and zero otherwise.

If  $i \neq j$ , then  $\text{tr}(E_{ij}) = 0$  and  $E_{ij} = [E_{ii}, E_{ij}] \in [M_n(k), M_n(k)]$ .

For  $2 \leq j \leq n$  denote by  $D_j$  the matrix with entry  $(1, 1)$  equal to 1, entry  $(j, j)$  equal to  $-1$  and zero everywhere else; that is,  $D_j = E_{11} - E_{jj}$ . Again  $\text{tr}(D_j) = 0$  and  $D_j = [E_{1j}, E_{j1}] \in [M_n(k), M_n(k)]$ .

The set of matrices  $E_{ij}$  with  $i \neq j$  and  $D_j$  with  $2 \leq j \leq n$  is linearly independent, contained in  $\ker(\text{tr})$  and has  $n^2 - 1$  elements.

This set is a  $k$ -basis of  $\ker(\text{tr})$  because together with  $E_{11}$  it is a  $k$ -basis of  $M_n(k)$ . This shows that  $\ker(\text{tr}) \subseteq [M_n(k), M_n(k)]$ , whence the first equality as stated.

## Proposition 1.5.7

### *Proof*

Since  $\text{tr}(MN) = \text{tr}(NM)$  it follows that for any  $M, N \in M_n(k)$  we have  $[M_n(k), M_n(k)] \subseteq \ker(\text{tr})$ .

For  $1 \leq i, j \leq n$  denote by  $E_{ij}$  the matrix with entry  $(i, j)$  equal to 1 and all other entries zero. Elementary verifications show that  $E_{ij}E_{rs} = E_{is}$  if  $j = r$  and zero otherwise.

If  $i \neq j$ , then  $\text{tr}(E_{ij}) = 0$  and  $E_{ij} = [E_{ii}, E_{ij}] \in [M_n(k), M_n(k)]$ .

For  $2 \leq j \leq n$  denote by  $D_j$  the matrix with entry  $(1, 1)$  equal to 1, entry  $(j, j)$  equal to  $-1$  and zero everywhere else; that is,  $D_j = E_{11} - E_{jj}$ . Again  $\text{tr}(D_j) = 0$  and  $D_j = [E_{1j}, E_{j1}] \in [M_n(k), M_n(k)]$ .

The set of matrices  $E_{ij}$  with  $i \neq j$  and  $D_j$  with  $2 \leq j \leq n$  is linearly independent, contained in  $\ker(\text{tr})$  and has  $n^2 - 1$  elements.

This set is a  $k$ -basis of  $\ker(\text{tr})$  because together with  $E_{11}$  it is a  $k$ -basis of  $M_n(k)$ . This shows that  $\ker(\text{tr}) \subseteq [M_n(k), M_n(k)]$ , whence the first equality as stated.

## Proposition 1.5.7

*(contd.)*

The last statement is trivial if  $n = 1$ . Assume that  $n \geq 2$ . It suffices to show that the matrices  $E_{ij}$  are in the  $k$ -span of  $GL_n(k)$ .



## Proposition 1.5.7

*(contd.)*

The last statement is trivial if  $n = 1$ . Assume that  $n \geq 2$ . It suffices to show that the matrices  $E_{ij}$  are in the  $k$ -span of  $GL_n(k)$ .

If  $i \neq j$ , then  $\text{Id} + E_{ij}$  is invertible, hence  $E_{ij} = (\text{Id} + E_{ij}) - \text{Id}$  is in the span of  $GL_n(k)$ .

## Proposition 1.5.7

*(contd.)*

The last statement is trivial if  $n = 1$ . Assume that  $n \geq 2$ . It suffices to show that the matrices  $E_{ij}$  are in the  $k$ -span of  $GL_n(k)$ .

If  $i \neq j$ , then  $\text{Id} + E_{ij}$  is invertible, hence  $E_{ij} = (\text{Id} + E_{ij}) - \text{Id}$  is in the span of  $GL_n(k)$ .

The matrix  $E_{ii}$  is conjugate to  $E_{11}$  via a permutation matrix, so it suffices to show that  $E_{11}$  is in the span of  $GL_n(k)$ .

## Proposition 1.5.7

(contd.)

The last statement is trivial if  $n = 1$ . Assume that  $n \geq 2$ . It suffices to show that the matrices  $E_{ij}$  are in the  $k$ -span of  $GL_n(k)$ .

If  $i \neq j$ , then  $\text{Id} + E_{ij}$  is invertible, hence  $E_{ij} = (\text{Id} + E_{ij}) - \text{Id}$  is in the span of  $GL_n(k)$ .

The matrix  $E_{ii}$  is conjugate to  $E_{11}$  via a permutation matrix, so it suffices to show that  $E_{11}$  is in the span of  $GL_n(k)$ .

Let  $J$  be the matrix with antidiagonal entries 1 and 0 elsewhere. Then  $J$  and  $J + E_{11}$  are invertible, hence  $E_{11} = (J + E_{11}) - J$  is in the span of  $GL_n(k)$ , completing the proof.

$$J = \begin{pmatrix} & & 1 \\ & & \\ 1 & & \end{pmatrix} \quad (-1)^{n+1}$$

### Remarks C., Definition 1.5.8

A  $k$ -algebra homomorphism  $\alpha : A \rightarrow B$  need not send  $Z(A)$  to  $Z(B)$ .

$\alpha: K G_2 \hookrightarrow K S_3, e \mapsto e, (12) \mapsto (12)$   
 $\alpha|_{Z(K G_2)}: K e + K(12) \rightarrow K e + K(12) + K(13) + K(23) + K(123) + K(132)$   
 $Z(K G_2) \not\rightarrow Z(K S_3)$

A  $k$ -algebra homomorphism  $\alpha : A \rightarrow B$  need not send  $Z(A)$  to  $Z(B)$ . But  $\alpha$  sends any commutator  $[a, b]$  in  $[A, A]$  to the commutator  $[\alpha(a), \alpha(b)]$  in  $[B, B]$  and hence induces a  $k$ -linear map  $A/[A, A] \rightarrow B/[B, B]$ . Thus the quotient  $A/[A, A]$  has better functoriality properties than the centre  $Z(A)$ .

## Remarks C., Definition 1.5.8

A  $k$ -algebra homomorphism  $\alpha : A \rightarrow B$  need not send  $Z(A)$  to  $Z(B)$ . But  $\alpha$  sends any commutator  $[a, b]$  in  $[A, A]$  to the commutator  $[\alpha(a), \alpha(b)]$  in  $[B, B]$  and hence induces a  $k$ -linear map  $A/[A, A] \rightarrow B/[B, B]$ . Thus the quotient  $A/[A, A]$  has better functoriality properties than the centre  $Z(A)$ .

If  $A, B$  are group algebras then by taking  $k$ -duals one gets from 1.5.5 at least a  $k$ -linear (but not necessarily multiplicative) map  $Z(B) \rightarrow Z(A)$ , so taking centres becomes contravariantly functorial.

A  $k$ -algebra homomorphism  $\alpha : A \rightarrow B$  need not send  $Z(A)$  to  $Z(B)$ . But  $\alpha$  sends any commutator  $[a, b]$  in  $[A, A]$  to the commutator  $[\alpha(a), \alpha(b)]$  in  $[B, B]$  and hence induces a  $k$ -linear map  $A/[A, A] \rightarrow B/[B, B]$ . Thus the quotient  $A/[A, A]$  has better functoriality properties than the centre  $Z(A)$ .

If  $A, B$  are group algebras then by taking  $k$ -duals one gets from 1.5.5 at least a  $k$ -linear (but not necessarily multiplicative) map  $Z(B) \rightarrow Z(A)$ , so taking centres becomes contravariantly functorial.

### Definition 1.5.8

Let  $A$  be a  $k$ -algebra, and let  $M$  be an  $(A, A)$ -bimodule.

## Remarks C., Definition 1.5.8

A  $k$ -algebra homomorphism  $\alpha : A \rightarrow B$  need not send  $Z(A)$  to  $Z(B)$ . But  $\alpha$  sends any commutator  $[a, b]$  in  $[A, A]$  to the commutator  $[\alpha(a), \alpha(b)]$  in  $[B, B]$  and hence induces a  $k$ -linear map  $A/[A, A] \rightarrow B/[B, B]$ . Thus the quotient  $A/[A, A]$  has better functoriality properties than the centre  $Z(A)$ .

If  $A, B$  are group algebras then by taking  $k$ -duals one gets from 1.5.5 at least a  $k$ -linear (but not necessarily multiplicative) map  $Z(B) \rightarrow Z(A)$ , so taking centres becomes contravariantly functorial.

### Definition 1.5.8

Let  $A$  be a  $k$ -algebra, and let  $M$  be an  $(A, A)$ -bimodule. We set

$$M^A = \{m \in M : am = ma \text{ for all } a \in A\}.$$



## Remarks C., Definition 1.5.8

A  $k$ -algebra homomorphism  $\alpha : A \rightarrow B$  need not send  $Z(A)$  to  $Z(B)$ . But  $\alpha$  sends any commutator  $[a, b]$  in  $[A, A]$  to the commutator  $[\alpha(a), \alpha(b)]$  in  $[B, B]$  and hence induces a  $k$ -linear map  $A/[A, A] \rightarrow B/[B, B]$ . Thus the quotient  $A/[A, A]$  has better functoriality properties than the centre  $Z(A)$ .

If  $A, B$  are group algebras then by taking  $k$ -duals one gets from 1.5.5 at least a  $k$ -linear (but not necessarily multiplicative) map  $Z(B) \rightarrow Z(A)$ , so taking centres becomes contravariantly functorial.

### Definition 1.5.8

Let  $A$  be a  $k$ -algebra, and let  $M$  be an  $(A, A)$ -bimodule. We set

$$M^A = \{m \in M : am = ma \text{ for all } a \in A\}.$$

We denote by  $[A, M]$  the  $k$ -submodule of  $M$  generated by the set of elements of the form  $am - ma$  in  $M$ , where  $a \in A$  and  $m \in M$ .

$$M/[A, M]$$

$$M_A$$

## Remark 1.5.9

### Remark 1.5.9

Let  $A$  be a  $k$ -algebra and let  $M, N$  be  $(A, A)$ -bimodules. Some of the earlier properties of centres and commutator spaces extend to fixed points and commutator spaces in bimodules.

## Remark 1.5.9

### Remark 1.5.9

Let  $A$  be a  $k$ -algebra and let  $M, N$  be  $(A, A)$ -bimodules. Some of the earlier properties of centres and commutator spaces extend to fixed points and commutator spaces in bimodules.

The sets  $M^A$  and  $[A, M]$  are both left and right  $Z(A)$ -submodules of  $M$ , and we have  $A^A = Z(A)$ .

### Remark 1.5.9

Let  $A$  be a  $k$ -algebra and let  $M, N$  be  $(A, A)$ -bimodules. Some of the earlier properties of centres and commutator spaces extend to fixed points and commutator spaces in bimodules.

The sets  $M^A$  and  $[A, M]$  are both left and right  $Z(A)$ -submodules of  $M$ , and we have  $A^A = Z(A)$ .

A bimodule homomorphism  $\theta : M \rightarrow N$  restricts to maps  $M^A \rightarrow N^A$  and  $[A, M] \rightarrow [A, N]$ .

### Remark 1.5.9

Let  $A$  be a  $k$ -algebra and let  $M, N$  be  $(A, A)$ -bimodules. Some of the earlier properties of centres and commutator spaces extend to fixed points and commutator spaces in bimodules.

The sets  $M^A$  and  $[A, M]$  are both left and right  $Z(A)$ -submodules of  $M$ , and we have  $A^A = Z(A)$ .

A bimodule homomorphism  $\theta : M \rightarrow N$  restricts to maps  $M^A \rightarrow N^A$  and  $[A, M] \rightarrow [A, N]$ .

If  $\theta$  is injective, then both of these restrictions are injective as well.

### Remark 1.5.9

Let  $A$  be a  $k$ -algebra and let  $M, N$  be  $(A, A)$ -bimodules. Some of the earlier properties of centres and commutator spaces extend to fixed points and commutator spaces in bimodules.

The sets  $M^A$  and  $[A, M]$  are both left and right  $Z(A)$ -submodules of  $M$ , and we have  $A^A = Z(A)$ .

A bimodule homomorphism  $\theta : M \rightarrow N$  restricts to maps  $M^A \rightarrow N^A$  and  $[A, M] \rightarrow [A, N]$ .

If  $\theta$  is injective, then both of these restrictions are injective as well. If  $\theta$  is surjective, then the induced map  $[A, M] \rightarrow [A, N]$  is surjective,

### Remark 1.5.9

Let  $A$  be a  $k$ -algebra and let  $M, N$  be  $(A, A)$ -bimodules. Some of the earlier properties of centres and commutator spaces extend to fixed points and commutator spaces in bimodules.

The sets  $M^A$  and  $[A, M]$  are both left and right  $Z(A)$ -submodules of  $M$ , and we have  $A^A = Z(A)$ .

A bimodule homomorphism  $\theta : M \rightarrow N$  restricts to maps  $M^A \rightarrow N^A$  and  $[A, M] \rightarrow [A, N]$ .

If  $\theta$  is injective, then both of these restrictions are injective as well. If  $\theta$  is surjective, then the induced map  $[A, M] \rightarrow [A, N]$  is surjective, but the induced map  $M^A \rightarrow N^A$  need not be surjective.

### Remark 1.5.9

Let  $A$  be a  $k$ -algebra and let  $M, N$  be  $(A, A)$ -bimodules. Some of the earlier properties of centres and commutator spaces extend to fixed points and commutator spaces in bimodules.

The sets  $M^A$  and  $[A, M]$  are both left and right  $Z(A)$ -submodules of  $M$ , and we have  $A^A = Z(A)$ .

A bimodule homomorphism  $\theta : M \rightarrow N$  restricts to maps  $M^A \rightarrow N^A$  and  $[A, M] \rightarrow [A, N]$ .

If  $\theta$  is injective, then both of these restrictions are injective as well. If  $\theta$  is surjective, then the induced map  $[A, M] \rightarrow [A, N]$  is surjective, but the induced map  $M^A \rightarrow N^A$  need not be surjective.

That is, taking  $A$ -fixed points on bimodules, viewed as a functor from the category of  $(A, A)$ -bimodules to the category of  $k$ -modules, is left exact but not right exact.



## Remark 1.5.10

### Remark 1.5.10

Let  $A$  be a  $k$ -algebra and  $U$  an  $A$ -module.

### Remark 1.5.10

Let  $A$  be a  $k$ -algebra and  $U$  an  $A$ -module. Then  $\text{End}_k(U)$  is an  $(A, A)$ -bimodule through composing and precomposing linear endomorphisms of  $U$  by the action of elements in  $A$  on  $U$ .

### Remark 1.5.10

Let  $A$  be a  $k$ -algebra and  $U$  an  $A$ -module. Then  $\text{End}_k(U)$  is an  $(A, A)$ -bimodule through composing and precomposing linear endomorphisms of  $U$  by the action of elements in  $A$  on  $U$ .

More precisely, for  $\phi \in \text{End}_k(U)$  and  $a, b \in A$  the  $k$ -endomorphism  $a \cdot \phi \cdot b$  of  $U$  is defined by  $(a \cdot \phi \cdot b)(u) = a\phi(bu)$  for all  $u \in U$ .

### Remark 1.5.10

Let  $A$  be a  $k$ -algebra and  $U$  an  $A$ -module. Then  $\text{End}_k(U)$  is an  $(A, A)$ -bimodule through composing and precomposing linear endomorphisms of  $U$  by the action of elements in  $A$  on  $U$ .

More precisely, for  $\phi \in \text{End}_k(U)$  and  $a, b \in A$  the  $k$ -endomorphism  $a \cdot \phi \cdot b$  of  $U$  is defined by  $(a \cdot \phi \cdot b)(u) = a\phi(bu)$  for all  $u \in U$ .

We have

$$(\text{End}_k(U))^A = \text{End}_A(U);$$

indeed, the equality  $a \cdot \phi = \phi \cdot a$  is equivalent to  $a\phi(u) = \phi(au)$  for all  $u \in U$ .

# Proposition 1.5.11

## Proposition 1.5.11

Let  $A$  be a  $k$ -algebra and  $M$  an  $(A, A)$ -bimodule.

## Proposition 1.5.11

### Proposition 1.5.11

Let  $A$  be a  $k$ -algebra and  $M$  an  $(A, A)$ -bimodule. The map sending an  $(A, A)$ -bimodule homomorphism  $\phi : A \rightarrow M$  to its evaluation  $\phi(1) \in M$  induces an isomorphism

# Proposition 1.5.11

## Proposition 1.5.11

Let  $A$  be a  $k$ -algebra and  $M$  an  $(A, A)$ -bimodule. The map sending an  $(A, A)$ -bimodule homomorphism  $\phi : A \rightarrow M$  to its evaluation  $\phi(1) \in M$  induces an isomorphism of  $Z(A)$ -modules  $\text{Hom}_{A \otimes_k A^{\text{op}}}(A, M) \cong M^A$ .

$$(k, M) = H^0$$

## Proposition 1.5.11

### Proposition 1.5.11

Let  $A$  be a  $k$ -algebra and  $M$  an  $(A, A)$ -bimodule. The map sending an  $(A, A)$ -bimodule homomorphism  $\phi : A \rightarrow M$  to its evaluation  $\phi(1) \in M$  induces an isomorphism of  $Z(A)$ -modules  $\text{Hom}_{A \otimes_k A^{\text{op}}}(A, M) \cong M^A$ . The inverse of this isomorphism sends an element  $m \in M^A$  to the homomorphism  $\psi : A \rightarrow M$  defined by  $\psi(a) = am$  for all  $a \in A$ .



## Proposition 1.5.11

### Proposition 1.5.11

Let  $A$  be a  $k$ -algebra and  $M$  an  $(A, A)$ -bimodule. The map sending an  $(A, A)$ -bimodule homomorphism  $\phi : A \rightarrow M$  to its evaluation  $\phi(1) \in M$  induces an isomorphism of  $Z(A)$ -modules  $\text{Hom}_{A \otimes_k A^{\text{op}}}(A, M) \cong M^A$ . The inverse of this isomorphism sends an element  $m \in M^A$  to the homomorphism  $\psi : A \rightarrow M$  defined by  $\psi(a) = am$  for all  $a \in A$ .

### *Proof*

For  $\phi : A \rightarrow M$  an  $(A, A)$ -bimodule homomorphism, we have  $a\phi(1) = \phi(a) = \phi(1)a$  for all  $a \in A$ ,

## Proposition 1.5.11

### Proposition 1.5.11

Let  $A$  be a  $k$ -algebra and  $M$  an  $(A, A)$ -bimodule. The map sending an  $(A, A)$ -bimodule homomorphism  $\phi : A \rightarrow M$  to its evaluation  $\phi(1) \in M$  induces an isomorphism of  $Z(A)$ -modules  $\text{Hom}_{A \otimes_k A^{\text{op}}}(A, M) \cong M^A$ . The inverse of this isomorphism sends an element  $m \in M^A$  to the homomorphism  $\psi : A \rightarrow M$  defined by  $\psi(a) = am$  for all  $a \in A$ .

### *Proof*

For  $\phi : A \rightarrow M$  an  $(A, A)$ -bimodule homomorphism, we have  $a\phi(1) = \phi(a) = \phi(1)a$  for all  $a \in A$ , which shows that  $\phi(1) \in M^A$ , so the map in the statement is well-defined,

## Proposition 1.5.11

### Proposition 1.5.11

Let  $A$  be a  $k$ -algebra and  $M$  an  $(A, A)$ -bimodule. The map sending an  $(A, A)$ -bimodule homomorphism  $\phi : A \rightarrow M$  to its evaluation  $\phi(1) \in M$  induces an isomorphism of  $Z(A)$ -modules  $\text{Hom}_{A \otimes_k A^{\text{op}}}(A, M) \cong M^A$ . The inverse of this isomorphism sends an element  $m \in M^A$  to the homomorphism  $\psi : A \rightarrow M$  defined by  $\psi(a) = am$  for all  $a \in A$ .

### *Proof*

For  $\phi : A \rightarrow M$  an  $(A, A)$ -bimodule homomorphism, we have  $a\phi(1) = \phi(a) = \phi(1)a$  for all  $a \in A$ , which shows that  $\phi(1) \in M^A$ , so the map in the statement is well-defined, and clearly this map is a  $Z(A)$ -module homomorphism.

## Proposition 1.5.11

### Proposition 1.5.11

Let  $A$  be a  $k$ -algebra and  $M$  an  $(A, A)$ -bimodule. The map sending an  $(A, A)$ -bimodule homomorphism  $\phi : A \rightarrow M$  to its evaluation  $\phi(1) \in M$  induces an isomorphism of  $Z(A)$ -modules  $\text{Hom}_{A \otimes_k A^{\text{op}}}(A, M) \cong M^A$ . The inverse of this isomorphism sends an element  $m \in M^A$  to the homomorphism  $\psi : A \rightarrow M$  defined by  $\psi(a) = am$  for all  $a \in A$ .

### *Proof*

For  $\phi : A \rightarrow M$  an  $(A, A)$ -bimodule homomorphism, we have  $a\phi(1) = \phi(a) = \phi(1)a$  for all  $a \in A$ , which shows that  $\phi(1) \in M^A$ , so the map in the statement is well-defined, and clearly this map is a  $Z(A)$ -module homomorphism.

The map  $\psi$ , as defined in the statement, is indeed an  $(A, A)$ -bimodule homomorphism, since we have  $am = ma$  for all  $a \in A$  as  $m \in M^A$ .

## Proposition 1.5.11

### Proposition 1.5.11

Let  $A$  be a  $k$ -algebra and  $M$  an  $(A, A)$ -bimodule. The map sending an  $(A, A)$ -bimodule homomorphism  $\phi : A \rightarrow M$  to its evaluation  $\phi(1) \in M$  induces an isomorphism of  $Z(A)$ -modules  $\text{Hom}_{A \otimes_k A^{\text{op}}}(A, M) \cong M^A$ . The inverse of this isomorphism sends an element  $m \in M^A$  to the homomorphism  $\psi : A \rightarrow M$  defined by  $\psi(a) = am$  for all  $a \in A$ .

### *Proof*

For  $\phi : A \rightarrow M$  an  $(A, A)$ -bimodule homomorphism, we have  $a\phi(1) = \phi(a) = \phi(1)a$  for all  $a \in A$ , which shows that  $\phi(1) \in M^A$ , so the map in the statement is well-defined, and clearly this map is a  $Z(A)$ -module homomorphism.

The map  $\psi$ , as defined in the statement, is indeed an  $(A, A)$ -bimodule homomorphism, since we have  $am = ma$  for all  $a \in A$  as  $m \in M^A$ . A trivial verification shows that the two maps in the statement are inverse to each other.

## Proposition 1.5.12

### Proposition 1.5.12

Let  $A$  be a  $k$ -algebra.

## Proposition 1.5.12

### Proposition 1.5.12

Let  $A$  be a  $k$ -algebra. The map sending an  $(A, A)$ -bimodule endomorphism  $\phi$  of  $A$  to  $\phi(1)$  is a  $k$ -algebra isomorphism  $\text{End}_{A \otimes_k A^{\text{op}}}(A) \cong Z(A)$ .

## Proposition 1.5.12

### Proposition 1.5.12

Let  $A$  be a  $k$ -algebra. The map sending an  $(A, A)$ -bimodule endomorphism  $\phi$  of  $A$  to  $\phi(1)$  is a  $k$ -algebra isomorphism  $\text{End}_{A \otimes_k A^{\text{op}}}(A) \cong Z(A)$ . Its inverse sends  $z \in Z(A)$  to the endomorphism  $\psi$  of  $A$  defined by  $\psi(a) = az$  for all  $a \in A$ .



# Proposition 1.5.12

## Proposition 1.5.12

Let  $A$  be a  $k$ -algebra. The map sending an  $(A, A)$ -bimodule endomorphism  $\phi$  of  $A$  to  $\phi(1)$  is a  $k$ -algebra isomorphism  $\text{End}_{A \otimes_k A^{\text{op}}}(A) \cong Z(A)$ . Its inverse sends  $z \in Z(A)$  to the endomorphism  $\psi$  of  $A$  defined by  $\psi(a) = az$  for all  $a \in A$ .

Corollary 1.5.12 implies that idempotents in  $Z(A)$  correspond bijectively to projections of  $A$  onto bimodule summands of  $A$ .

- [1] Linckelmann, Markus (2018). *The Block Theory of Finite Group Algebras*. Cambridge University Press.