

Knots, Links, and a Little Magic


Amethyst Price
UCSC Graduate Student
Spring 2022

Department of Mathematics
UCSC Undergraduate Colloquium by AWM




Let's Begin with Some Magic!

Case 1:



Case 2:



Amazing! Why does it work?

Amazing! Why does it work?

Tools we will need

- ◇ Some Topology
- ◇ Some (Abstract) Algebra
- ◇ Knots and Links
- ◇ Knot and Link Complements
- ◇ The Knot Group
- ◇ The Wirtinger Presentation

Definitions

Background

- ◇ **Topology** is the study of geometric properties and spatial relations unaffected by the *continuous* change of shape or size of figures.

Background

- ◇ **Topology** is the study of geometric properties and spatial relations unaffected by the *continuous* change of shape or size of figures.
- ◇ We define the **0-sphere, 1-sphere, 2-sphere, and 3-sphere** (resp.) as follows: $\mathcal{S}^0 = \{-1, 1\}$, \mathcal{S}^1 = the unit circle, \mathcal{S}^2 = the unit sphere, and $\mathcal{S}^3 = \mathcal{S}^1 \times \mathcal{S}^1 \times \mathcal{S}^1 = \{(x, y, z) | x, y, z \in \mathcal{S}^1\}$.

Background

- ◇ **Topology** is the study of geometric properties and spatial relations unaffected by the *continuous* change of shape or size of figures.
- ◇ We define the **0-sphere**, **1-sphere**, **2-sphere**, and **3-sphere** (resp.) as follows: $\mathcal{S}^0 = \{-1, 1\}$, \mathcal{S}^1 = the unit circle, \mathcal{S}^2 = the unit sphere, and $\mathcal{S}^3 = \mathcal{S}^1 \times \mathcal{S}^1 \times \mathcal{S}^1 = \{(x, y, z) | x, y, z \in \mathcal{S}^1\}$.
- ◇ A **homeomorphism** is a continuous bijective map $f : X \rightarrow Y$ between topological spaces X and Y . Two spaces with a homeomorphism between them are called homeomorphic, and from a topological viewpoint they are "the same".

Background

- ◇ **Topology** is the study of geometric properties and spatial relations unaffected by the *continuous* change of shape or size of figures.
- ◇ We define the **0-sphere**, **1-sphere**, **2-sphere**, and **3-sphere** (resp.) as follows: $\mathcal{S}^0 = \{-1, 1\}$, \mathcal{S}^1 = the unit circle, \mathcal{S}^2 = the unit sphere, and $\mathcal{S}^3 = \mathcal{S}^1 \times \mathcal{S}^1 \times \mathcal{S}^1 = \{(x, y, z) | x, y, z \in \mathcal{S}^1\}$.
- ◇ A **homeomorphism** is a continuous bijective map $f : X \rightarrow Y$ between topological spaces X and Y . Two spaces with a homeomorphism between them are called homeomorphic, and from a topological viewpoint they are "the same".
- ◇ Two continuous functions from one topological space to another are called **homotopic** if one can be "continuously deformed" into the other. This is a weaker notion than being homeomorphic.

Background

- ◇ While a homeomorphism takes one object and continuously distorts it into another, an **ambient isotopy** is a distortion of the ambient space surrounding an object which morphs the object into another.

Background

- ◇ While a homeomorphism takes one object and continuously distorts it into another, an **ambient isotopy** is a distortion of the ambient space surrounding an object which morphs the object into another.
- ◇ This is a stronger notion than being homeomorphic.

Background

- ◇ While a homeomorphism takes one object and continuously distorts it into another, an **ambient isotopy** is a distortion of the ambient space surrounding an object which morphs the object into another.
- ◇ This is a stronger notion than being homeomorphic.
- ◇ A **(topological) embedding** is an injective continuous map

$$f : X \rightarrow Y$$

that yields a homeomorphism between X and $f(X)$

What is a knot?

What is a knot?

A subset K in \mathbb{R}^3 (or S^3) is a **knot** if it is an embedding of S^1 .

What is a knot?

A subset K in \mathbb{R}^3 (or S^3) is a **knot** if it is an embedding of S^1 .

We call K the **unknot** if $K = S^1$. In otherwords, K is a trivial embedding in a 3-dimensional space.


What is a knot?

A subset K in \mathbb{R}^3 (or S^3) is a **knot** if it is an embedding of S^1 .


We call K the **unknot** if $K = S^1$. In otherwords, K is a trivial embedding in a 3-dimensional space.

What is the difference?

The Trefoil:



A Trivial Knot:




Knot Diagrams

In $X = \mathbb{R}^3$ to draw knots, we use **regular projections** to some plane in X with corresponding *over-under crossings*.


Knot Diagrams

In $X = \mathbb{R}^3$ to draw knots, we use **regular projections** to some plane in X with corresponding *over-under crossings*.

Example



Examples



What is a link?


A subset L of a 3-dimensional space X is a **link** if it is an embedding of $\mathcal{S}^0 \cup \dots \cup \mathcal{S}^0 \cup \mathcal{S}^1 \cup \dots \cup \mathcal{S}^1$.

What is a link?

A subset L of a 3-dimensional space X is a **link** if it is an embedding of $\mathcal{S}^0 \cup \dots \cup \mathcal{S}^0 \cup \mathcal{S}^1 \cup \dots \cup \mathcal{S}^1$.

Remark: The number of components need not be finite.

Example: Antione's Necklace



What makes two knots (or links) the "same"?

What makes two knots (or links) the "same"?

Two knots (links) K and K' are **equivalent**, denoted $K \sim K'$, if there exists a homeomorphism

$$h : X \rightarrow X$$

$$K \mapsto K'$$

What makes two knots (or links) the "same"?

Two knots (links) K and K' are **equivalent**, denoted $K \sim K'$, if there exists a homeomorphism

$$h : X \rightarrow X$$

$$K \mapsto K'$$

Equivalent knots are said to have the same **knot type**.

What makes two knots (or links) the "same"?

Two knots (links) K and K' are **equivalent**, denoted $K \sim K'$, if there exists a homeomorphism

$$h : X \rightarrow X$$

$$K \mapsto K'$$


Equivalent knots are said to have the same **knot type**.

We are often more interested in **knot types modulo ambient isotopies**.

What about knots/links in 2D spaces?


What about knots/links in 2D spaces?

Example: Non-Equivalent links in \mathcal{S}^2



What about knots/links in 2D spaces?


Example: Non-Equivalent links in \mathcal{S}^2



However, all knots in \mathbb{R}^2 are trivial.

What about knots/links in 2D spaces?


Example: Non-Equivalent links in \mathcal{S}^2




However, all knots in \mathbb{R}^2 are trivial.




A Trivial Knot in \mathbb{R}^3



A Trivial Knot in \mathbb{R}^3



A Trivial Knot in \mathbb{R}^3




Reidemeister Moves

In the previous example we used what we call **Reidemeister moves** to re-draw the knot:

Reidemeister Moves


In the previous example we used what we call **Reidemeister moves** to re-draw the knot:

Type I



Reidemeister Moves


In the previous example we used what we call **Reidemeister moves** to re-draw the knot:




Reidemeister Moves

In the previous example we used what we call **Reidemeister moves** to re-draw the knot:


Type I



Type II



Type III



Torus Knots

More Background

- ◇ A (simple) **loop** in a space X is a non self-intersecting path that begins and ends at the same point. We define paths using functions.

More Background

- ◇ A (simple) **loop** in a space X is a non self-intersecting path that begins and ends at the same point. We define paths using functions.
- ◇ The loops on a space modulo homotopy equivalence give us a lot of information about the space itself.

More Background

- ◇ A (simple) **loop** in a space X is a non self-intersecting path that begins and ends at the same point. We define paths using functions.
- ◇ The loops on a space modulo homotopy equivalence give us a lot of information about the space itself.
- ◇ The group of all loops on a space X modulo homotopy equivalence is called the **fundamental group of X** , and is denoted $\pi_1(X)$.

Background

- ◇ A **group**, G , is a set of elements with an associated binary operator (*like* multiplication or addition) that follow certain axioms.

Background

- ◇ A **group**, G , is a set of elements with an associated binary operator (*like* multiplication or addition) that follow certain axioms.
- ◇ A group's elements may commute or they may not! If a group is commutative we call it **Abelian**, otherwise we call it **non-Abelian**.

Background

- ◇ A **group**, G , is a set of elements with an associated binary operator (*like* multiplication or addition) that follow certain axioms.
- ◇ A group's elements may commute or they may not! If a group is commutative we call it **Abelian**, otherwise we call it **non-Abelian**.
- ◇ For elements $x, y \in G$, there is a special element $[x, y] = xyx^{-1}y^{-1} \in G$ called the **commutator**. If G is Abelian, then $[x, y] = yxx^{-1}y^{-1} = yy^{-1} = 1$.

Background

- ◇ A **group**, G , is a set of elements with an associated binary operator (*like* multiplication or addition) that follow certain axioms.
- ◇ A group's elements may commute or they may not! If a group is commutative we call it **Abelian**, otherwise we call it **non-Abelian**.
- ◇ For elements $x, y \in G$, there is a special element $[x, y] = xyx^{-1}y^{-1} \in G$ called the **commutator**. If G is Abelian, then $[x, y] = yxx^{-1}y^{-1} = yy^{-1} = 1$.
- ◇ A **group presentation** is a way to specify a group with a set of *generators*, \mathcal{S} , and a set of *relations*, \mathcal{R} , denoted $G = \langle \mathcal{S} \mid \mathcal{R} \rangle$.

The Fundamental Group of the Torus

The Fundamental Group of the Torus

$$\pi_1(\mathbb{T}) = \mathbb{Z}^2 = \langle x, y \mid xy = yx \rangle$$

The Fundamental Group of the Torus

$$\pi_1(\mathbb{T}) = \mathbb{Z}^2 = \langle x, y \mid xy = yx \rangle$$

For a loop on the torus, ℓ , the associated element in π_1 is denoted $[\ell] = \langle a, b \rangle$ where $a, b \in \mathbb{Z}$.

The Fundamental Group of the Torus

$$\pi_1(\mathbb{T}) = \mathbb{Z}^2 = \langle x, y \mid xy = yx \rangle$$

For a loop on the torus, ℓ , the associated element in π_1 is denoted $[\ell] = \langle a, b \rangle$ where $a, b \in \mathbb{Z}$.

Examples

$$[M] = \langle 1, 0 \rangle$$



The Fundamental Group of the Torus

$$\pi_1(\mathbb{T}) = \mathbb{Z}^2 = \langle x, y \mid xy = yx \rangle$$

For a loop on the torus, ℓ , the associated element in π_1 is denoted $[\ell] = \langle a, b \rangle$ where $a, b \in \mathbb{Z}$.

Examples

$$[M] = \langle 1, 0 \rangle$$

$$[L] = \langle 0, 1 \rangle$$



The Fundamental Group of the Torus

$$\pi_1(\mathbb{T}) = \mathbb{Z}^2 = \langle x, y \mid xy = yx \rangle$$


For a loop on the torus, ℓ , the associated element in π_1 is denoted $[\ell] = \langle a, b \rangle$ where $a, b \in \mathbb{Z}$.

Examples

$$[M] = \langle 1, 0 \rangle$$

$$[L] = \langle 0, 1 \rangle$$

$$[J] = \langle 0, 0 \rangle$$



The Fundamental Group of the Torus

$$\pi_1(\mathbb{T}) = \mathbb{Z}^2 = \langle x, y \mid xy = yx \rangle$$


For a loop on the torus, ℓ , the associated element in π_1 is denoted $[\ell] = \langle a, b \rangle$ where $a, b \in \mathbb{Z}$.

Examples

$$[M] = \langle 1, 0 \rangle$$



$$[L] = \langle 0, 1 \rangle$$



$$[J] = \langle 0, 0 \rangle$$



$$\langle 3, 0 \rangle$$



The Fundamental Group of the Torus

$$\pi_1(\mathbb{T}) = \mathbb{Z}^2 = \langle x, y \mid xy = yx \rangle$$


For a loop on the torus, ℓ , the associated element in π_1 is denoted $[\ell] = \langle a, b \rangle$ where $a, b \in \mathbb{Z}$.

Examples

$$[M] = \langle 1, 0 \rangle$$



$$[L] = \langle 0, 1 \rangle$$



$$[J] = \langle 0, 0 \rangle$$



$$\langle 3, 0 \rangle$$



$$\langle 3, 1 \rangle$$



Torus Knots

Claim: There are only two knot types modulo ambient isotopy on the torus.

Torus Knots

Claim: There are only two knot types modulo ambient isotopy on the torus.

Any guesses what they are?

Torus Knots

Claim: There are only two knot types modulo ambient isotopy on the torus.

Any guesses what they are?

An important observation: Simple loops are knots!

Torus Knots

Claim: There are only two knot types modulo ambient isotopy on the torus.

Any guesses what they are?

An important observation: Simple loops are knots!

Some natural guesses:

Torus Knots

Claim: There are only two knot types modulo ambient isotopy on the torus.

Any guesses what they are?

An important observation: Simple loops are knots!

Some natural guesses:

$$\begin{aligned}
 [M] &= [\text{Meridian}] \\
 &= \langle 0, 1 \rangle
 \end{aligned}$$



Torus Knots

Claim: There are only two knot types modulo ambient isotopy on the torus.

Any guesses what they are?

An important observation: Simple loops are knots!

Some natural guesses:

$$\begin{aligned}
 [M] &= [\text{Meridian}] \\
 &= \langle 0, 1 \rangle
 \end{aligned}$$

$$\begin{aligned}
 [L] &= [\text{Longitude}] \\
 &= \langle 1, 0 \rangle
 \end{aligned}$$



Torus Knots

Claim: There are only two knot types modulo ambient isotopy on the torus.

Any guesses what they are?

An important observation: Simple loops are knots!

Some natural guesses:

$$\begin{aligned}[M] &= [\text{Meridian}] \\ &= \langle 0, 1 \rangle\end{aligned}$$

$$\begin{aligned}[L] &= [\text{Longitude}] \\ &= \langle 1, 0 \rangle\end{aligned}$$

$$\begin{aligned}[J] &= [\text{Inessential}] \\ &= \langle 0, 0 \rangle\end{aligned}$$



$$M \approx J$$

We can see that $M \approx J$ because J is a separating curve and M is not.

$$M \approx J$$

We can see that $M \approx J$ because J is a separating curve and M is not.



$$M \approx J$$

We can see that $M \approx J$ because J is a separating curve and M is not.




$$L \approx J$$

We can see that $L \approx J$ for the same reason.

$$L \approx J$$

We can see that $L \approx J$ for the same reason.



Torus Knots

Behind the Scenes: We exploit more topology, abstract algebra, and some linear algebra.

Torus Knots

Behind the Scenes: We exploit more topology, abstract algebra, and some linear algebra.

Results

- ◇ Any two knots of class $\langle 0, \pm 1 \rangle$ are ambient isotopic.

Torus Knots

Behind the Scenes: We exploit more topology, abstract algebra, and some linear algebra.

Results

- ◇ Any two knots of class $\langle 0, \pm 1 \rangle$ are ambient isotopic.
- ◇ There exists a composition of twist homeomorphisms taking L to M .
Therefore $L \sim M$.

Torus Knots

Behind the Scenes: We exploit more topology, abstract algebra, and some linear algebra.

Results

- ◇ Any two knots of class $\langle 0, \pm 1 \rangle$ are ambient isotopic.
- ◇ There exists a composition of twist homeomorphisms taking L to M . Therefore $L \sim M$.
- ◇ For all $\langle a, b \rangle \in \pi_1(\mathbb{T}^2)$, \exists a composition of twist homeomorphisms taking $\langle a, b \rangle$ to $\langle 0, d \rangle$ for $\gcd(a, b) = d$ (unless $a = 0$ or $b = 0$).

Torus Knots

Behind the Scenes: We exploit more topology, abstract algebra, and some linear algebra.

Results

- ◇ Any two knots of class $\langle 0, \pm 1 \rangle$ are ambient isotopic.
- ◇ There exists a composition of twist homeomorphisms taking L to M . Therefore $L \sim M$.
- ◇ For all $\langle a, b \rangle \in \pi_1(\mathbb{T}^2)$, \exists a composition of twist homeomorphisms taking $\langle a, b \rangle$ to $\langle 0, d \rangle$ for $\gcd(a, b) = d$ (unless $a = 0$ or $b = 0$).
- ◇ For any K such that $[K] \neq \langle 0, 0 \rangle$, \exists a homeomorphism taking K to M .

Torus Knots

Behind the Scenes: We exploit more topology, abstract algebra, and some linear algebra.

Results

- ◇ Any two knots of class $\langle 0, \pm 1 \rangle$ are ambient isotopic.
- ◇ There exists a composition of twist homeomorphisms taking L to M . Therefore $L \sim M$.
- ◇ For all $\langle a, b \rangle \in \pi_1(\mathbb{T}^2)$, \exists a composition of twist homeomorphisms taking $\langle a, b \rangle$ to $\langle 0, d \rangle$ for $\gcd(a, b) = d$ (unless $a = 0$ or $b = 0$).
- ◇ For any K such that $[K] \neq \langle 0, 0 \rangle$, \exists a homeomorphism taking K to M .
- ◇ Knots K and K' are ambient isotopic $\iff [K] = [\pm K'] \in \pi_1(\mathbb{T}^2)$.

Torus Knots

Behind the Scenes: We exploit more topology, abstract algebra, and some linear algebra.

Results

- ◇ Any two knots of class $\langle 0, \pm 1 \rangle$ are ambient isotopic.
- ◇ There exists a composition of twist homeomorphisms taking L to M . Therefore $L \sim M$.
- ◇ For all $\langle a, b \rangle \in \pi_1(\mathbb{T}^2)$, \exists a composition of twist homeomorphisms taking $\langle a, b \rangle$ to $\langle 0, d \rangle$ for $\gcd(a, b) = d$ (unless $a = 0$ or $b = 0$).
- ◇ For any K such that $[K] \neq \langle 0, 0 \rangle$, \exists a homeomorphism taking K to M .
- ◇ Knots K and K' are ambient isotopic $\iff [K] = [\pm K'] \in \pi_1(\mathbb{T}^2)$.

Therefore, there are only two knot types on \mathbb{T}^2 : the inessential, and everything else!

The Solid Torus

Let $V = \mathcal{S}^1 \times D^1 =$ a solid torus.

The Solid Torus

Let $V = \mathcal{S}^1 \times D^1 =$ a solid torus.

We still consider a meridian M and a longitude L , simple closed curves on ∂V .

The Solid Torus

Let $V = S^1 \times D^1 =$ a solid torus.

We still consider a meridian M and a longitude L , simple closed curves on ∂V .

- ◇ Any two meridians are ambient isotopic.

The Solid Torus

Let $V = S^1 \times D^1 =$ a solid torus.

We still consider a meridian M and a longitude L , simple closed curves on ∂V .

- ◇ Any two meridians are ambient isotopic.
- ◇ There are infinitely many ambient isotopies of longitudes.

The Solid Torus

Let $V = \mathcal{S}^1 \times D^1 =$ a solid torus.

We still consider a meridian M and a longitude L , simple closed curves on ∂V .

- ◇ Any two meridians are ambient isotopic.
- ◇ There are infinitely many ambient isotopies of longitudes.
- ◇ If we embedded V into \mathcal{S}^3 and take the **complement** $X = \mathcal{S}^3 \setminus V$, $\pi_1(X)$ can tell us a lot about the embedding.

The Solid Torus

The complement X may or may not also be a solid torus.

The Solid Torus

The complement X may or may not also be a solid torus.

Theorem

The complement X is a solid torus $\iff \pi_1(X) = \mathbb{Z}$


The Solid Torus

The complement X may or may not also be a solid torus.

Theorem

The complement X is a solid torus $\iff \pi_1(X) = \mathbb{Z}$

If X is not a solid torus, it is sometimes called a **cube-with-knotted-hole**



The Knot Group

The Knot Group

A thickened knot is an embedding of a solid torus. So it is enlightening to consider the fundamental group of **knot complements**, $X = \mathbb{R}^3 \setminus K$ (or $S^3 \setminus K$).

The Knot Group

A thickened knot is an embedding of a solid torus. So it is enlightening to consider the fundamental group of **knot complements**, $X = \mathbb{R}^3 \setminus K$ (or $S^3 \setminus K$).

We call $\pi_1(X)$ the **knot group** of K .

The Knot Group

A thickened knot is an embedding of a solid torus. So it is enlightening to consider the fundamental group of **knot complements**, $X = \mathbb{R}^3 \setminus K$ (or $S^3 \setminus K$).

We call $\pi_1(X)$ the **knot group** of K .

Remark: The knot complement is a complete topological invariant for knots, but not for links. Though we still utilize this for many examples.

The Wirtinger Presentation

An algorithm to compute the knot group:

The Wirtinger Presentation

An algorithm to compute the knot group:

- ◇ Choose an orientation for K .



The Wirtinger Presentation

An algorithm to compute the knot group:

◇ Choose an orientation for K .




◇ The knot diagram consists of n arcs, separated by n over-under crossings. Label each arc $\alpha_1, \alpha_2, \dots, \alpha_n$ so that α_i connects to α_{i+1} and α_n to α_1 .




The Wirtinger Presentation

- At each α_i , draw an arrow labeled x_i passing under the arc using a *right-hand rule*. These arrows represent loops in the knot complement.




The Wirtinger Presentation

- At each α_i , draw an arrow labeled x_i passing under the arc using a *right-hand rule*. These arrows represent loops in the knot complement.



Case 1:


- At each over-under crossing, observe the relations described by the x_i 's.



$$r_i : x_k x_{i+1} = x_i x_k$$


The Wirtinger Presentation

- At each α_i , draw an arrow labeled x_i passing under the arc using a *right-hand rule*. These arrows represent loops in the knot complement.




Case 1:

- At each over-under crossing, observe the relations described by the x_i 's.



$$r_i : x_k x_{i+1} = x_i x_k$$


Case 2:



$$r_i : x_{i+1} x_k = x_k x_i$$


The Wirtinger Presentation

- At each α_i , draw an arrow labeled x_i passing under the arc using a *right-hand rule*. These arrows represent loops in the knot complement.




Case 1:

- At each over-under crossing, observe the relations described by the x_i 's.




$$r_i : x_k x_{i+1} = x_i x_k$$

Case 2:




$$r_i : x_{i+1} x_k = x_k x_i$$


This yields $\pi_1(X) = \langle x_1, \dots, x_n \mid r_1, \dots, r_n \rangle$.




Example: The Trefoil




Example: The Trefoil




$$r_1 : x_1 x_3 = x_2 x_1$$



Example: The Trefoil




$$r_1 : x_1 x_3 = x_2 x_1$$




$$r_1 : x_1 x_3 = x_3 x_2$$




Example: The Trefoil



$$r_1 : x_1 x_3 = x_2 x_1$$




$$r_1 : x_1 x_3 = x_3 x_2$$




$$r_1 : x_3 x_2 = x_2 x_1$$




Example: The Trefoil



$$r_1 : x_1 x_3 = x_2 x_1$$



$$r_1 : x_1 x_3 = x_3 x_2$$




$$r_1 : x_3 x_2 = x_2 x_1$$

$$\begin{aligned} \implies \pi_1(X) &= \langle x_1, x_2, x_3 \mid x_1 x_3 = x_2 x_1 = x_3 x_2 \rangle \\ &= \langle x_1, x_2 \mid x_1 x_2 x_1 = x_2 x_1 x_2 \rangle \end{aligned}$$


The Magic Trick Revealed

The Magic Trick Revealed

Case 1:




Case 2:



The Magic Trick Revealed


Let $L_1 = A \cup B$ and $X_1 = \mathbb{R}^3 \setminus L_1$.



We have $\pi_1(X_1) = \langle x, y | - \rangle$.


The Magic Trick Revealed

Let $L_1 = A \cup B$ and $X_1 = \mathbb{R}^3 \setminus L_1$.




We have $\pi_1(X_1) = \langle x, y | - \rangle$.

Now consider what $C \in \pi_1(X_1)$:




The Magic Trick Revealed

Let $L_1 = A \cup B$ and $X_1 = \mathbb{R}^3 \setminus L_1$.



We have $\pi_1(X_1) = \langle x, y | - \rangle$.


Now consider what $C \in \pi_1(X_1)$:



By observation $C = xyx^{-1}y^{-1} = [x, y] \in \pi_1(X_1)$, the commutator.


The Magic Trick Revealed

Now let $L_2 = A \cup B$ with A and B linked and $X_2 = \mathbb{R}^3 \setminus L_2$.




The Magic Trick Revealed

Now let $L_2 = A \cup B$ with A and B linked and $X_2 = \mathbb{R}^3 \setminus L_2$.




Consider the relation defined at the over-under crossings.




The Magic Trick Revealed

Now let $L_2 = A \cup B$ with A and B linked and $X_2 = \mathbb{R}^3 \setminus L_2$.



Consider the relation defined at the over-under crossings.



This yields $\pi_1(X_2) = \langle x, y | xy = yx \rangle$.

The Magic Trick Revealed


- ◇ We preserved the configuration of how C was linked to $A \cup B$, and therefore C still represents the commutator in $\pi_1(X_2)$.

The Magic Trick Revealed

- ◇ We preserved the configuration of how C was linked to $A \cup B$, and therefore C still represents the commutator in $\pi_1(X_2)$.
- ◇ Since the knot group of L_2 is abelian, the commutator is trivial and therefore C is unlinked from $A \cup B$.

The Magic Trick Revealed

- ◇ We preserved the configuration of how C was linked to $A \cup B$, and therefore C still represents the commutator in $\pi_1(X_2)$.
- ◇ Since the knot group of L_2 is abelian, the commutator is trivial and therefore C is unlinked from $A \cup B$.



Bibliography

References

- [1] D. Rolfsen, *Knots and Links*. AMS Chelsea Publishing, Rhode Island, 1990, Reprinted with corrections: 2003, pp. 1–65.

Thank you!

