

Cohomology on Sites

UCSC Graduate Colloquium

Deewang Bhamidipati

10th January 2022

Sheaves

× × ×

Category of Open Sets

Let (X, \mathcal{T}_X) be a topological space,

Category of Open Sets

Let (X, \mathcal{T}_X) be a topological space, then \mathcal{T}_X becomes a poset with respect to the partial order given by set containment.

Category of Open Sets

Let (X, \mathcal{T}_X) be a topological space, then \mathcal{T}_X becomes a poset with respect to the partial order given by set containment. We can always categorify a poset, which in the case of \mathcal{T}_X we denote as $\text{Open}(X)$ and is described as follows:

Category of Open Sets

Let (X, \mathcal{T}_X) be a topological space, then \mathcal{T}_X becomes a poset with respect to the partial order given by set containment. We can always categorify a poset, which in the case of \mathcal{T}_X we denote as $\text{Open}(X)$ and is described as follows:

- The objects of $\text{Open}(X)$ are the open sets of X .

Category of Open Sets

Let (X, \mathcal{T}_X) be a topological space, then \mathcal{T}_X becomes a poset with respect to the partial order given by set containment. We can always categorify a poset, which in the case of \mathcal{T}_X we denote as $\text{Open}(X)$ and is described as follows:

- The objects of $\text{Open}(X)$ are the open sets of X .
- Given two open sets U, V , say

$$\text{Hom}_{\text{Open}(X)}(U, V) = \begin{cases} \{U \hookrightarrow V\} & \text{if } U \subseteq V \\ \emptyset & \text{otherwise} \end{cases}$$

Category of Open Sets

Let (X, \mathcal{T}_X) be a topological space, then \mathcal{T}_X becomes a poset with respect to the partial order given by set containment. We can always categorify a poset, which in the case of \mathcal{T}_X we denote as $\text{Open}(X)$ and is described as follows:

- The objects of $\text{Open}(X)$ are the open sets of X .
- Given two open sets U, V , say

$$\text{Hom}_{\text{Open}(X)}(U, V) = \begin{cases} \{U \hookrightarrow V\} & \text{if } U \subseteq V \\ \emptyset & \text{otherwise} \end{cases}$$

Note that X is the final object of $\text{Open}(X)$

Category of Open Sets

Let (X, \mathcal{T}_X) be a topological space, then \mathcal{T}_X becomes a poset with respect to the partial order given by set containment. We can always categorify a poset, which in the case of \mathcal{T}_X we denote as $\text{Open}(X)$ and is described as follows:

- The objects of $\text{Open}(X)$ are the open sets of X .
- Given two open sets U, V , say

$$\text{Hom}_{\text{Open}(X)}(U, V) = \begin{cases} \{U \hookrightarrow V\} & \text{if } U \subseteq V \\ \emptyset & \text{otherwise} \end{cases}$$

Note that X is the final object of $\text{Open}(X)$ and finite (fibered) products exist and are given by finite intersections.

Category of Open Sets

Let (X, \mathcal{T}_X) be a topological space, then \mathcal{T}_X becomes a poset with respect to the partial order given by set containment. We can always categorify a poset, which in the case of \mathcal{T}_X we denote as $\text{Open}(X)$ and is described as follows:

- The objects of $\text{Open}(X)$ are the open sets of X .
- Given two open sets U, V , say

$$\text{Hom}_{\text{Open}(X)}(U, V) = \begin{cases} \{U \hookrightarrow V\} & \text{if } U \subseteq V \\ \emptyset & \text{otherwise} \end{cases}$$

Note that X is the final object of $\text{Open}(X)$ and finite (fibered) products exist and are given by finite intersections. We reiterate,

Topology Categorized

Category of Open Sets

Let (X, \mathcal{T}_X) be a topological space, then \mathcal{T}_X becomes a poset with respect to the partial order given by set containment. We can always categorify a poset, which in the case of \mathcal{T}_X we denote as $\text{Open}(X)$ and is described as follows:

- The objects of $\text{Open}(X)$ are the open sets of X .
- Given two open sets U, V , say

$$\text{Hom}_{\text{Open}(X)}(U, V) = \begin{cases} \{U \hookrightarrow V\} & \text{if } U \subseteq V \\ \emptyset & \text{otherwise} \end{cases}$$

Note that X is the final object of $\text{Open}(X)$ and finite (fibered) products exist and are given by finite intersections. We reiterate, the following is a Cartesian square in $\text{Open}(X)$,

$$\begin{array}{ccc} U \cap V & \hookrightarrow & U \\ \downarrow \lrcorner & & \downarrow \\ V & \hookrightarrow & X \end{array}$$

that is, $U \times_X V = U \cap V$ in $\text{Open}(X)$.

Presheaves

A *presheaf of abelian groups on X* , or simply a *presheaf*, is a functor $\mathcal{F} : \text{Open}(X)^{\text{op}} \rightarrow \text{Ab}$.

Presheaves

A *presheaf of abelian groups on X* , or simply a *presheaf*, is a functor $\mathcal{F} : \text{Open}(X)^{\text{op}} \rightarrow \text{Ab}$. That is, it consists of the following data

Presheaves

A *presheaf of abelian groups on X* , or simply a *presheaf*, is a functor $\mathcal{F} : \text{Open}(X)^{\text{op}} \rightarrow \text{Ab}$. That is, it consists of the following data

- For each open set U , an abelian group $\mathcal{F}(U)$.

Presheaves

A *presheaf of abelian groups on X* , or simply a *presheaf*, is a functor $\mathcal{F} : \text{Open}(X)^{\text{op}} \rightarrow \text{Ab}$. That is, it consists of the following data

- For each open set U , an abelian group $\mathcal{F}(U)$. Other notations include $\Gamma(U, \mathcal{F})$ and $H^0(U, \mathcal{F})$.

Presheaves

A *presheaf of abelian groups on X* , or simply a *presheaf*, is a functor $\mathcal{F} : \text{Open}(X)^{\text{op}} \rightarrow \text{Ab}$. That is, it consists of the following data

- For each open set U , an abelian group $\mathcal{F}(U)$. Other notations include $\Gamma(U, \mathcal{F})$ and $H^0(U, \mathcal{F})$. The elements of $\mathcal{F}(U)$ are called *sections of \mathcal{F} over U* ;

Presheaves

A *presheaf of abelian groups on X* , or simply a *presheaf*, is a functor $\mathcal{F} : \text{Open}(X)^{\text{op}} \rightarrow \text{Ab}$. That is, it consists of the following data

- For each open set U , an abelian group $\mathcal{F}(U)$. Other notations include $\Gamma(U, \mathcal{F})$ and $H^0(U, \mathcal{F})$. The elements of $\mathcal{F}(U)$ are called *sections of \mathcal{F} over U* ; in the particular case when $U = X$, the elements are called *global sections*.

Presheaves

A *presheaf of abelian groups on X* , or simply a *presheaf*, is a functor $\mathcal{F} : \text{Open}(X)^{\text{op}} \rightarrow \text{Ab}$. That is, it consists of the following data

- For each open set U , an abelian group $\mathcal{F}(U)$. Other notations include $\Gamma(U, \mathcal{F})$ and $H^0(U, \mathcal{F})$. The elements of $\mathcal{F}(U)$ are called *sections of \mathcal{F} over U* ; in the particular case when $U = X$, the elements are called *global sections*.
- Whenever $U \hookrightarrow V$, a group homomorphism $\text{res}_{V,U} : \mathcal{F}(V) \rightarrow \mathcal{F}(U)$, which is usually called a *restriction map*.

Presheaves

A *presheaf of abelian groups on X* , or simply a *presheaf*, is a functor $\mathcal{F} : \text{Open}(X)^{\text{op}} \rightarrow \text{Ab}$. That is, it consists of the following data

- For each open set U , an abelian group $\mathcal{F}(U)$. Other notations include $\Gamma(U, \mathcal{F})$ and $H^0(U, \mathcal{F})$. The elements of $\mathcal{F}(U)$ are called *sections of \mathcal{F} over U* ; in the particular case when $U = X$, the elements are called *global sections*.
- Whenever $U \hookrightarrow V$, a group homomorphism $\text{res}_{V,U} : \mathcal{F}(V) \rightarrow \mathcal{F}(U)$, which is usually called a *restriction map*. For an element $f \in \mathcal{F}(V)$, if V is clear from context, we simply write $f|_U$ for its image under the restriction map $\text{res}_{V,U}$.

Presheaves

A *presheaf of abelian groups on X* , or simply a *presheaf*, is a functor $\mathcal{F} : \text{Open}(X)^{\text{op}} \rightarrow \text{Ab}$. That is, it consists of the following data

- For each open set U , an abelian group $\mathcal{F}(U)$. Other notations include $\Gamma(U, \mathcal{F})$ and $H^0(U, \mathcal{F})$. The elements of $\mathcal{F}(U)$ are called *sections of \mathcal{F} over U* ; in the particular case when $U = X$, the elements are called *global sections*.
- Whenever $U \hookrightarrow V$, a group homomorphism $\text{res}_{V,U} : \mathcal{F}(V) \rightarrow \mathcal{F}(U)$, which is usually called a *restriction map*. For an element $f \in \mathcal{F}(V)$, if V is clear from context, we simply write $f|_U$ for its image under the restriction map $\text{res}_{V,U}$.
- In particular, $\text{res}_{U,U} = \text{id}_{\mathcal{F}(U)}$ for any open set U .

Presheaves

A *presheaf of abelian groups on X* , or simply a *presheaf*, is a functor $\mathcal{F} : \text{Open}(X)^{\text{op}} \rightarrow \text{Ab}$. That is, it consists of the following data

- For each open set U , an abelian group $\mathcal{F}(U)$. Other notations include $\Gamma(U, \mathcal{F})$ and $H^0(U, \mathcal{F})$. The elements of $\mathcal{F}(U)$ are called *sections of \mathcal{F} over U* ; in the particular case when $U = X$, the elements are called *global sections*.
- Whenever $U \hookrightarrow V$, a group homomorphism $\text{res}_{V,U} : \mathcal{F}(V) \rightarrow \mathcal{F}(U)$, which is usually called a *restriction map*. For an element $f \in \mathcal{F}(V)$, if V is clear from context, we simply write $f|_U$ for its image under the restriction map $\text{res}_{V,U}$.
- In particular, $\text{res}_{U,U} = \text{id}_{\mathcal{F}(U)}$ for any open set U .
- Finally, whenever $U \hookrightarrow V \hookrightarrow W$, the following diagram commutes in Ab

$$\begin{array}{ccc} \mathcal{F}(W) & \xrightarrow{\text{res}_{W,U}} & \mathcal{F}(U) \\ & \searrow \text{res}_{W,V} & \nearrow \text{res}_{V,U} \\ & \mathcal{F}(V) & \end{array}$$

Presheaves

A *presheaf of abelian groups on X* , or simply a *presheaf*, is a functor $\mathcal{F} : \text{Open}(X)^{\text{op}} \rightarrow \text{Ab}$. That is, it consists of the following data

- For each open set U , an abelian group $\mathcal{F}(U)$. Other notations include $\Gamma(U, \mathcal{F})$ and $H^0(U, \mathcal{F})$. The elements of $\mathcal{F}(U)$ are called *sections of \mathcal{F} over U* ; in the particular case when $U = X$, the elements are called *global sections*.
- Whenever $U \hookrightarrow V$, a group homomorphism $\text{res}_{V,U} : \mathcal{F}(V) \rightarrow \mathcal{F}(U)$, which is usually called a *restriction map*. For an element $f \in \mathcal{F}(V)$, if V is clear from context, we simply write $f|_U$ for its image under the restriction map $\text{res}_{V,U}$.
- In particular, $\text{res}_{U,U} = \text{id}_{\mathcal{F}(U)}$ for any open set U .
- Finally, whenever $U \hookrightarrow V \hookrightarrow W$, the following diagram commutes in Ab

$$\begin{array}{ccc} \mathcal{F}(W) & \xrightarrow{\text{res}_{W,U}} & \mathcal{F}(U) \\ & \searrow \text{res}_{W,V} & \nearrow \text{res}_{V,U} \\ & \mathcal{F}(V) & \end{array}$$

One can suitably replace the target category to obtain *presheaf of sets, groups, rings, modules, algebras* etc.

Examples I

The examples to keep in mind are the following, these will also be our motivating examples of sheaves.

Examples I

The examples to keep in mind are the following, these will also be our motivating examples of sheaves.

- Consider a topological space (resp. smooth manifold) X ,

Examples I

The examples to keep in mind are the following, these will also be our motivating examples of sheaves.

- Consider a topological space (resp. smooth manifold) X , then associating to each open set U the ring of \mathbb{R} -valued continuous (resp. smooth) functions on U gives us a presheaf (of rings).

$$\begin{aligned}\mathcal{F}(U) &= \{f: U \rightarrow \mathbb{R} \mid f|_W \text{ smooth}\} \\ f &\mapsto f|_U\end{aligned}$$

Examples I

The examples to keep in mind are the following, these will also be our motivating examples of sheaves.

- Consider a topological space (resp. smooth manifold) X , then associating to each open set U the ring of \mathbb{R} -valued continuous (resp. smooth) functions on U gives us a presheaf (of rings).
- If X is a complex manifold,

Examples I

The examples to keep in mind are the following, these will also be our motivating examples of sheaves.

- Consider a topological space (resp. smooth manifold) X , then associating to each open set U the ring of \mathbb{R} -valued continuous (resp. smooth) functions on U gives us a presheaf (of rings).
- If X is a complex manifold, similarly associating to each open set the ring of \mathbb{C} -valued holomorphic functions gives us a presheaf (of rings).

Examples I

The examples to keep in mind are the following, these will also be our motivating examples of sheaves.

- Consider a topological space (resp. smooth manifold) X , then associating to each open set U the ring of \mathbb{R} -valued continuous (resp. smooth) functions on U gives us a presheaf (of rings).
- If X is a complex manifold, similarly associating to each open set the ring of \mathbb{C} -valued holomorphic functions gives us a presheaf (of rings).
- If X is a smooth (resp. complex) manifold,

Examples I

The examples to keep in mind are the following, these will also be our motivating examples of sheaves.

- Consider a topological space (resp. smooth manifold) X , then associating to each open set U the ring of \mathbb{R} -valued continuous (resp. smooth) functions on U gives us a presheaf (of rings).
- If X is a complex manifold, similarly associating to each open set the ring of \mathbb{C} -valued holomorphic functions gives us a presheaf (of rings).
- If X is a smooth (resp. complex) manifold, associating to each open set U the abelian group of smooth (resp. holomorphic) differential forms on U gives us a presheaf (of abelian groups).

$$U \mapsto \Omega^p(U)$$

Examples I

The examples to keep in mind are the following, these will also be our motivating examples of sheaves.

- Consider a topological space (resp. smooth manifold) X , then associating to each open set U the ring of \mathbb{R} -valued continuous (resp. smooth) functions on U gives us a presheaf (of rings).
- If X is a complex manifold, similarly associating to each open set the ring of \mathbb{C} -valued holomorphic functions gives us a presheaf (of rings).
- If X is a smooth (resp. complex) manifold, associating to each open set U the abelian group of smooth (resp. holomorphic) differential forms on U gives us a presheaf (of abelian groups).

In all these examples, the restriction maps are just that, honest restriction of functions.

Examples II

Some other important examples are as following:

Examples II

Some other important examples are as following:

- *Constant presheaf*: Let X be a topological space, and A an abelian group.

Examples II

Some other important examples are as following:

- *Constant presheaf*: Let X be a topological space, and A an abelian group. Then $\underline{A}_{\text{pre}}$ defined as $\underline{A}_{\text{pre}}(U) = A$,

Examples II

Some other important examples are as following:

- *Constant presheaf*: Let X be a topological space, and A an abelian group. Then $\underline{A}_{\text{pre}}$ defined as $\underline{A}_{\text{pre}}(U) = A$, equivalently

$$\underline{A}_{\text{pre}}(U) = \{f : U \rightarrow A \mid f \text{ a constant function}\}, //$$

for all open sets U , is a presheaf of abelian groups

Examples II

Some other important examples are as following:

- *Constant presheaf*: Let X be a topological space, and A an abelian group. Then $\underline{A}_{\text{pre}}$ defined as $\underline{A}_{\text{pre}}(U) = A$, equivalently

$$\underline{A}_{\text{pre}}(U) = \{f : U \rightarrow A \mid f \text{ a constant function}\},$$

for all open sets U , is a presheaf of abelian groups called the *constant presheaf associated to A* .

Examples II

Some other important examples are as following:

- *Constant presheaf*: Let X be a topological space, and A an abelian group. Then $\underline{A}_{\text{pre}}$ defined as $\underline{A}_{\text{pre}}(U) = A$, equivalently

$$\underline{A}_{\text{pre}}(U) = \{f : U \rightarrow A \mid f \text{ a constant function}\},$$

for all open sets U , is a presheaf of abelian groups called the *constant presheaf associated to A* .

For the first interpretation, the restriction maps are simply id_A ;

Examples II

Some other important examples are as following:

- *Constant presheaf*: Let X be a topological space, and A an abelian group. Then $\underline{A}_{\text{pre}}$ defined as $\underline{A}_{\text{pre}}(U) = A$, equivalently

$$\underline{A}_{\text{pre}}(U) = \{f : U \rightarrow A \mid f \text{ a constant function}\},$$

for all open sets U , is a presheaf of abelian groups called the *constant presheaf associated to A* .

For the first interpretation, the restriction maps are simply id_A ; while for the second the restrictions maps are again restriction maps of functions.

Examples II

Some other important examples are as following:

- *Constant presheaf*: Let X be a topological space, and A an abelian group. Then $\underline{A}_{\text{pre}}$ defined as $\underline{A}_{\text{pre}}(U) = A$, equivalently

$$\underline{A}_{\text{pre}}(U) = \{f : U \rightarrow A \mid f \text{ a constant function}\},$$

for all open sets U , is a presheaf of abelian groups called the *constant presheaf associated to A* .

For the first interpretation, the restriction maps are simply id_A ; while for the second the restrictions maps are again restriction maps of functions.

- *Skyscraper presheaf*: Suppose X is a topological space, with $p \in X$, and A is an abelian group.

Examples II

Some other important examples are as following:

- *Constant presheaf*: Let X be a topological space, and A an abelian group. Then $\underline{A}_{\text{pre}}$ defined as $\underline{A}_{\text{pre}}(U) = A$, equivalently

$$\underline{A}_{\text{pre}}(U) = \{f : U \rightarrow A \mid f \text{ a constant function}\},$$

for all open sets U , is a presheaf of abelian groups called the *constant presheaf associated to A* .

For the first interpretation, the restriction maps are simply id_A ; while for the second the restrictions maps are again restriction maps of functions.

- *Skyscraper presheaf*: Suppose X is a topological space, with $p \in X$, and A is an abelian group. Let $i(p) : p \hookrightarrow X$ be the inclusion.

Examples II

Some other important examples are as following:

- *Constant presheaf*: Let X be a topological space, and A an abelian group. Then $\underline{A}_{\text{pre}}$ defined as $\underline{A}_{\text{pre}}(U) = A$, equivalently

$$\underline{A}_{\text{pre}}(U) = \{f : U \rightarrow A \mid f \text{ a constant function}\},$$

for all open sets U , is a presheaf of abelian groups called the *constant presheaf associated to A* .

For the first interpretation, the restriction maps are simply id_A ; while for the second the restrictions maps are again restriction maps of functions.

- *Skyscraper presheaf*: Suppose X is a topological space, with $p \in X$, and A is an abelian group. Let $i(p) : p \hookrightarrow X$ be the inclusion. Then $i(p)_* \underline{A}$ defined by

$$[i(p)_* \underline{A}](U) = \begin{cases} A & \text{if } p \in U \\ \{0\} & \text{otherwise} \end{cases}$$

forms a presheaf.

Examples II

Some other important examples are as following:

- *Constant presheaf*: Let X be a topological space, and A an abelian group. Then $\underline{A}_{\text{pre}}$ defined as $\underline{A}_{\text{pre}}(U) = A$, equivalently

$$\underline{A}_{\text{pre}}(U) = \{f : U \rightarrow A \mid f \text{ a constant function}\},$$

for all open sets U , is a presheaf of abelian groups called the *constant presheaf associated to A* .

For the first interpretation, the restriction maps are simply id_A ; while for the second the restrictions maps are again restriction maps of functions.

- *Skyscraper presheaf*: Suppose X is a topological space, with $p \in X$, and A is an abelian group. Let $i(p) : p \hookrightarrow X$ be the inclusion. Then $i(p)_* \underline{A}$ defined by

$$[i(p)_* \underline{A}](U) = \begin{cases} A & \text{if } p \in U \\ \{0\} & \text{otherwise} \end{cases}$$

forms a presheaf. The restriction maps are either the 0 map or id_A .


Sheaves

A *sheaf of abelian groups on X* , or simply a *sheaf*, is a presheaf \mathcal{F} which satisfies the following two axioms

Sheaves

A *sheaf of abelian groups on X* , or simply a *sheaf*, is a presheaf \mathcal{F} which satisfies the following two axioms

(S1) *Identity axiom.* Suppose $\{U_i\}_{i \in I}$ is an open cover of an open set U , and $f \in \mathcal{F}(U)$ is such that $f|_{U_i} = 0$ for all i ,



Sheaves

A *sheaf of abelian groups on X* , or simply a *sheaf*, is a presheaf \mathcal{F} which satisfies the following two axioms

(S1) *Identity axiom.* Suppose $\{U_i\}_{i \in I}$ is an open cover of an open set U , and $f \in \mathcal{F}(U)$ is such that $f|_{U_i} = 0$ for all i , then $f = 0$.

Sheaves

A *sheaf of abelian groups on X* , or simply a *sheaf*, is a presheaf \mathcal{F} which satisfies the following two axioms

(S1) *Identity axiom.* Suppose $\{U_i\}_{i \in I}$ is an open cover of an open set U , and $f \in \mathcal{F}(U)$ is such that $f|_{U_i} = 0$ for all i , then $f = 0$.

A presheaf that only satisfies this axiom is called a *separated presheaf*.

Sheaves

A *sheaf of abelian groups on X* , or simply a *sheaf*, is a presheaf \mathcal{F} which satisfies the following two axioms

(S1) *Identity axiom.* Suppose $\{U_i\}_{i \in I}$ is an open cover of an open set U , and $f \in \mathcal{F}(U)$ is such that $f|_{U_i} = 0$ for all i , then $f = 0$.

A presheaf that only satisfies this axiom is called a *separated presheaf*.

(S2) *Gluability axiom.* Suppose $\{U_i\}_{i \in I}$ is an open cover of an open set U , and $f_i \in \mathcal{F}(U_i)$ are such that $f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$ for all i, j ,



Sheaves

A *sheaf of abelian groups on X* , or simply a *sheaf*, is a presheaf \mathcal{F} which satisfies the following two axioms

(S1) *Identity axiom.* Suppose $\{U_i\}_{i \in I}$ is an open cover of an open set U , and $f \in \mathcal{F}(U)$ is such that $f|_{U_i} = 0$ for all i , then $f = 0$.

A presheaf that only satisfies this axiom is called a *separated presheaf*.

(S2) *Glueability axiom.* Suppose $\{U_i\}_{i \in I}$ is an open cover of an open set U , and $f_i \in \mathcal{F}(U_i)$ are such that $f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$ for all i, j , then there exists a $f \in \mathcal{F}(U)$ such that $f|_{U_i} = f_i$ for all i .

Sheaves

A *sheaf of abelian groups on X* , or simply a *sheaf*, is a presheaf \mathcal{F} which satisfies the following two axioms

(S1) *Identity axiom.* Suppose $\{U_i\}_{i \in I}$ is an open cover of an open set U , and $f \in \mathcal{F}(U)$ is such that $f|_{U_i} = 0$ for all i , then $f = 0$.

A presheaf that only satisfies this axiom is called a *separated presheaf*.

(S2) *Glueability axiom.* Suppose $\{U_i\}_{i \in I}$ is an open cover of an open set U , and $f_i \in \mathcal{F}(U_i)$ are such that $f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$ for all i, j , then there exists a $f \in \mathcal{F}(U)$ such that $f|_{U_i} = f_i$ for all i . The previous axiom tells us that such an f is unique.

(...)

Sheaves

(...) Equivalently, these axioms can be succinctly stated as saying the following sequence

$$0 \longrightarrow \mathcal{F}(U) \xrightarrow{f} \prod_i \mathcal{F}(U_i) \xrightarrow{(\text{fla})} \prod_{i,j} \mathcal{F}(U_i \cap U_j)$$

$\rho := (\text{res}_{U_i, U_i \cap U_j})_{i,j}$
 $\rho_i := (\text{res}_{U_i, U_i \cap U_j})_{i,j}$
 $\rho_j := (\text{res}_{U_j, U_i \cap U_j})_{i,j}$

is exact for a given covering of $\{U_i\}_{i \in I}$ of U .

Sheaves

(...) Equivalently, these axioms can be succinctly stated as saying the following sequence

$$0 \longrightarrow \mathcal{F}(U) \xrightarrow{\rho := (\text{res}_{U, U_i})_i} \prod_i \mathcal{F}(U_i) \begin{array}{c} \xrightarrow{\rho_i := (\text{res}_{U_i, U_i \cap U_j})_{i,j}} \\ \xrightarrow{\rho_j := (\text{res}_{U_j, U_i \cap U_j})_{i,j}} \end{array} \prod_{i,j} \mathcal{F}(U_i \cap U_j)$$

is exact for a given covering of $\{U_i\}_{i \in I}$ of U .

(S1) corresponds to exactness at $\mathcal{F}(U)$, that is the injectivity of ρ .

Sheaves

(...) Equivalently, these axioms can be succinctly stated as saying the following sequence

$$0 \longrightarrow \mathcal{F}(U) \xrightarrow{\rho := (\text{res}_{U, U_i})_i} \prod_i \mathcal{F}(U_i) \begin{array}{c} \xrightarrow{\rho_i := (\text{res}_{U_i, U_i \cap U_j})_{i,j}} \\ \xrightarrow{\rho_j := (\text{res}_{U_j, U_i \cap U_j})_{i,j}} \end{array} \prod_{i,j} \mathcal{F}(U_i \cap U_j)$$

is exact for a given covering of $\{U_i\}_{i \in I}$ of U .

(S1) corresponds to exactness at $\mathcal{F}(U)$, that is the injectivity of ρ . While (S2) corresponds to exactness at $\prod_i \mathcal{F}(U_i)$.

Sheaves

(...) Equivalently, these axioms can be succinctly stated as saying the following sequence

$$0 \longrightarrow \mathcal{F}(U) \xrightarrow{\rho := (\text{res}_{U, U_i})_i} \prod_i \mathcal{F}(U_i) \begin{array}{c} \xrightarrow{\rho_i := (\text{res}_{U_i, U_i \cap U_j})_{i,j}} \\ \xrightarrow{\rho_j := (\text{res}_{U_j, U_i \cap U_j})_{i,j}} \end{array} \prod_{i,j} \mathcal{F}(U_i \cap U_j)$$

is exact for a given covering of $\{U_i\}_{i \in I}$ of U .

(S1) corresponds to exactness at $\mathcal{F}(U)$, that is the injectivity of ρ . While (S2) corresponds to exactness at $\prod_i \mathcal{F}(U_i)$. This can be interpreted as saying $\text{im}(\rho) = \text{eq}(\rho_i, \rho_j)$, the equaliser;

Sheaves

(...) Equivalently, these axioms can be succinctly stated as saying the following sequence

$$0 \longrightarrow \mathcal{F}(U) \xrightarrow{\rho := (\text{res}_{U, U_i})_i} \prod_i \mathcal{F}(U_i) \xrightleftharpoons[\rho_j := (\text{res}_{U_j, U_i \cap U_j})_{i,j}]{\rho_i := (\text{res}_{U_i, U_i \cap U_j})_{i,j}} \prod_{i,j} \mathcal{F}(U_i \cap U_j)$$

is exact for a given covering of $\{U_i\}_{i \in I}$ of U .

(S1) corresponds to exactness at $\mathcal{F}(U)$, that is the injectivity of ρ . While (S2) corresponds to exactness at $\prod_i \mathcal{F}(U_i)$. This can be interpreted as saying $\text{im}(\rho) = \text{eq}(\rho_i, \rho_j)$, the equaliser; or equivalently $\text{im}(\rho) = \ker(\rho_i - \rho_j)$, the difference kernel.

$$f \longmapsto f|_{U_i \cap U_j} - f|_{U_j \cap U_i} = 0 \quad \forall i, j$$

Examples

Amongst the examples given previously the only presheaf that is not a sheaf is the constant presheaf,

Examples

Amongst the examples given previously the only presheaf that is not a sheaf is the constant presheaf, morally because constant functions don't necessarily glue to give constant functions.

Examples

Amongst the examples given previously the only presheaf that is not a sheaf is the constant presheaf, morally because constant functions don't necessarily glue to give constant functions.

- For a topological space X , and an abelian group A topologised with the discrete topology,

$$\{f: U \rightarrow A \mid f \text{ is } \cancel{\text{constant}}\}$$

locally constant.

Examples

Amongst the examples given previously the only presheaf that is not a sheaf is the constant presheaf, morally because constant functions don't necessarily glue to give constant functions.

- For a topological space X , and an abelian group A topologised with the discrete topology, define \underline{A} as follows

$$\underline{A}(U) = \{f : U \rightarrow A \mid f \text{ is continuous}\},$$

$\mathcal{U} = f^{-1}(a) \leftarrow a$

Examples

Amongst the examples given previously the only presheaf that is not a sheaf is the constant presheaf, morally because constant functions don't necessarily glue to give constant functions.

- For a topological space X , and an abelian group A topologised with the discrete topology, define \underline{A} as follows

$$\underline{A}(U) = \{f : U \rightarrow A \mid f \text{ is continuous}\},$$

that is, $\underline{A}(U)$ is the abelian group of *locally constant* (A -valued) functions on U .

Examples

Amongst the examples given previously the only presheaf that is not a sheaf is the constant presheaf, morally because constant functions don't necessarily glue to give constant functions.

- For a topological space X , and an abelian group A topologised with the discrete topology, define \underline{A} as follows

$$\underline{A}(U) = \{f : U \rightarrow A \mid f \text{ is continuous}\},$$

that is, $\underline{A}(U)$ is the abelian group of *locally constant* (A -valued) functions on U . \underline{A} is a sheaf, called the *constant sheaf associated to A* .

Examples

Amongst the examples given previously the only presheaf that is not a sheaf is the constant presheaf, morally because constant functions don't necessarily glue to give constant functions.

- For a topological space X , and an abelian group A topologised with the discrete topology, define \underline{A} as follows

$$\underline{A}(U) = \{f : U \rightarrow A \mid f \text{ is continuous}\},$$

that is, $\underline{A}(U)$ is the abelian group of *locally constant* (A -valued) functions on U . \underline{A} is a sheaf, called the *constant sheaf associated to A* .

The following is an example of a sheaf that motivates our use of the term sections:

Examples

Amongst the examples given previously the only presheaf that is not a sheaf is the constant presheaf, morally because constant functions don't necessarily glue to give constant functions.

- For a topological space X , and an abelian group A topologised with the discrete topology, define \underline{A} as follows

$$\underline{A}(U) = \{f : U \rightarrow A \mid f \text{ is continuous}\},$$

that is, $\underline{A}(U)$ is the abelian group of *locally constant* (A -valued) functions on U . \underline{A} is a sheaf, called the *constant sheaf associated to A* .

The following is an example of a sheaf that motivates our use of the term sections:

- Given a continuous map $\mu : Y \rightarrow X$

Examples

Amongst the examples given previously the only presheaf that is not a sheaf is the constant presheaf, morally because constant functions don't necessarily glue to give constant functions.

- For a topological space X , and an abelian group A topologised with the discrete topology, define \underline{A} as follows

$$\underline{A}(U) = \{f : U \rightarrow A \mid f \text{ is continuous}\},$$

that is, $\underline{A}(U)$ is the abelian group of *locally constant* (A -valued) functions on U . \underline{A} is a sheaf, called the *constant sheaf associated to A* .

The following is an example of a sheaf that motivates our use of the term sections:

- Given a continuous map $\mu : Y \rightarrow X$ then for an open set U the collection of local sections of μ on U

Examples

Amongst the examples given previously the only presheaf that is not a sheaf is the constant presheaf, morally because constant functions don't necessarily glue to give constant functions.

- For a topological space X , and an abelian group A topologised with the discrete topology, define \underline{A} as follows

$$\underline{A}(U) = \{f : U \rightarrow A \mid f \text{ is continuous}\},$$

that is, $\underline{A}(U)$ is the abelian group of *locally constant* (A -valued) functions on U . \underline{A} is a sheaf, called the *constant sheaf associated to A* .

The following is an example of a sheaf that motivates our use of the term sections:

- Given a continuous map $\mu : Y \rightarrow X$ then for an open set U the collection of local sections of μ on U

$$\Gamma(U, \mu) = \{\sigma : U \rightarrow Y \mid \sigma \text{ is continuous and } \mu \circ \sigma = \text{id}|_U\}$$

defines a sheaf.

(...)

Examples

(...) Some more examples of presheaves that are not sheaves are:

- The presheaf of bounded functions on \mathbb{R} (or \mathbb{C}).
- The presheaf of holomorphic functions on \mathbb{C} that admit a holomorphic square root.

Examples

(...) Some more examples of presheaves that are not sheaves are:

- The presheaf of bounded functions on \mathbb{R} (or \mathbb{C}).
- The presheaf of holomorphic functions on \mathbb{C} that admit a holomorphic square root.

If A is a commutative ring with unity, we can consider the set of prime ideals $\text{Spec } A$. The *Zariski topology* on $\text{Spec } A$ has as its basis of topology the collection

$$D(f) = \{\mathfrak{p} \in \text{Spec } A \mid f \notin \mathfrak{p}\}$$

Examples

(...) Some more examples of presheaves that are not sheaves are:

- The presheaf of bounded functions on \mathbb{R} (or \mathbb{C}).
- The presheaf of holomorphic functions on \mathbb{C} that admit a holomorphic square root.

If A is a commutative ring with unity, we can consider the set of prime ideals $\text{Spec } A$. The *Zariski topology* on $\text{Spec } A$ has as its basis of topology the collection

$$D(f) = \{\mathfrak{p} \in \text{Spec } A \mid f \notin \mathfrak{p}\}$$

It's a fact that it suffices to define a sheaf on the basis of the topology.

Examples

(...) Some more examples of presheaves that are not sheaves are:

- The presheaf of bounded functions on \mathbb{R} (or \mathbb{C}).
- The presheaf of holomorphic functions on \mathbb{C} that admit a holomorphic square root.

If A is a commutative ring with unity, we can consider the set of prime ideals $\text{Spec } A$. The *Zariski topology* on $\text{Spec } A$ has as its basis of topology the collection

$$D(f) = \{\mathfrak{p} \in \text{Spec } A \mid f \notin \mathfrak{p}\}$$

It's a fact that it suffices to define a sheaf on the basis of the topology.

- Associate to each $D(f)$, the localisation A_f .

Examples

(...) Some more examples of presheaves that are not sheaves are:

- The presheaf of bounded functions on \mathbb{R} (or \mathbb{C}).
- The presheaf of holomorphic functions on \mathbb{C} that admit a holomorphic square root.

If A is a commutative ring with unity, we can consider the set of prime ideals $\text{Spec } A$. The *Zariski topology* on $\text{Spec } A$ has as its basis of topology the collection

$$D(f) = \{\mathfrak{p} \in \text{Spec } A \mid f \notin \mathfrak{p}\}$$

It's a fact that it suffices to define a sheaf on the basis of the topology.

- Associate to each $D(f)$, the localisation A_f . Whenever $D(f) \subseteq D(g)$, or equivalently $f^n \in (g)$ for some positive n ,

Examples

(...) Some more examples of presheaves that are not sheaves are:

- The presheaf of bounded functions on \mathbb{R} (or \mathbb{C}).
- The presheaf of holomorphic functions on \mathbb{C} that admit a holomorphic square root.

If A is a commutative ring with unity, we can consider the set of prime ideals $\text{Spec } A$. The *Zariski topology* on $\text{Spec } A$ has as its basis of topology the collection

$$D(f) = \{\mathfrak{p} \in \text{Spec } A \mid f \notin \mathfrak{p}\}$$

It's a fact that it suffices to define a sheaf on the basis of the topology.

- Associate to each $D(f)$, the localisation A_f . Whenever $D(f) \subseteq D(g)$, or equivalently $f^n \in (g)$ for some positive n , let the localisation maps $A_g \rightarrow A_f$ be the restriction maps.

Examples

(...) Some more examples of presheaves that are not sheaves are:

- The presheaf of bounded functions on \mathbb{R} (or \mathbb{C}).
- The presheaf of holomorphic functions on \mathbb{C} that admit a holomorphic square root.

If A is a commutative ring with unity, we can consider the set of prime ideals $\text{Spec } A$. The *Zariski topology* on $\text{Spec } A$ has as its basis of topology the collection

$$D(f) = \{\mathfrak{p} \in \text{Spec } A \mid f \notin \mathfrak{p}\}$$

It's a fact that it suffices to define a sheaf on the basis of the topology.

- Associate to each $D(f)$, the localisation A_f . Whenever $D(f) \subseteq D(g)$, or equivalently $f^n \in (g)$ for some positive n , let the localisation maps $A_g \rightarrow A_f$ be the restriction maps. This describes a sheaf (of rings) on $\text{Spec } A$.

Examples

(...) Some more examples of presheaves that are not sheaves are:

- The presheaf of bounded functions on \mathbb{R} (or \mathbb{C}).
- The presheaf of holomorphic functions on \mathbb{C} that admit a holomorphic square root.

If A is a commutative ring with unity, we can consider the set of prime ideals $\text{Spec } A$. The *Zariski topology* on $\text{Spec } A$ has as its basis of topology the collection

$$D(f) = \{\mathfrak{p} \in \text{Spec } A \mid f \notin \mathfrak{p}\}$$

It's a fact that it suffices to define a sheaf on the basis of the topology.

- Associate to each $D(f)$, the localisation A_f . Whenever $D(f) \subseteq D(g)$, or equivalently $f^n \in (g)$ for some positive n , let the localisation maps $A_g \rightarrow A_f$ be the restriction maps. This describes a sheaf (of rings) on $\text{Spec } A$. In particular, the ring of global sections is precisely the ring A .

$$\text{Spec } A = D(1) \mapsto A_1 = A$$

Category of Sheaves

The category of presheaves on X is simply $\mathbf{PSh}(X) := [\mathbf{Open}(X)^{\mathrm{op}}, \mathbf{Ab}]$,

Category of Sheaves

The category of presheaves on X is simply $\mathbf{PSh}(X) := [\mathbf{Open}(X)^{\mathbf{op}}, \mathbf{Ab}]$, and the category of sheaves $\mathbf{Sh}(X)$ is the full subcategory of $\mathbf{PSh}(X)$ with objects as sheaves

$$i : \mathbf{Sh}(X) \hookrightarrow \mathbf{PSh}(X)$$

Category of Sheaves

The category of presheaves on X is simply $\mathbf{PSh}(X) := [\mathbf{Open}(X)^{\mathbf{op}}, \mathbf{Ab}]$, and the category of sheaves $\mathbf{Sh}(X)$ is the full subcategory of $\mathbf{PSh}(X)$ with objects as sheaves

$$i : \mathbf{Sh}(X) \hookrightarrow \mathbf{PSh}(X)$$

(you can interpret i as the forgetful functor).

Category of Sheaves

The category of presheaves on X is simply $\mathbf{PSh}(X) := [\mathbf{Open}(X)^{\mathbf{op}}, \mathbf{Ab}]$, and the category of sheaves $\mathbf{Sh}(X)$ is the full subcategory of $\mathbf{PSh}(X)$ with objects as sheaves

$$i : \mathbf{Sh}(X) \hookrightarrow \mathbf{PSh}(X)$$

(you can interpret i as the forgetful functor).

Therefore, a morphism of sheaves $\phi : \mathcal{F} \rightarrow \mathcal{G}$ is a natural transformation of functors.

Category of Sheaves

The category of presheaves on X is simply $\mathbf{PSh}(X) := [\mathbf{Open}(X)^{\mathbf{op}}, \mathbf{Ab}]$, and the category of sheaves $\mathbf{Sh}(X)$ is the full subcategory of $\mathbf{PSh}(X)$ with objects as sheaves

$$i : \mathbf{Sh}(X) \hookrightarrow \mathbf{PSh}(X)$$

(you can interpret i as the forgetful functor).

Therefore, a morphism of sheaves $\phi : \mathcal{F} \rightarrow \mathcal{G}$ is a natural transformation of functors. That is, for each open set U , we have a group homomorphism $\phi(U) : \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ such that whenever $U \hookrightarrow V$ the following diagram commutes

$$\begin{array}{ccc} \mathcal{F}(V) & \xrightarrow{\phi(V)} & \mathcal{G}(V) \\ \text{res}_{V,U}^{\mathcal{F}} \downarrow & & \downarrow \text{res}_{V,U}^{\mathcal{G}} \\ \mathcal{F}(U) & \xrightarrow{\phi(U)} & \mathcal{G}(U) \end{array}$$

Sheafification

The functor $i : \text{Sh}(X) \hookrightarrow \text{PSh}(X)$ has a left adjoint

Sheafification

The functor $i : \mathbf{Sh}(X) \hookrightarrow \mathbf{PSh}(X)$ has a left adjoint

$$a : \mathbf{PSh}(X) \rightarrow \mathbf{Sh}(X)$$

called the sheafification functor,

Sheafification

The functor $i : \text{Sh}(X) \hookrightarrow \text{PSh}(X)$ has a left adjoint

$$a : \text{PSh}(X) \rightarrow \text{Sh}(X)$$

called the sheafification functor, where for a presheaf \mathcal{F} , $a(\mathcal{F})$ or \mathcal{F}^{sh} is called the associated sheaf of \mathcal{F} or its sheafification.

Sheafification

The functor $i : \text{Sh}(X) \hookrightarrow \text{PSh}(X)$ has a left adjoint

$$a : \text{PSh}(X) \rightarrow \text{Sh}(X)$$

called the sheafification functor, where for a presheaf \mathcal{F} , $a(\mathcal{F})$ or \mathcal{F}^{sh} is called the associated sheaf of \mathcal{F} or its sheafification. That is, we have a bijection, natural in its entries

$$\text{Mor}_{\text{PSh}(X)}(\mathcal{F}, i(\mathcal{G})) \cong \text{Mor}_{\text{Sh}(X)}(a(\mathcal{F}), \mathcal{G}).$$

Sheafification

The functor $i : \text{Sh}(X) \hookrightarrow \text{PSh}(X)$ has a left adjoint

$$a : \text{PSh}(X) \rightarrow \text{Sh}(X)$$

called the sheafification functor, where for a presheaf \mathcal{F} , $a(\mathcal{F})$ or \mathcal{F}^{sh} is called the associated sheaf of \mathcal{F} or its sheafification. That is, we have a bijection, natural in its entries

$$\text{Mor}_{\text{PSh}(X)}(\mathcal{F}, i(\mathcal{G})) \cong \text{Mor}_{\text{Sh}(X)}(a(\mathcal{F}), \mathcal{G}).$$

Example

For any abelian group A , the constant sheaf \underline{A} is the sheafification of the constant presheaf $\underline{A}_{\text{pre}}$.

Sheaf Cohomology

× × ×

Kernels and Cokernels

Fix a topological space X and our presheaves and sheaves on X are all presheaves and sheaves of abelian groups.

Category of Sheaves is Abelian

Kernels and Cokernels

Fix a topological space X and our presheaves and sheaves on X are all presheaves and sheaves of abelian groups.

Given a morphism of sheaves on X , say $\phi : \mathcal{F} \rightarrow \mathcal{G}$, one can define the presheaves $\ker \phi$, $\operatorname{coker} \phi$ and $\operatorname{im} \phi$ in the obvious manner.

Kernels and Cokernels

Fix a topological space X and our presheaves and sheaves on X are all presheaves and sheaves of abelian groups.

Given a morphism of sheaves on X , say $\phi : \mathcal{F} \rightarrow \mathcal{G}$, one can define the presheaves $\ker \phi$, $\operatorname{coker} \phi$ and $\operatorname{im} \phi$ in the obvious manner. That is,

$$(\ker \phi)(U) := \ker(\phi(U)), \quad (\operatorname{coker} \phi)(U) = \operatorname{coker} \phi(U), \quad (\operatorname{im} \phi)(U) = \operatorname{im} \phi(U).$$

Kernels and Cokernels

Fix a topological space X and our presheaves and sheaves on X are all presheaves and sheaves of abelian groups.

Given a morphism of sheaves on X , say $\phi : \mathcal{F} \rightarrow \mathcal{G}$, one can define the presheaves $\ker \phi$, $\operatorname{coker} \phi$ and $\operatorname{im} \phi$ in the obvious manner. That is,

$$(\ker \phi)(U) := \ker(\phi(U)), \quad (\operatorname{coker} \phi)(U) = \operatorname{coker} \phi(U), \quad (\operatorname{im} \phi)(U) = \operatorname{im} \phi(U).$$

Out of these only $\ker \phi$ is a sheaf, while $\operatorname{coker} \phi$ (resp. $\operatorname{im} \phi$) fail to be.

Kernels and Cokernels

Fix a topological space X and our presheaves and sheaves on X are all presheaves and sheaves of abelian groups.

Given a morphism of sheaves on X , say $\phi : \mathcal{F} \rightarrow \mathcal{G}$, one can define the presheaves $\ker \phi$, $\operatorname{coker} \phi$ and $\operatorname{im} \phi$ in the obvious manner. That is,

$$(\ker \phi)(U) := \ker(\phi(U)), \quad (\operatorname{coker} \phi)(U) = \operatorname{coker} \phi(U), \quad (\operatorname{im} \phi)(U) = \operatorname{im} \phi(U).$$

Out of these only $\ker \phi$ is a sheaf, while $\operatorname{coker} \phi$ (resp. $\operatorname{im} \phi$) fail to be. We simply rectify this by considering $(\operatorname{coker} \phi)^{\operatorname{sh}}$ (resp. $(\operatorname{im} \phi)^{\operatorname{sh}}$) and this does turn out to be the cokernel (resp. image) of ϕ in $\operatorname{Sh}(X)$.

Category of Sheaves is Abelian

Kernels and Cokernels

Fix a topological space X and our presheaves and sheaves on X are all presheaves and sheaves of abelian groups.

Given a morphism of sheaves on X , say $\phi : \mathcal{F} \rightarrow \mathcal{G}$, one can define the presheaves $\ker \phi$, $\operatorname{coker} \phi$ and $\operatorname{im} \phi$ in the obvious manner. That is,

$$(\ker \phi)(U) := \ker(\phi(U)), \quad (\operatorname{coker} \phi)(U) = \operatorname{coker} \phi(U), \quad (\operatorname{im} \phi)(U) = \operatorname{im} \phi(U).$$

Out of these only $\ker \phi$ is a sheaf, while $\operatorname{coker} \phi$ (resp. $\operatorname{im} \phi$) fail to be. We simply rectify this by considering $(\operatorname{coker} \phi)^{\operatorname{sh}}$ (resp. $(\operatorname{im} \phi)^{\operatorname{sh}}$) and this does turn out to be the cokernel (resp. image) of ϕ in $\operatorname{Sh}(X)$.

Category of Sheaves is Abelian

The category $\operatorname{Sh}(X)$ is an abelian category with enough injectives.

Category of Sheaves is Abelian

Kernels and Cokernels

Fix a topological space X and our presheaves and sheaves on X are all presheaves and sheaves of abelian groups.

Given a morphism of sheaves on X , say $\phi : \mathcal{F} \rightarrow \mathcal{G}$, one can define the presheaves $\ker \phi$, $\operatorname{coker} \phi$ and $\operatorname{im} \phi$ in the obvious manner. That is,

$$(\ker \phi)(U) := \ker(\phi(U)), \quad (\operatorname{coker} \phi)(U) = \operatorname{coker} \phi(U), \quad (\operatorname{im} \phi)(U) = \operatorname{im} \phi(U).$$

Out of these only $\ker \phi$ is a sheaf, while $\operatorname{coker} \phi$ (resp. $\operatorname{im} \phi$) fail to be. We simply rectify this by considering $(\operatorname{coker} \phi)^{\operatorname{sh}}$ (resp. $(\operatorname{im} \phi)^{\operatorname{sh}}$) and this does turn out to be the cokernel (resp. image) of ϕ in $\operatorname{Sh}(X)$.

Category of Sheaves is Abelian

The category $\operatorname{Sh}(X)$ is an abelian category with enough injectives. The latter is a consequence of Ab having enough injectives.

Global Sections Functor

We can consider the *global section functor*

$$\Gamma(X, -) : \mathbf{Sh}(X) \rightarrow \mathbf{Ab}$$

Global Sections Functor

We can consider the *global section functor*

$$\Gamma(X, -) : \mathbf{Sh}(X) \rightarrow \mathbf{Ab}$$

This functor is left exact.

Global Sections Functor

We can consider the *global section functor*

$$\Gamma(X, -) : \mathbf{Sh}(X) \rightarrow \mathbf{Ab}$$

This functor is left exact. That is, given an exact sequence in $\mathbf{Sh}(X)$

$$0 \longrightarrow \mathcal{F} \xrightarrow{\phi} \mathcal{G} \xrightarrow{\psi} \mathcal{H} \longrightarrow 0$$

Global Sections Functor

We can consider the *global section functor*

$$\Gamma(X, -) : \mathbf{Sh}(X) \rightarrow \mathbf{Ab}$$

This functor is left exact. That is, given an exact sequence in $\mathbf{Sh}(X)$

$$0 \longrightarrow \mathcal{F} \xrightarrow{\phi} \mathcal{G} \xrightarrow{\psi} \mathcal{H} \longrightarrow 0$$

the following induced sequence is exact in \mathbf{Ab}

$$0 \longrightarrow \Gamma(X, \mathcal{F}) \xrightarrow{\phi(X)} \Gamma(X, \mathcal{G}) \xrightarrow{\psi(X)} \Gamma(X, \mathcal{H})$$

Global Sections Functor

We can consider the *global section functor*

$$\Gamma(X, -) : \mathbf{Sh}(X) \rightarrow \mathbf{Ab}$$

This functor is left exact. That is, given an exact sequence in $\mathbf{Sh}(X)$

$$0 \longrightarrow \mathcal{F} \xrightarrow{\phi} \mathcal{G} \xrightarrow{\psi} \mathcal{H} \longrightarrow 0$$

the following induced sequence is exact in \mathbf{Ab}

$$0 \longrightarrow \Gamma(X, \mathcal{F}) \xrightarrow{\phi(X)} \Gamma(X, \mathcal{G}) \xrightarrow{\psi(X)} \Gamma(X, \mathcal{H})$$

The takeaway being that $\psi(X)$ is not necessarily surjective.

The standard example that illustrates this is the *exponential exact sequence*.

Exponential Exact Sequence

Let $X = \mathbb{C}^*$ (the punctured complex plane) with the standard topology, and define \mathcal{O}_X to be the sheaf of holomorphic functions. Also let \mathcal{O}_X^* to be the sheaf of invertible (nowhere zero) holomorphic functions (this is a sheaf of abelian groups under multiplication).

Exponential Exact Sequence

Let $X = \mathbb{C}^*$ (the punctured complex plane) with the standard topology, and define \mathcal{O}_X to be the sheaf of holomorphic functions. Also let \mathcal{O}_X^* to be the sheaf of invertible (nowhere zero) holomorphic functions (this is a sheaf of abelian groups under multiplication).

Then it's a fact that the following is an exact sequence of sheaves

$$0 \longrightarrow \underline{\mathbb{Z}} \xrightarrow{2\pi i} \mathcal{O}_X \xrightarrow{\exp} \mathcal{O}_X^* \longrightarrow 1$$

Exponential Exact Sequence

Let $X = \mathbb{C}^*$ (the punctured complex plane) with the standard topology, and define \mathcal{O}_X to be the sheaf of holomorphic functions. Also let \mathcal{O}_X^* to be the sheaf of invertible (nowhere zero) holomorphic functions (this is a sheaf of abelian groups under multiplication).

Then it's a fact that the following is an exact sequence of sheaves

$$0 \longrightarrow \underline{\mathbb{Z}} \xrightarrow{2\pi i} \mathcal{O}_X \xrightarrow{\exp} \mathcal{O}_X^* \longrightarrow 1$$

But

$$\exp : \Gamma(X, \mathcal{O}_X) \rightarrow \Gamma(X, \mathcal{O}_X^*) = \Gamma(X, \mathcal{O}_X)^*$$

is not surjective since there does not exist a global complex logarithm.

Sheaf Cohomology

Since the *global section functor* $\Gamma(X, -)$ is left exact and $\text{Sh}(X)$ has enough injectives,

Sheaf Cohomology

Since the *global section functor* $\Gamma(X, -)$ is left exact and $\text{Sh}(X)$ has enough injectives, we define the *sheaf cohomology* of X with coefficients in \mathcal{F} as

$$H^i(X, \mathcal{F}) := R^i\Gamma(X, \mathcal{F}),$$

Sheaf Cohomology

Since the *global section functor* $\Gamma(X, -)$ is left exact and $\text{Sh}(X)$ has enough injectives, we define the *sheaf cohomology* of X with coefficients in \mathcal{F} as

$$H^i(X, \mathcal{F}) := R^i\Gamma(X, \mathcal{F}),$$

that is, the right derived functor of $\Gamma(X, -)$.

Sheaf Cohomology

Since the *global section functor* $\Gamma(X, -)$ is left exact and $\text{Sh}(X)$ has enough injectives, we define the *sheaf cohomology* of X with coefficients in \mathcal{F} as

$$H^i(X, \mathcal{F}) := R^i\Gamma(X, \mathcal{F}),$$

that is, the right derived functor of $\Gamma(X, -)$. Therefore, given any short exact sequence of sheaves

$$0 \longrightarrow \mathcal{F} \xrightarrow{\phi} \mathcal{G} \xrightarrow{\psi} \mathcal{H} \longrightarrow 0$$

Sheaf Cohomology

Since the *global section functor* $\Gamma(X, -)$ is left exact and $\mathbf{Sh}(X)$ has enough injectives, we define the *sheaf cohomology* of X with coefficients in \mathcal{F} as

$$H^i(X, \mathcal{F}) := R^i\Gamma(X, \mathcal{F}),$$

that is, the right derived functor of $\Gamma(X, -)$. Therefore, given any short exact sequence of sheaves

$$0 \longrightarrow \mathcal{F} \xrightarrow{\phi} \mathcal{G} \xrightarrow{\psi} \mathcal{H} \longrightarrow 0$$

we obtain a long exact sequence of abelian groups

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Gamma(X, \mathcal{F}) & \longrightarrow & \Gamma(X, \mathcal{G}) & \longrightarrow & \Gamma(X, \mathcal{H}) \\ & & & & & \searrow & \\ & & & & & H^1(X, \mathcal{F}) & \longrightarrow H^1(X, \mathcal{G}) \longrightarrow H^1(X, \mathcal{H}) \\ & & & & & & \searrow \\ & & & & & & H^2(X, \mathcal{F}) \longrightarrow \dots \end{array}$$

Exponential Exact Sequence

Let X be a complex manifold, we again consider as before the exponential exact sequence

$$0 \longrightarrow \underline{\mathbb{Z}} \xrightarrow{2\pi i} \mathcal{O}_X \xrightarrow{\exp} \mathcal{O}_X^* \longrightarrow 1$$


Exponential Exact Sequence

Let X be a complex manifold, we again consider as before the exponential exact sequence

$$0 \longrightarrow \underline{\mathbb{Z}} \xrightarrow{2\pi i} \mathcal{O}_X \xrightarrow{\exp} \mathcal{O}_X^* \longrightarrow 1$$

which then gives us the long exact sequence in cohomology

$$0 \longrightarrow \Gamma(X, \underline{\mathbb{Z}}) \longrightarrow \Gamma(X, \mathcal{O}_X) \longrightarrow \Gamma(X, \mathcal{O}_X^*)$$


$$H^1(X, \underline{\mathbb{Z}}) \longrightarrow H^1(X, \mathcal{O}_X) \longrightarrow H^1(X, \mathcal{O}_X^*)$$


$$H^2(X, \underline{\mathbb{Z}}) \longrightarrow \dots$$


Exponential Exact Sequence

Let X be a complex manifold, we again consider as before the exponential exact sequence

$$0 \longrightarrow \underline{\mathbb{Z}} \xrightarrow{2\pi i} \mathcal{O}_X \xrightarrow{\exp} \mathcal{O}_X^* \longrightarrow 1$$

which then gives us the long exact sequence in cohomology

$$0 \longrightarrow \Gamma(X, \underline{\mathbb{Z}}) \longrightarrow \Gamma(X, \mathcal{O}_X) \longrightarrow \Gamma(X, \mathcal{O}_X^*)$$


$$H^1(X, \underline{\mathbb{Z}}) \longrightarrow H^1(X, \mathcal{O}_X) \longrightarrow H^1(X, \mathcal{O}_X^*)$$


$$H^2(X, \underline{\mathbb{Z}}) \longrightarrow \dots$$

A comparison theorem tells us that $H^i(X, \underline{\mathbb{Z}}) \cong \underline{H_{\text{sing}}^i(X, \mathbb{Z})}$, the singular cohomology group of X ;

Exponential Exact Sequence

Let X be a complex manifold, we again consider as before the exponential exact sequence

$$0 \longrightarrow \underline{\mathbb{Z}} \xrightarrow{2\pi i} \mathcal{O}_X \xrightarrow{\exp} \mathcal{O}_X^* \longrightarrow 1$$

which then gives us the long exact sequence in cohomology

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Gamma(X, \underline{\mathbb{Z}}) & \longrightarrow & \Gamma(X, \mathcal{O}_X) & \longrightarrow & \Gamma(X, \mathcal{O}_X^*)^* \\ & & & & & & \downarrow \\ & & & & & & H^1(X, \underline{\mathbb{Z}}) \longrightarrow H^1(X, \mathcal{O}_X) \longrightarrow H^1(X, \mathcal{O}_X^*) \\ & & & & & & \downarrow \\ & & & & & & H^2(X, \underline{\mathbb{Z}}) \longrightarrow \dots \end{array}$$

A comparison theorem tells us that $H^i(X, \underline{\mathbb{Z}}) \cong H_{\text{sing}}^i(X, \mathbb{Z})$, the singular cohomology group of X ; in particular,

$$H^1(X, \underline{\mathbb{Z}}) \cong H_{\text{sing}}^1(X, \mathbb{Z}) = \text{Hom}(H_1^{\text{sing}}(X, \mathbb{Z}), \mathbb{Z}) \cong \text{Hom}(\pi_1(X)^{\text{ab}}, \mathbb{Z}).$$

(...)

Exponential Exact Sequence

(...) Another standard fact is that $H^1(X, \mathcal{O}_X^*) \cong \text{Pic}(X)$, the Picard group of X which is the group of isomorphism classes of complex line bundles on X .

Exponential Exact Sequence

(...) Another standard fact is that $H^1(X, \mathcal{O}_X^*) \cong \text{Pic}(X)$, the Picard group of X which is the group of isomorphism classes of complex line bundles on X .

Let's concentrate on the connecting morphisms

$$w : H^0(X, \mathcal{O}_X^*) \rightarrow H^1(X, \underline{\mathbb{Z}})$$

$$c : H^1(X, \mathcal{O}_X^*) \rightarrow H^2(X, \underline{\mathbb{Z}})$$

Exponential Exact Sequence

(...) Another standard fact is that $H^1(X, \mathcal{O}_X^*) \cong \text{Pic}(X)$, the Picard group of X which is the group of isomorphism classes of complex line bundles on X .

Let's concentrate on the connecting morphisms

$$w : H^0(X, \mathcal{O}_X^*) \rightarrow H^1(X, \underline{\mathbb{Z}})$$

$$c : H^1(X, \mathcal{O}_X^*) \rightarrow H^2(X, \underline{\mathbb{Z}})$$

Under the above identifications, w can be understood as taking a winding number of a global section of \mathcal{O}_X^* ,

Exponential Exact Sequence

(...) Another standard fact is that $H^1(X, \mathcal{O}_X^*) \cong \text{Pic}(X)$, the Picard group of X which is the group of isomorphism classes of complex line bundles on X .

Let's concentrate on the connecting morphisms

$$w : H^0(X, \mathcal{O}_X^*) \rightarrow H^1(X, \underline{\mathbb{Z}})$$

$$c : H^1(X, \mathcal{O}_X^*) \rightarrow H^2(X, \underline{\mathbb{Z}})$$

Under the above identifications, w can be understood as taking a winding number of a global section of \mathcal{O}_X^* , detecting the obstruction to taking a global logarithm of a non-vanishing holomorphic function.

Exponential Exact Sequence

(...) Another standard fact is that $H^1(X, \mathcal{O}_X^*) \cong \text{Pic}(X)$, the Picard group of X which is the group of isomorphism classes of complex line bundles on X .

Let's concentrate on the connecting morphisms

$$w : H^0(X, \mathcal{O}_X^*) \rightarrow H^1(X, \underline{\mathbb{Z}})$$

$$c : H^1(X, \mathcal{O}_X^*) \rightarrow H^2(X, \underline{\mathbb{Z}})$$

Under the above identifications, w can be understood as taking a winding number of a global section of \mathcal{O}_X^* , detecting the obstruction to taking a global logarithm of a non-vanishing holomorphic function. That is,

$$w(g) := \left(\gamma \mapsto \int_{\gamma} \frac{dg}{g} \right)$$

Exponential Exact Sequence

(...) Another standard fact is that $H^1(X, \mathcal{O}_X^*) \cong \text{Pic}(X)$, the Picard group of X which is the group of isomorphism classes of complex line bundles on X .

Let's concentrate on the connecting morphisms

$$w : H^0(X, \mathcal{O}_X^*) \rightarrow H^1(X, \underline{\mathbb{Z}})$$

$$c : H^1(X, \mathcal{O}_X^*) \rightarrow H^2(X, \underline{\mathbb{Z}})$$

Under the above identifications, w can be understood as taking a winding number of a global section of \mathcal{O}_X^* , detecting the obstruction to taking a global logarithm of a non-vanishing holomorphic function. That is,

$$w(g) := \left(\gamma \mapsto \int_{\gamma} \frac{dg}{g} \right)$$

While c associates to each (isomorphism class of) line bundle \mathcal{L} its first Chern class $c_1(\mathcal{L})$.

(...)

Exponential Exact Sequence

(...) Furthermore, suppose $X = C$ was a smooth projective curve, then because of dimension reasons we have $H^2(X, \mathcal{O}_C) = 0$

Exponential Exact Sequence

(...) Furthermore, suppose $X = C$ was a smooth projective curve, then because of dimension reasons we have $H^2(X, \mathcal{O}_C) = 0$ and the first three terms of the long exact sequence are simply

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{C} \rightarrow \mathbb{C}^* \rightarrow 1$$

Exponential Exact Sequence

(...) Furthermore, suppose $X = C$ was a smooth projective curve, then because of dimension reasons we have $H^2(X, \mathcal{O}_C) = 0$ and the first three terms of the long exact sequence are simply

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{C} \rightarrow \mathbb{C}^* \rightarrow 1$$

This gives us the exact sequence

$$0 \rightarrow H_{\text{sing}}^1(C, \mathbb{Z}) \rightarrow H^1(C, \mathcal{O}_C) \rightarrow \text{Pic}(C) \rightarrow H_{\text{sing}}^2(C, \mathbb{Z}) \rightarrow 0$$

Exponential Exact Sequence

(...) Furthermore, suppose $X = C$ was a smooth projective curve, then because of dimension reasons we have $H^2(X, \mathcal{O}_C) = 0$ and the first three terms of the long exact sequence are simply

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{C} \rightarrow \mathbb{C}^* \rightarrow 1$$

This gives us the exact sequence

$$0 \rightarrow H_{\text{sing}}^1(C, \mathbb{Z}) \rightarrow H^1(C, \mathcal{O}_C) \rightarrow \text{Pic}(C) \rightarrow H_{\text{sing}}^2(C, \mathbb{Z}) \rightarrow 0$$

We have $H^1(C, \mathcal{O}_C) \cong \mathbb{C}^g$, where g is the genus of C , by Serre duality;

Exponential Exact Sequence

(...) Furthermore, suppose $X = C$ was a smooth projective curve, then because of dimension reasons we have $H^2(X, \mathcal{O}_C) = 0$ and the first three terms of the long exact sequence are simply

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{C} \rightarrow \mathbb{C}^* \rightarrow 1$$

This gives us the exact sequence

$$0 \rightarrow H_{\text{sing}}^1(C, \mathbb{Z}) \rightarrow H^1(C, \mathcal{O}_C) \rightarrow \text{Pic}(C) \rightarrow H_{\text{sing}}^2(C, \mathbb{Z}) \rightarrow 0$$

We have $H^1(C, \mathcal{O}_C) \cong \mathbb{C}^g$, where g is the genus of C , by Serre duality; and $H_{\text{sing}}^1(C, \mathbb{Z}) \subseteq H^1(C, \mathcal{O}_C)$ is a lattice.

Exponential Exact Sequence

(...) Furthermore, suppose $X = C$ was a smooth projective curve, then because of dimension reasons we have $H^2(X, \mathcal{O}_C) = 0$ and the first three terms of the long exact sequence are simply

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{C} \rightarrow \mathbb{C}^* \rightarrow 1$$

This gives us the exact sequence

$$0 \rightarrow H_{\text{sing}}^1(C, \mathbb{Z}) \rightarrow H^1(C, \mathcal{O}_C) \rightarrow \text{Pic}(C) \rightarrow H_{\text{sing}}^2(C, \mathbb{Z}) \rightarrow 0$$

We have $H^1(C, \mathcal{O}_C) \cong \mathbb{C}^g$, where g is the genus of C , by Serre duality; and $H_{\text{sing}}^1(C, \mathbb{Z}) \subseteq H^1(C, \mathcal{O}_C)$ is a lattice.

Then $\text{Jac}(C) := H^1(C, \mathcal{O}_C) / H_{\text{sing}}^1(C, \mathbb{Z})$ is a complex torus of dimension g and is called the *Jacobian of C* .

Exponential Exact Sequence

(...) Furthermore, suppose $X = C$ was a smooth projective curve, then because of dimension reasons we have $H^2(X, \mathcal{O}_C) = 0$ and the first three terms of the long exact sequence are simply

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{C} \rightarrow \mathbb{C}^* \rightarrow 1$$

This gives us the exact sequence

$$0 \rightarrow H_{\text{sing}}^1(C, \mathbb{Z}) \rightarrow H^1(C, \mathcal{O}_C) \rightarrow \text{Pic}(C) \rightarrow H_{\text{sing}}^2(C, \mathbb{Z}) \rightarrow 0$$

We have $H^1(C, \mathcal{O}_C) \cong \mathbb{C}^g$, where g is the genus of C , by Serre duality; and $H_{\text{sing}}^1(C, \mathbb{Z}) \subseteq H^1(C, \mathcal{O}_C)$ is a lattice.

Then $\text{Jac}(C) := H^1(C, \mathcal{O}_C) / H_{\text{sing}}^1(C, \mathbb{Z})$ is a complex torus of dimension g and is called the *Jacobian of C* .

By exactness, $\text{Jac}(C)$ can be identified with the kernel of $c_1 : \text{Pic}(C) \rightarrow H_{\text{sing}}^2(C, \mathbb{Z})$, equivalently the degree map.

Exponential Exact Sequence

(...) Furthermore, suppose $X = C$ was a smooth projective curve, then because of dimension reasons we have $H^2(X, \mathcal{O}_C) = 0$ and the first three terms of the long exact sequence are simply

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{C} \rightarrow \mathbb{C}^* \rightarrow 1$$

This gives us the exact sequence

$$0 \rightarrow H_{\text{sing}}^1(C, \mathbb{Z}) \rightarrow H^1(C, \mathcal{O}_C) \rightarrow \text{Pic}(C) \rightarrow H_{\text{sing}}^2(C, \mathbb{Z}) \rightarrow 0$$

We have $H^1(C, \mathcal{O}_C) \cong \mathbb{C}^g$, where g is the genus of C , by Serre duality; and $H_{\text{sing}}^1(C, \mathbb{Z}) \subseteq H^1(C, \mathcal{O}_C)$ is a lattice.

Then $\text{Jac}(C) := H^1(C, \mathcal{O}_C) / H_{\text{sing}}^1(C, \mathbb{Z})$ is a complex torus of dimension g and is called the *Jacobian of C* .

By exactness, $\text{Jac}(C)$ can be identified with the kernel of $c_1 : \text{Pic}(C) \rightarrow H_{\text{sing}}^2(C, \mathbb{Z})$, equivalently the degree map. That is, it's the group of degree 0 line bundles,

Exponential Exact Sequence

(...) Furthermore, suppose $X = C$ was a smooth projective curve, then because of dimension reasons we have $H^2(X, \mathcal{O}_C) = 0$ and the first three terms of the long exact sequence are simply

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{C} \rightarrow \mathbb{C}^* \rightarrow 1$$

This gives us the exact sequence

$$0 \rightarrow H_{\text{sing}}^1(C, \mathbb{Z}) \rightarrow H^1(C, \mathcal{O}_C) \rightarrow \text{Pic}(C) \rightarrow H_{\text{sing}}^2(C, \mathbb{Z}) \rightarrow 0$$

We have $H^1(C, \mathcal{O}_C) \cong \mathbb{C}^g$, where g is the genus of C , by Serre duality; and $H_{\text{sing}}^1(C, \mathbb{Z}) \subseteq H^1(C, \mathcal{O}_C)$ is a lattice.

Then $\text{Jac}(C) := H^1(C, \mathcal{O}_C) / H_{\text{sing}}^1(C, \mathbb{Z})$ is a complex torus of dimension g and is called the *Jacobian of C* .

By exactness, $\text{Jac}(C)$ can be identified with the kernel of $c_1 : \text{Pic}(C) \rightarrow H_{\text{sing}}^2(C, \mathbb{Z})$, equivalently the degree map. That is, it's the group of degree 0 line bundles, which can be identified with the connected component of $\text{Pic}(C)$.

Sites

× × ×

Let's go back to Sheaves,

Let's go back to Sheaves, in our definition of sheaves what was really important was the notion of coverings and intersections.

Let's go back to Sheaves, in our definition of sheaves what was really important was the notion of coverings and intersections. With that observation we make the following general definition, where we replace the notion of intersections with the more general notion of fibered products and weaken the notion of unions to a notion of covers.

Let's go back to Sheaves, in our definition of sheaves what was really important was the notion of coverings and intersections. With that observation we make the following general definition, where we replace the notion of intersections with the more general notion of fibered products and weaken the notion of unions to a notion of covers.

Grothendieck Topology

Let \mathcal{C} be a category, a *Grothendieck topology* on \mathcal{C} is an assignment, for each object U of \mathcal{C} , of a set $\mathbf{cov}(U)$ of families of morphisms $\{U_i \rightarrow U\}_{i \in I}$ (called *coverings*) such that

Let's go back to Sheaves, in our definition of sheaves what was really important was the notion of coverings and intersections. With that observation we make the following general definition, where we replace the notion of intersections with the more general notion of fibered products and weaken the notion of unions to a notion of covers.

Grothendieck Topology

Let \mathcal{C} be a category, a *Grothendieck topology* on \mathcal{C} is an assignment, for each object U of \mathcal{C} , of a set $\mathbf{cov}(U)$ of families of morphisms $\{U_i \rightarrow U\}_{i \in I}$ (called *coverings*) such that


- if $\{U' \rightarrow U\}$ is an isomorphism in \mathcal{C} , then it's in $\mathbf{cov}(U)$;

Let's go back to Sheaves, in our definition of sheaves what was really important was the notion of coverings and intersections. With that observation we make the following general definition, where we replace the notion of intersections with the more general notion of fibered products and weaken the notion of unions to a notion of covers.

Grothendieck Topology

Let \mathcal{C} be a category, a *Grothendieck topology* on \mathcal{C} is an assignment, for each object U of \mathcal{C} , of a set $\mathbf{cov}(U)$ of families of morphisms $\{U_i \rightarrow U\}_{i \in I}$ (called *coverings*) such that

- if $\{U' \rightarrow U\}$ is an isomorphism in \mathcal{C} , then it's in $\mathbf{cov}(U)$;
- if $\{U_i \rightarrow U\}_i$ is in $\mathbf{cov}(U)$ and $V \rightarrow U$ is a morphism in \mathcal{C} , then all fibered products $V \times_U U_i$ exist and $\{V \times_U U_i \rightarrow V\}_i$ is in $\mathbf{cov}(V)$.



Let's go back to Sheaves, in our definition of sheaves what was really important was the notion of coverings and intersections. With that observation we make the following general definition, where we replace the notion of intersections with the more general notion of fibered products and weaken the notion of unions to a notion of covers.

Grothendieck Topology

Let \mathcal{C} be a category, a *Grothendieck topology* on \mathcal{C} is an assignment, for each object U of \mathcal{C} , of a set $\mathbf{cov}(U)$ of families of morphisms $\{U_i \rightarrow U\}_{i \in I}$ (called *coverings*) such that

- if $\{U' \rightarrow U\}$ is an isomorphism in \mathcal{C} , then it's in $\mathbf{cov}(U)$;
- if $\{U_i \rightarrow U\}_i$ is in $\mathbf{cov}(U)$ and $V \rightarrow U$ is a morphism in \mathcal{C} , then all fibered products $V \times_U U_i$ exist and $\{V \times_U U_i \rightarrow V\}_i$ is in $\mathbf{cov}(V)$.
- if $\{U_i \rightarrow U\}_i$ is in $\mathbf{cov}(U)$ and $\{V_{ij} \rightarrow U_i\}_j$ is in $\mathbf{cov}(U_i)$, then $\{V_{ij} \rightarrow U_i \rightarrow U\}_{i,j}$, obtained by composition, is in $\mathbf{cov}(U)$.

$$\boxed{V_{ij} \rightarrow U_i \rightarrow U} \quad \text{and} \quad \textcircled{V_{ij} \rightarrow U}$$

Let's go back to Sheaves, in our definition of sheaves what was really important was the notion of coverings and intersections. With that observation we make the following general definition, where we replace the notion of intersections with the more general notion of fibered products and weaken the notion of unions to a notion of covers.

Grothendieck Topology

Let \mathcal{C} be a category, a *Grothendieck topology* on \mathcal{C} is an assignment, for each object U of \mathcal{C} , of a set $\mathbf{cov}(U)$ of families of morphisms $\{U_i \rightarrow U\}_{i \in I}$ (called *coverings*) such that

- if $\{U' \rightarrow U\}$ is an isomorphism in \mathcal{C} , then it's in $\mathbf{cov}(U)$;
- if $\{U_i \rightarrow U\}_i$ is in $\mathbf{cov}(U)$ and $V \rightarrow U$ is a morphism in \mathcal{C} , then all fibered products $V \times_U U_i$ exist and $\{V \times_U U_i \rightarrow V\}_i$ is in $\mathbf{cov}(V)$.
- if $\{U_i \rightarrow U\}_i$ is in $\mathbf{cov}(U)$ and $\{V_{ij} \rightarrow U_i\}_j$ is in $\mathbf{cov}(U_i)$, then $\{V_{ij} \rightarrow U_i \rightarrow U\}_{i,j}$, obtained by composition, is in $\mathbf{cov}(U)$.

A category equipped with a Grothendieck topology is called a *site*.

Let's go back to Sheaves, in our definition of sheaves what was really important was the notion of coverings and intersections. With that observation we make the following general definition, where we replace the notion of intersections with the more general notion of fibered products and weaken the notion of unions to a notion of covers.

Grothendieck Topology

Let \mathcal{C} be a category, a *Grothendieck topology* on \mathcal{C} is an assignment, for each object U of \mathcal{C} , of a set $\mathbf{cov}(U)$ of families of morphisms $\{U_i \rightarrow U\}_{i \in I}$ (called *coverings*) such that

- if $\{U' \rightarrow U\}$ is an isomorphism in \mathcal{C} , then it's in $\mathbf{cov}(U)$;
- if $\{U_i \rightarrow U\}_i$ is in $\mathbf{cov}(U)$ and $V \rightarrow U$ is a morphism in \mathcal{C} , then all fibered products $V \times_U U_i$ exist and $\{V \times_U U_i \rightarrow V\}_i$ is in $\mathbf{cov}(V)$.
- if $\{U_i \rightarrow U\}_i$ is in $\mathbf{cov}(U)$ and $\{V_{ij} \rightarrow U_i\}_j$ is in $\mathbf{cov}(U_i)$, then $\{V_{ij} \rightarrow U_i \rightarrow U\}_{i,j}$, obtained by composition, is in $\mathbf{cov}(U)$.

A category equipped with a Grothendieck topology is called a *site*.

So, given a site \mathcal{T} , we have the underlying category denoted as $\mathbf{Cat}(\mathcal{T})$ and a collection of covers for each object which we denote $\mathbf{Cov}(\mathcal{T})$.

Comma Topology

Given an object X of $\mathcal{C} = \text{Cat}(\mathcal{T})$, for a site \mathcal{T} , we can consider the comma category \mathcal{C}/X .

Comma Topology

Given an object X of $\mathcal{C} = \text{Cat}(\mathcal{T})$, for a site \mathcal{T} , we can consider the comma category \mathcal{C}/X . We can describe an induced Grothendieck topology on it, called the *comma topology*, to obtain the site that we denote $X_{\mathcal{T}}$ as follows:

Comma Topology

Given an object X of $\mathcal{C} = \text{Cat}(\mathcal{T})$, for a site \mathcal{T} , we can consider the comma category \mathcal{C}/X . We can describe an induced Grothendieck topology on it, called the *comma topology*, to obtain the site that we denote $X_{\mathcal{T}}$ as follows:

- Recall that objects of \mathcal{C}/X are morphisms $U \rightarrow X$,

Comma Topology

Given an object X of $\mathcal{C} = \text{Cat}(\mathcal{T})$, for a site \mathcal{T} , we can consider the comma category \mathcal{C}/X . We can describe an induced Grothendieck topology on it, called the *comma topology*, to obtain the site that we denote $X_{\mathcal{T}}$ as follows:

- Recall that objects of \mathcal{C}/X are morphisms $U \rightarrow X$, and that given another object $V \rightarrow X$, a morphism $(U \rightarrow X) \rightarrow (V \rightarrow X)$

Comma Topology

Given an object X of $\mathcal{C} = \text{Cat}(\mathcal{T})$, for a site \mathcal{T} , we can consider the comma category \mathcal{C}/X . We can describe an induced Grothendieck topology on it, called the *comma topology*, to obtain the site that we denote $X_{\mathcal{T}}$ as follows:

- Recall that objects of \mathcal{C}/X are morphisms $U \rightarrow X$, and that given another object $V \rightarrow X$, a morphism $(U \rightarrow X) \rightarrow (V \rightarrow X)$ is a morphism $\phi : U \rightarrow V$ in \mathcal{C} such that the diagram

$$\begin{array}{ccc} U & \xrightarrow{\phi} & V \\ & \searrow & \swarrow \\ & X & \end{array}$$

commutes.

Comma Topology

Given an object X of $\mathcal{C} = \text{Cat}(\mathcal{T})$, for a site \mathcal{T} , we can consider the comma category \mathcal{C}/X . We can describe an induced Grothendieck topology on it, called the *comma topology*, to obtain the site that we denote $X_{\mathcal{T}}$ as follows:

- Recall that objects of \mathcal{C}/X are morphisms $U \rightarrow X$, and that given another object $V \rightarrow X$, a morphism $(U \rightarrow X) \rightarrow (V \rightarrow X)$ is a morphism $\phi : U \rightarrow V$ in \mathcal{C} such that the diagram

$$\begin{array}{ccc} U & \xrightarrow{\phi} & V \\ & \searrow & \swarrow \\ & X & \end{array}$$

commutes.

Note that $(\text{id}_X : X \rightarrow X)$ is the terminal object,

Comma Topology

Given an object X of $\mathcal{C} = \text{Cat}(\mathcal{T})$, for a site \mathcal{T} , we can consider the comma category \mathcal{C}/X . We can describe an induced Grothendieck topology on it, called the *comma topology*, to obtain the site that we denote $X_{\mathcal{T}}$ as follows:

- Recall that objects of \mathcal{C}/X are morphisms $U \rightarrow X$, and that given another object $V \rightarrow X$, a morphism $(U \rightarrow X) \rightarrow (V \rightarrow X)$ is a morphism $\phi : U \rightarrow V$ in \mathcal{C} such that the diagram

$$\begin{array}{ccc} U & \xrightarrow{\phi} & V \\ & \searrow & \swarrow \\ & X & \end{array}$$

commutes.

Note that $(\text{id}_X : X \rightarrow X)$ is the terminal object, one thinks of this as " X is the terminal object".

Comma Topology

Given an object X of $\mathcal{C} = \text{Cat}(\mathcal{T})$, for a site \mathcal{T} , we can consider the comma category \mathcal{C}/X . We can describe an induced Grothendieck topology on it, called the *comma topology*, to obtain the site that we denote $X_{\mathcal{T}}$ as follows:


- Recall that objects of \mathcal{C}/X are morphisms $U \rightarrow X$, and that given another object $V \rightarrow X$, a morphism $(U \rightarrow X) \rightarrow (V \rightarrow X)$ is a morphism $\phi : U \rightarrow V$ in \mathcal{C} such that the diagram

$$\begin{array}{ccc} U & \xrightarrow{\phi} & V \\ & \searrow & \swarrow \\ & X & \end{array}$$

commutes.

Note that $(\text{id}_X : X \rightarrow X)$ is the terminal object, one thinks of this as " X is the terminal object".

- For any object $U \rightarrow X$, the covering $\mathbf{cov}(U \rightarrow X)$ consists of families of morphisms $\{(U_i \rightarrow X) \rightarrow (U \rightarrow X)\}_i$ such that $\{U_i \rightarrow U\}$ is in $\mathbf{cov}(U)$ in $\text{Cov}(\mathcal{T})$.



Comma Topology

Given an object X of $\mathcal{C} = \text{Cat}(\mathcal{T})$, for a site \mathcal{T} , we can consider the comma category \mathcal{C}/X . We can describe an induced Grothendieck topology on it, called the *comma topology*, to obtain the site that we denote $X_{\mathcal{T}}$ as follows:

- Recall that objects of \mathcal{C}/X are morphisms $U \rightarrow X$, and that given another object $V \rightarrow X$, a morphism $(U \rightarrow X) \rightarrow (V \rightarrow X)$ is a morphism $\phi : U \rightarrow V$ in \mathcal{C} such that the diagram

$$\begin{array}{ccc} U & \xrightarrow{\phi} & V \\ & \searrow & \swarrow \\ & X & \end{array}$$

commutes.

Note that $(\text{id}_X : X \rightarrow X)$ is the terminal object, one thinks of this as " X is the terminal object".

- For any object $U \rightarrow X$, the covering $\mathbf{cov}(U \rightarrow X)$ consists of families of morphisms $\{(U_i \rightarrow X) \rightarrow (U \rightarrow X)\}_i$ such that $\{U_i \rightarrow U\}$ is in $\mathbf{cov}(U)$ in $\text{Cov}(\mathcal{T})$.

Note that if $\text{Cat}(\mathcal{T})$ has a terminal object \star , then $\mathcal{T} = \star_{\mathcal{T}}$.

Morphism of Sites

Consider two sites \mathcal{T} and \mathcal{S} , a *continuous function* $f : \mathcal{T} \rightarrow \mathcal{S}$

Morphism of Sites

Consider two sites \mathcal{T} and \mathcal{S} , a *continuous function* $f : \mathcal{T} \rightarrow \mathcal{S}$ is a functor

$$f^* : \text{Cat}(\mathcal{S}) \rightarrow \text{Cat}(\mathcal{T})$$

Morphism of Sites

Consider two sites \mathcal{T} and \mathcal{S} , a *continuous function* $f : \mathcal{T} \rightarrow \mathcal{S}$ is a functor

$$f^* : \text{Cat}(\mathcal{S}) \rightarrow \text{Cat}(\mathcal{T})$$

such that for any object U of $\text{Cat}(\mathcal{S})$

Morphism of Sites

Consider two sites \mathcal{T} and \mathcal{S} , a *continuous function* $f : \mathcal{T} \rightarrow \mathcal{S}$ is a functor

$$f^* : \text{Cat}(\mathcal{S}) \rightarrow \text{Cat}(\mathcal{T})$$

such that for any object U of $\text{Cat}(\mathcal{S})$

- (*takes covers to covers*) if $\{U_i \rightarrow U\}_i$ is a covering of U , then $\{f^*(U_i) \rightarrow f^*(U)\}_i$ is a covering of $f^*(U)$.

Morphism of Sites

Consider two sites \mathcal{T} and \mathcal{S} , a *continuous function* $f : \mathcal{T} \rightarrow \mathcal{S}$ is a functor

$$f^* : \text{Cat}(\mathcal{S}) \rightarrow \text{Cat}(\mathcal{T})$$

such that for any object U of $\text{Cat}(\mathcal{S})$

- (*takes covers to covers*) if $\{U_i \rightarrow U\}_i$ is a covering of U , then $\{f^*(U_i) \rightarrow f^*(U)\}_i$ is a covering of $f^*(U)$.
- (*preserves fibered products*) if $\{U_i \rightarrow U\}_i$ is a covering of U , then for a morphism $V \rightarrow U$ in $\text{Cat}(\mathcal{S})$ the canonical morphism

$$f^*(U_i \times_U V) \rightarrow f^*(U_i) \times_{f^*(U)} f^*(V)$$

is an isomorphism for all i .

Morphism of Sites

Consider two sites \mathcal{T} and \mathcal{S} , a *continuous function* $f : \mathcal{T} \rightarrow \mathcal{S}$ is a functor

$$f^* : \text{Cat}(\mathcal{S}) \rightarrow \text{Cat}(\mathcal{T})$$

such that for any object U of $\text{Cat}(\mathcal{S})$

- (*takes covers to covers*) if $\{U_i \rightarrow U\}_i$ is a covering of U , then $\{f^*(U_i) \rightarrow f^*(U)\}_i$ is a covering of $f^*(U)$.
- (*preserves fibered products*) if $\{U_i \rightarrow U\}_i$ is a covering of U , then for a morphism $V \rightarrow U$ in $\text{Cat}(\mathcal{S})$ the canonical morphism

$$f^*(U_i \times_U V) \rightarrow f^*(U_i) \times_{f^*(U)} f^*(V)$$

is an isomorphism for all i .

Remark

To be precise, what we've described here is really what is called a *Grothendieck pretopology*; this notion suffices for our purposes. So, as is commonly done, we will abuse terminology and keep using the term Grothendieck topology.

Topological Spaces are Sites

For a topological space X , we have the category of open sets $\text{Open}(X)$.

Topological Spaces are Sites

For a topological space X , we have the category of open sets $\text{Open}(X)$. We describe a site, which we denote X_{top} and is called the *small site of X* .

Topological Spaces are Sites

For a topological space X , we have the category of open sets $\text{Open}(X)$. We describe a site, which we denote X_{top} and is called the *small site of X* .

We let $\text{Cat}(X_{\text{top}}) = \text{Open}(X)$ and for any open set U , $\text{cov}(U)$ consists of families of morphisms $\{U_i \hookrightarrow U\}_i$ such that $U = \cup_i U_i$.

Topological Spaces are Sites

For a topological space X , we have the category of open sets $\text{Open}(X)$. We describe a site, which we denote X_{top} and is called the *small site of X* .

We let $\text{Cat}(X_{\text{top}}) = \text{Open}(X)$ and for any open set U , $\mathbf{cov}(U)$ consists of families of morphisms $\{U_i \hookrightarrow U\}_i$ such that $U = \cup_i U_i$.

Let X and Y be topological spaces and let $f : X \rightarrow Y$ be a continuous map (that is, a morphism in the category of topological spaces),

Topological Spaces are Sites

For a topological space X , we have the category of open sets $\text{Open}(X)$. We describe a site, which we denote X_{top} and is called the *small site of X* .

We let $\text{Cat}(X_{\text{top}}) = \text{Open}(X)$ and for any open set U , $\mathbf{cov}(U)$ consists of families of morphisms $\{U_i \hookrightarrow U\}_i$ such that $U = \cup_i U_i$.

Let X and Y be topological spaces and let $f : X \rightarrow Y$ be a continuous map (that is, a morphism in the category of topological spaces), then this gives us a continuous function of sites

$$f_{\text{top}} : X_{\text{top}} \rightarrow Y_{\text{top}}$$

given by the functor $f^{-1} : \text{Open}(Y) \rightarrow \text{Open}(X)$.

Topological Spaces are Sites

For a topological space X , we have the category of open sets $\text{Open}(X)$. We describe a site, which we denote X_{top} and is called the *small site of X* .

We let $\text{Cat}(X_{\text{top}}) = \text{Open}(X)$ and for any open set U , $\text{cov}(U)$ consists of families of morphisms $\{U_i \hookrightarrow U\}_i$ such that $U = \cup_i U_i$.

Let X and Y be topological spaces and let $f : X \rightarrow Y$ be a continuous map (that is, a morphism in the category of topological spaces), then this gives us a continuous function of sites

$$f_{\text{top}} : X_{\text{top}} \rightarrow Y_{\text{top}}$$

given by the functor $f^{-1} : \text{Open}(Y) \rightarrow \text{Open}(X)$.

That this is a functor is simply the fact that the inverse image map respects inclusion.

Topological Spaces are Sites

For a topological space X , we have the category of open sets $\text{Open}(X)$. We describe a site, which we denote X_{top} and is called the *small site of X* .

We let $\text{Cat}(X_{\text{top}}) = \text{Open}(X)$ and for any open set U , $\text{cov}(U)$ consists of families of morphisms $\{U_i \hookrightarrow U\}_i$ such that $U = \cup_i U_i$.

Let X and Y be topological spaces and let $f : X \rightarrow Y$ be a continuous map (that is, a morphism in the category of topological spaces), then this gives us a continuous function of sites

$$f_{\text{top}} : X_{\text{top}} \rightarrow Y_{\text{top}}$$

given by the functor $f^{-1} : \text{Open}(Y) \rightarrow \text{Open}(X)$.

That this is a functor is simply the fact that the inverse image map respects inclusion. That it describes a continuous function of sites is because if $U = \cup_i U_i$, then $f^{-1}(U) = \cup_i f^{-1}(U_i)$ (takes covers to covers)

Topological Spaces are Sites

For a topological space X , we have the category of open sets $\text{Open}(X)$. We describe a site, which we denote X_{top} and is called the *small site of X* .

We let $\text{Cat}(X_{\text{top}}) = \text{Open}(X)$ and for any open set U , $\text{cov}(U)$ consists of families of morphisms $\{U_i \hookrightarrow U\}_i$ such that $U = \cup_i U_i$.

Let X and Y be topological spaces and let $f : X \rightarrow Y$ be a continuous map (that is, a morphism in the category of topological spaces), then this gives us a continuous function of sites

$$f_{\text{top}} : X_{\text{top}} \rightarrow Y_{\text{top}}$$

given by the functor $f^{-1} : \text{Open}(Y) \rightarrow \text{Open}(X)$.

That this is a functor is simply the fact that the inverse image map respects inclusion. That it describes a continuous function of sites is because if $U = \cup_i U_i$, then $f^{-1}(U) = \cup_i f^{-1}(U_i)$ (takes covers to covers) and $f^{-1}(U \cap V) = f^{-1}(U) \cap f^{-1}(V)$ (preserves fibered products).

Topological Spaces are Sites

For a topological space X , we have the category of open sets $\text{Open}(X)$. We describe a site, which we denote X_{top} and is called the *small site of X* .

We let $\text{Cat}(X_{\text{top}}) = \text{Open}(X)$ and for any open set U , $\text{cov}(U)$ consists of families of morphisms $\{U_i \hookrightarrow U\}_i$ such that $U = \cup_i U_i$.

Let X and Y be topological spaces and let $f : X \rightarrow Y$ be a continuous map (that is, a morphism in the category of topological spaces), then this gives us a continuous function of sites

$$f_{\text{top}} : X_{\text{top}} \rightarrow Y_{\text{top}}$$

given by the functor $f^{-1} : \text{Open}(Y) \rightarrow \text{Open}(X)$.

That this is a functor is simply the fact that the inverse image map respects inclusion. That it describes a continuous function of sites is because if $U = \cup_i U_i$, then $f^{-1}(U) = \cup_i f^{-1}(U_i)$ (takes covers to covers) and $f^{-1}(U \cap V) = f^{-1}(U) \cap f^{-1}(V)$ (preserves fibered products).

It's this example that motivates the notion of a continuous function of sites.

Category of Topological Spaces is a Site

Given the category of topological spaces \mathbf{Top} , we give two examples of Grothendieck topology on it as follows, where the corresponding sites are denoted as $\ast_{\mathbf{Top}}$ and $\ast_{\mathbf{Lh}}$:

Category of Topological Spaces is a Site

Given the category of topological spaces \mathbf{Top} , we give two examples of Grothendieck topology on it as follows, where the corresponding sites are denoted as $*_{\mathbf{Top}}$ and $*_{\mathbf{Lh}}$:

- $\mathbf{Cov}(*_{\mathbf{Top}})$.

To any space X , let $\mathbf{cov}(X)$ be the collection of families of morphisms $\{U_i \hookrightarrow X\}_i$ which are jointly surjective open embeddings.

Category of Topological Spaces is a Site

Given the category of topological spaces \mathbf{Top} , we give two examples of Grothendieck topology on it as follows, where the corresponding sites are denoted as $*_{\mathbf{Top}}$ and $*_{\mathbf{Lh}}$:

- $\mathbf{Cov}(*_{\mathbf{Top}})$.

To any space X , let $\mathbf{cov}(X)$ be the collection of families of morphisms $\{U_i \hookrightarrow X\}_i$ which are jointly surjective open embeddings. That is, $U_i \hookrightarrow X$ is an injective open continuous map, for each i , such that the induced map $\coprod_i U_i \rightarrow X$ is surjective.

Category of Topological Spaces is a Site

Given the category of topological spaces \mathbf{Top} , we give two examples of Grothendieck topology on it as follows, where the corresponding sites are denoted as $*_{\mathbf{Top}}$ and $*_{\mathbf{Lh}}$:

- $\mathbf{Cov}(*_{\mathbf{Top}})$.

To any space X , let $\mathbf{cov}(X)$ be the collection of families of morphisms $\{U_i \hookrightarrow X\}_i$ which are jointly surjective open embeddings. That is, $U_i \hookrightarrow X$ is an injective open continuous map, for each i , such that the induced map $\coprod_i U_i \rightarrow X$ is surjective.

The induced site with the underlying category \mathbf{Top}/X is denoted $X_{\mathbf{Top}}$ and called the *big site of X* .

If one considers the full subcategory \mathbf{Open}/X of \mathbf{Top}/X where the structure maps are open embeddings, we recover (equivalent to) $X_{\mathbf{top}}$, the *small site of X* .

Category of Topological Spaces is a Site

Given the category of topological spaces \mathbf{Top} , we give two examples of Grothendieck topology on it as follows, where the corresponding sites are denoted as $*_{\mathbf{Top}}$ and $*_{\mathbf{Lh}}$:

- $\mathbf{Cov}(*_{\mathbf{Top}})$.

To any space X , let $\mathbf{cov}(X)$ be the collection of families of morphisms $\{U_i \hookrightarrow X\}_i$ which are jointly surjective open embeddings. That is, $U_i \hookrightarrow X$ is an injective open continuous map, for each i , such that the induced map $\coprod_i U_i \rightarrow X$ is surjective.

The induced site with the underlying category \mathbf{Top}/X is denoted $X_{\mathbf{Top}}$ and called the *big site of X* .

If one considers the full subcategory \mathbf{Open}/X of \mathbf{Top}/X where the structure maps are open embeddings, we recover (equivalent to) $X_{\mathbf{top}}$, the *small site of X* .

Given a continuous map $f : Y \rightarrow X$, we get a continuous function of sites

$$f_{\mathbf{Top}} : Y_{\mathbf{Top}} \rightarrow X_{\mathbf{Top}}$$

Category of Topological Spaces is a Site

Given the category of topological spaces \mathbf{Top} , we give two examples of Grothendieck topology on it as follows, where the corresponding sites are denoted as $*_{\mathbf{Top}}$ and $*_{\mathbf{Lh}}$:

- $\mathbf{Cov}(*_{\mathbf{Top}})$.

To any space X , let $\mathbf{cov}(X)$ be the collection of families of morphisms $\{U_i \hookrightarrow X\}_i$ which are jointly surjective open embeddings. That is, $U_i \hookrightarrow X$ is an injective open continuous map, for each i , such that the induced map $\coprod_i U_i \rightarrow X$ is surjective.

The induced site with the underlying category \mathbf{Top}/X is denoted $X_{\mathbf{Top}}$ and called the *big site of X* .

If one considers the full subcategory \mathbf{Open}/X of \mathbf{Top}/X where the structure maps are open embeddings, we recover (equivalent to) $X_{\mathbf{top}}$, the small site of X .

Given a continuous map $f : Y \rightarrow X$, we get a continuous function of sites

$$f_{\mathbf{Top}} : Y_{\mathbf{Top}} \rightarrow X_{\mathbf{Top}}$$

given by the functor

$$f^* : \mathbf{Top}/X \rightarrow \mathbf{Top}/Y, \quad (U \rightarrow X) \mapsto (U \times_X Y \rightarrow Y)$$

(...)

Category of Topological Spaces is a Site

(...)

- $\mathbf{Cov}(*_{\mathbf{Lh}})$.

To any X , let $\mathbf{cov}(X)$ be the collection of morphisms $\{C \rightarrow X\}$ which are local homeomorphisms.

Category of Topological Spaces is a Site

(...)

- $\mathbf{Cov}(*_{\mathbf{Lh}})$.

To any X , let $\mathbf{cov}(X)$ be the collection of morphisms $\{C \rightarrow X\}$ which are local homeomorphisms.

The induced site with the underlying category \mathbf{Top}/X is denoted $X_{\mathbf{Lh}}$.

Category of Topological Spaces is a Site

(...)

- $\mathbf{Cov}(*_{\mathbf{Lh}})$.

To any X , let $\mathbf{cov}(X)$ be the collection of morphisms $\{C \rightarrow X\}$ which are local homeomorphisms.

The induced site with the underlying category \mathbf{Top}/X is denoted $X_{\mathbf{Lh}}$.

Similarly as above, a continuous map $f : Y \rightarrow X$ induces a continuous function of sites $f_{\mathbf{Lh}} : Y_{\mathbf{Lh}} \rightarrow X_{\mathbf{Lh}}$, given by the pullback.

Category of Topological Spaces is a Site

(...)

- $\mathbf{Cov}(*_{\mathbf{Lh}})$.

To any X , let $\mathbf{cov}(X)$ be the collection of morphisms $\{C \rightarrow X\}$ which are local homeomorphisms.

The induced site with the underlying category \mathbf{Top}/X is denoted $X_{\mathbf{Lh}}$.

Similarly as above, a continuous map $f : Y \rightarrow X$ induces a continuous function of sites $f_{\mathbf{Lh}} : Y_{\mathbf{Lh}} \rightarrow X_{\mathbf{Lh}}$, given by the pullback.

Category of Smooth Manifolds is a Site

Given the category of smooth manifolds \mathbf{SmMan} , we can define two site similar to the ones above.

Category of Topological Spaces is a Site

(...)

- $\mathbf{Cov}(*_{\mathbf{Lh}})$.

To any X , let $\mathbf{cov}(X)$ be the collection of morphisms $\{C \rightarrow X\}$ which are local homeomorphisms.

The induced site with the underlying category \mathbf{Top}/X is denoted $X_{\mathbf{Lh}}$.

Similarly as above, a continuous map $f : Y \rightarrow X$ induces a continuous function of sites $f_{\mathbf{Lh}} : Y_{\mathbf{Lh}} \rightarrow X_{\mathbf{Lh}}$, given by the pullback.

Category of Smooth Manifolds is a Site

Given the category of smooth manifolds \mathbf{SmMan} , we can define two site similar to the ones above.

The site $*_{\mathbf{Man}}^{\infty}$ has as coverings jointly surjective open embeddings as above,

Category of Topological Spaces is a Site

(...)

- $\mathbf{Cov}(*_{\text{Lh}})$.

To any X , let $\mathbf{cov}(X)$ be the collection of morphisms $\{C \rightarrow X\}$ which are local homeomorphisms.

The induced site with the underlying category \mathbf{Top}/X is denoted X_{Lh} .

Similarly as above, a continuous map $f : Y \rightarrow X$ induces a continuous function of sites $f_{\text{Lh}} : Y_{\text{Lh}} \rightarrow X_{\text{Lh}}$, given by the pullback.

Category of Smooth Manifolds is a Site

Given the category of smooth manifolds \mathbf{SmMan} , we can define two site similar to the ones above.

The site $*_{\text{Man}}^{\infty}$ has as coverings jointly surjective open embeddings as above, while the site $*_{\text{Ld}}^{\infty}$ has as coverings local diffeomorphisms.

Category of Topological Spaces is a Site

(...)

- $\text{Cov}(*_{\text{Lh}})$.

To any X , let $\mathbf{cov}(X)$ be the collection of morphisms $\{C \rightarrow X\}$ which are local homeomorphisms.

The induced site with the underlying category Top/X is denoted X_{Lh} .

Similarly as above, a continuous map $f : Y \rightarrow X$ induces a continuous function of sites $f_{\text{Lh}} : Y_{\text{Lh}} \rightarrow X_{\text{Lh}}$, given by the pullback.

Category of Smooth Manifolds is a Site

Given the category of smooth manifolds SmMan , we can define two site similar to the ones above.

The site $*_{\text{Man}}^{\infty}$ has as coverings jointly surjective open embeddings as above, while the site $*_{\text{Ld}}^{\infty}$ has as coverings local diffeomorphisms.

The induced sites with the underlying category SmMan/M are denoted M_{Man} and M_{Ld} respectively.

(...)

Category of Smooth Manifolds is a Site

(...) We give another example of a site which we denote $*_{\text{Subm}}^{\infty}$:

Category of Smooth Manifolds is a Site

(...) We give another example of a site which we denote $*_{\text{Subm}}^{\infty}$:

- $\text{Cov}(*_{\text{Subm}}^{\infty})$.

To any M , let $\text{cov}(M)$ be the collection of morphisms $\{N \rightarrow M\}$ which are submersions, that is, where the induced map on tangents spaces at each point is surjective.

The induced site with the underlying category SmMan/M is denoted M_{Sub} .

Category of Smooth Manifolds is a Site

(...) We give another example of a site which we denote $*_{\text{Subm}}^{\infty}$:

- $\text{Cov}(*_{\text{Subm}}^{\infty})$.

To any M , let $\text{cov}(M)$ be the collection of morphisms $\{N \rightarrow M\}$ which are submersions, that is, where the induced map on tangents spaces at each point is surjective.

The induced site with the underlying category SmMan/M is denoted M_{Sub} .

There's a continuous function of sites

$$M_{\text{Ld}} \rightarrow M_{\text{Subm}}$$

induced by the identity map on M , since any submersion is a local diffeomorphism.

Category of Smooth Manifolds is a Site

(...) We give another example of a site which we denote $*_{\text{Subm}}^{\infty}$:

- $\text{Cov}(*_{\text{Subm}}^{\infty})$.

To any M , let $\text{cov}(M)$ be the collection of morphisms $\{N \rightarrow M\}$ which are submersions, that is, where the induced map on tangents spaces at each point is surjective.

The induced site with the underlying category SmMan/M is denoted M_{Sub} .

There's a continuous function of sites

$$M_{\text{Ld}} \rightarrow M_{\text{Subm}}$$

induced by the identity map on M , since any submersion is a local diffeomorphism.

We then say that the Grothendieck topology of local diffeomorphisms is a finer topology than the Grothendieck topology of submersions.

Category of Smooth Manifolds is a Site

(...) We give another example of a site which we denote $*_{\text{Subm}}^{\infty}$:

- $\text{Cov}(*_{\text{Subm}}^{\infty})$.

To any M , let $\text{cov}(M)$ be the collection of morphisms $\{N \rightarrow M\}$ which are submersions, that is, where the induced map on tangents spaces at each point is surjective.

The induced site with the underlying category SmMan/M is denoted M_{Sub} .

There's a continuous function of sites

$$M_{\text{Ld}} \rightarrow M_{\text{Subm}}$$

induced by the identity map on M , since any submersion is a local diffeomorphism.

We then say that the Grothendieck topology of local diffeomorphisms is a finer topology than the Grothendieck topology of submersions.

A word of warning is in order: for the topologies defined above there's a subtle point at play, that the fibered products (as defined in the category of topological spaces) exists in the category of smooth manifolds for the chosen morphisms.

Category of Smooth Manifolds is a Site

(...) We give another example of a site which we denote $*_{\text{Subm}}^{\infty}$:

- $\text{Cov}(*_{\text{Subm}}^{\infty})$.

To any M , let $\text{cov}(M)$ be the collection of morphisms $\{N \rightarrow M\}$ which are submersions, that is, where the induced map on tangents spaces at each point is surjective.

The induced site with the underlying category SmMan/M is denoted M_{Sub} .

There's a continuous function of sites

$$M_{\text{Ld}} \rightarrow M_{\text{Subm}}$$

induced by the identity map on M , since any submersion is a local diffeomorphism.

We then say that the Grothendieck topology of local diffeomorphisms is a finer topology than the Grothendieck topology of submersions.

A word of warning is in order: for the topologies defined above there's a subtle point at play, that the fibered products (as defined in the category of topological spaces) exists in the category of smooth manifolds for the chosen morphisms. It's not true that these fibered products exist in general (take a smooth map with a critical value as an example for this fact).

Category of finite G -sets is a Site

Paul Balmer describes a Grothendieck topology on the category of finite G -sets, for a fixed prime p in [2].

Category of finite G -sets is a Site

Paul Balmer describes a Grothendieck topology on the category of finite G -sets, for a fixed prime p in [2]. He calls it the *sipp topology* where sipp stands for *prime-to- p index stabilisers* (in French).

Category of finite G -sets is a Site

Paul Balmer describes a Grothendieck topology on the category of finite G -sets, for a fixed prime p in [2]. He calls it the *sipp topology* where *sipp* stands for *prime-to- p index stabilisers* (in French). This site can reasonably be denoted as e_{sipp} , where e is the neutral element of G , since e can be taken to be the terminal object.

Category of finite G -sets is a Site

Paul Balmer describes a Grothendieck topology on the category of finite G -sets, for a fixed prime p in [2]. He calls it the *sipp topology* where sipp stands for *prime-to- p index stabilisers* (in French). This site can reasonably be denoted as e_{sipp} , where e is the neutral element of G , since e can be taken to be the terminal object.

Let X be a finite G -set, $\mathbf{cov}(X)$ is taken to be the collection of families of G -equivariant maps $\{\alpha_i : U_i \rightarrow X\}_i$ such that

Category of finite G -sets is a Site

Paul Balmer describes a Grothendieck topology on the category of finite G -sets, for a fixed prime p in [2]. He calls it the *sipp topology* where *sipp* stands for *prime-to- p index stabilisers* (in French). This site can reasonably be denoted as e_{sipp} , where e is the neutral element of G , since e can be taken to be the terminal object.

Let X be a finite G -set, $\mathbf{cov}(X)$ is taken to be the collection of families of G -equivariant maps $\{\alpha_i : U_i \rightarrow X\}_i$ such that

- for every $x \in X$ there exists an i and $u \in U_i$ such that $\alpha_i(u) = x$;

Category of finite G -sets is a Site

Paul Balmer describes a Grothendieck topology on the category of finite G -sets, for a fixed prime p in [2]. He calls it the *sipp topology* where *sipp* stands for *prime-to- p index stabilisers* (in French). This site can reasonably be denoted as e_{sipp} , where e is the neutral element of G , since e can be taken to be the terminal object.

Let X be a finite G -set, $\mathbf{cov}(X)$ is taken to be the collection of families of G -equivariant maps $\{\alpha_i : U_i \rightarrow X\}_i$ such that

- for every $x \in X$ there exists an i and $u \in U_i$ such that $\alpha_i(u) = x$; and
- the index $[\text{Stab}_G(x) : \text{Stab}_G(u)]$ is prime to p .

Category of finite G -sets is a Site

Paul Balmer describes a Grothendieck topology on the category of finite G -sets, for a fixed prime p in [2]. He calls it the *sipp topology* where *sipp* stands for *prime-to- p index stabilisers* (in French). This site can reasonably be denoted as e_{sipp} , where e is the neutral element of G , since e can be taken to be the terminal object.

Let X be a finite G -set, $\mathbf{cov}(X)$ is taken to be the collection of families of G -equivariant maps $\{\alpha_i : U_i \rightarrow X\}_i$ such that

- for every $x \in X$ there exists an i and $u \in U_i$ such that $\alpha_i(u) = x$; and
- the index $[\text{Stab}_G(x) : \text{Stab}_G(u)]$ is prime to p .

This site enjoys some nice properties,

Category of finite G -sets is a Site

Paul Balmer describes a Grothendieck topology on the category of finite G -sets, for a fixed prime p in [2]. He calls it the *sipp topology* where *sipp* stands for *prime-to- p index stabilisers* (in French). This site can reasonably be denoted as e_{sipp} , where e is the neutral element of G , since e can be taken to be the terminal object.

Let X be a finite G -set, $\mathbf{cov}(X)$ is taken to be the collection of families of G -equivariant maps $\{\alpha_i : U_i \rightarrow X\}_i$ such that

- for every $x \in X$ there exists an i and $u \in U_i$ such that $\alpha_i(u) = x$; and
- the index $[\text{Stab}_G(x) : \text{Stab}_G(u)]$ is prime to p .

This site enjoys some nice properties, for example it's a Noetherian (or quasi-compact) site: every covering has a finite sub-coverings, since we're working with finite G -sets;

Category of finite G -sets is a Site

Paul Balmer describes a Grothendieck topology on the category of finite G -sets, for a fixed prime p in [2]. He calls it the *sipp topology* where *sipp* stands for *prime-to- p index stabilisers* (in French). This site can reasonably be denoted as e_{sipp} , where e is the neutral element of G , since e can be taken to be the terminal object.

Let X be a finite G -set, $\mathbf{cov}(X)$ is taken to be the collection of families of G -equivariant maps $\{\alpha_i : U_i \rightarrow X\}_i$ such that

- for every $x \in X$ there exists an i and $u \in U_i$ such that $\alpha_i(u) = x$; and
- the index $[\text{Stab}_G(x) : \text{Stab}_G(u)]$ is prime to p .

This site enjoys some nice properties, for example it's a Noetherian (or quasi-compact) site: every covering has a finite sub-coverings, since we're working with finite G -sets; and the Grothendieck topology is subcanonical: every representable presheaf is a sheaf.

Sites associated to the category of Schemes

Let's quickly recall certain important classes of morphisms in \mathbf{Sch} , the category of Schemes:

Sites associated to the category of Schemes

Let's quickly recall certain important classes of morphisms in Sch , the category of Schemes:

- A morphism $f : X \rightarrow Y$ is *locally of finite presentation*,

Sites associated to the category of Schemes

Let's quickly recall certain important classes of morphisms in Sch , the category of Schemes:

- A morphism $f : X \rightarrow Y$ is *locally of finite presentation*, if for any affine open set $\text{Spec } A \subseteq Y$ and affine open set $\text{Spec } B \subseteq f^{-1}(\text{Spec } A)$ the corresponding ring map $A \rightarrow B$ makes B an A -algebra of finite presentation.

Sites associated to the category of Schemes

Let's quickly recall certain important classes of morphisms in Sch , the category of Schemes:

- A morphism $f : X \rightarrow Y$ is *locally of finite presentation*, if for any affine open set $\text{Spec } A \subseteq Y$ and affine open set $\text{Spec } B \subseteq f^{-1}(\text{Spec } A)$ the corresponding ring map $A \rightarrow B$ makes B an A -algebra of finite presentation. That is, there exists a surjective ring map $A[x_1, \dots, x_n] \twoheadrightarrow B$ such that the kernel is a finitely generated ideal of $A[x_1, \dots, x_n]$.

Sites associated to the category of Schemes

Let's quickly recall certain important classes of morphisms in Sch , the category of Schemes:

- A morphism $f : X \rightarrow Y$ is *locally of finite presentation*, if for any affine open set $\text{Spec } A \subseteq Y$ and affine open set $\text{Spec } B \subseteq f^{-1}(\text{Spec } A)$ the corresponding ring map $A \rightarrow B$ makes B an A -algebra of finite presentation. That is, there exists a surjective ring map $A[x_1, \dots, x_n] \twoheadrightarrow B$ such that the kernel is a finitely generated ideal of $A[x_1, \dots, x_n]$.
- A morphism $f : X \rightarrow Y$ that is locally of finite presentation is *unramified*,

Sites associated to the category of Schemes

Let's quickly recall certain important classes of morphisms in Sch , the category of Schemes:

- A morphism $f : X \rightarrow Y$ is *locally of finite presentation*, if for any affine open set $\text{Spec } A \subseteq Y$ and affine open set $\text{Spec } B \subseteq f^{-1}(\text{Spec } A)$ the corresponding ring map $A \rightarrow B$ makes B an A -algebra of finite presentation. That is, there exists a surjective ring map $A[x_1, \dots, x_n] \twoheadrightarrow B$ such that the kernel is a finitely generated ideal of $A[x_1, \dots, x_n]$.
- A morphism $f : X \rightarrow Y$ that is locally of finite presentation is *unramified*, if for each point $y \in Y$, the fiber $f^{-1}(y) = X \times_Y \text{Spec } \kappa(y) \cong \coprod_i \text{Spec } k_i$, where k_i 's are finite separable extensions of the residue field at y , $\kappa(y)$.

Sites associated to the category of Schemes

Let's quickly recall certain important classes of morphisms in Sch , the category of Schemes:

- A morphism $f : X \rightarrow Y$ is *locally of finite presentation*, if for any affine open set $\text{Spec } A \subseteq Y$ and affine open set $\text{Spec } B \subseteq f^{-1}(\text{Spec } A)$ the corresponding ring map $A \rightarrow B$ makes B an A -algebra of finite presentation. That is, there exists a surjective ring map $A[x_1, \dots, x_n] \twoheadrightarrow B$ such that the kernel is a finitely generated ideal of $A[x_1, \dots, x_n]$.
- A morphism $f : X \rightarrow Y$ that is locally of finite presentation is *unramified*, if for each point $y \in Y$, the fiber $f^{-1}(y) = X \times_Y \text{Spec } \kappa(y) \cong \coprod_i \text{Spec } k_i$, where k_i 's are finite separable extensions of the residue field at y , $\kappa(y)$.
- A morphism $f : X \rightarrow Y$ is *flat*,

Sites associated to the category of Schemes

Let's quickly recall certain important classes of morphisms in Sch , the category of Schemes:

- A morphism $f : X \rightarrow Y$ is *locally of finite presentation*, if for any affine open set $\text{Spec } A \subseteq Y$ and affine open set $\text{Spec } B \subseteq f^{-1}(\text{Spec } A)$ the corresponding ring map $A \rightarrow B$ makes B an A -algebra of finite presentation. That is, there exists a surjective ring map $A[x_1, \dots, x_n] \twoheadrightarrow B$ such that the kernel is a finitely generated ideal of $A[x_1, \dots, x_n]$.
- A morphism $f : X \rightarrow Y$ that is locally of finite presentation is *unramified*, if for each point $y \in Y$, the fiber $f^{-1}(y) = X \times_Y \text{Spec } \kappa(y) \cong \coprod_i \text{Spec } k_i$, where k_i 's are finite separable extensions of the residue field at y , $\kappa(y)$.
- A morphism $f : X \rightarrow Y$ is *flat*, if for each point $x \in X$ the map of local rings $\mathcal{O}_{Y,f(x)} \rightarrow \mathcal{O}_{X,x}$ makes $\mathcal{O}_{X,x}$ a flat $\mathcal{O}_{Y,f(x)}$ -module.

Sites associated to the category of Schemes

Let's quickly recall certain important classes of morphisms in Sch , the category of Schemes:

- A morphism $f : X \rightarrow Y$ is *locally of finite presentation*, if for any affine open set $\text{Spec } A \subseteq Y$ and affine open set $\text{Spec } B \subseteq f^{-1}(\text{Spec } A)$ the corresponding ring map $A \rightarrow B$ makes B an A -algebra of finite presentation. That is, there exists a surjective ring map $A[x_1, \dots, x_n] \twoheadrightarrow B$ such that the kernel is a finitely generated ideal of $A[x_1, \dots, x_n]$.
- A morphism $f : X \rightarrow Y$ that is locally of finite presentation is *unramified*, if for each point $y \in Y$, the fiber $f^{-1}(y) = X \times_Y \text{Spec } \kappa(y) \cong \coprod_i \text{Spec } k_i$, where k_i 's are finite separable extensions of the residue field at y , $\kappa(y)$.
- A morphism $f : X \rightarrow Y$ is *flat*, if for each point $x \in X$ the map of local rings $\mathcal{O}_{Y,f(x)} \rightarrow \mathcal{O}_{X,x}$ makes $\mathcal{O}_{X,x}$ a flat $\mathcal{O}_{Y,f(x)}$ -module.
- A morphism $f : X \rightarrow Y$ is *étale*, if it's flat and unramified.

Sites associated to the category of Schemes

Let's quickly recall certain important classes of morphisms in Sch , the category of Schemes:

- A morphism $f : X \rightarrow Y$ is *locally of finite presentation*, if for any affine open set $\text{Spec } A \subseteq Y$ and affine open set $\text{Spec } B \subseteq f^{-1}(\text{Spec } A)$ the corresponding ring map $A \rightarrow B$ makes B an A -algebra of finite presentation. That is, there exists a surjective ring map $A[x_1, \dots, x_n] \twoheadrightarrow B$ such that the kernel is a finitely generated ideal of $A[x_1, \dots, x_n]$.
- A morphism $f : X \rightarrow Y$ that is locally of finite presentation is *unramified*, if for each point $y \in Y$, the fiber $f^{-1}(y) = X \times_Y \text{Spec } \kappa(y) \cong \coprod_i \text{Spec } k_i$, where k_i 's are finite separable extensions of the residue field at y , $\kappa(y)$.
- A morphism $f : X \rightarrow Y$ is *flat*, if for each point $x \in X$ the map of local rings $\mathcal{O}_{Y,f(x)} \rightarrow \mathcal{O}_{X,x}$ makes $\mathcal{O}_{X,x}$ a flat $\mathcal{O}_{Y,f(x)}$ -module.
- A morphism $f : X \rightarrow Y$ is *étale*, if it's flat and unramified.
- A morphism $f : X \rightarrow Y$ is *fppf*, if it's flat and locally of finite presentation; where fppf stands for *faithfully flat and of finite presentation* (in French)

(...)

Sites associated to the category of Schemes

We now describe three sites associated to the category of schemes

- The big Zariski site, which we denote Sch_{Zar} ,

Sites associated to the category of Schemes

We now describe three sites associated to the category of schemes

- The big Zariski site, which we denote Sch_{Zar} , is the site where for a given scheme T , $\text{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective open embeddings.

Sites associated to the category of Schemes

We now describe three sites associated to the category of schemes

- The big Zariski site, which we denote Sch_{Zar} , is the site where for a given scheme T , $\text{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective open embeddings.

For a scheme X , we get the induced *big Zariski site of X* , $(\text{Sch}/X)_{\text{Zar}}$.

Sites associated to the category of Schemes

We now describe three sites associated to the category of schemes

- The big Zariski site, which we denote Sch_{Zar} , is the site where for a given scheme T , $\text{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective open embeddings.

For a scheme X , we get the induced *big Zariski site of X* , $(\text{Sch}/X)_{\text{Zar}}$.

The *small Zariski site of X* , denoted X_{Zar} , is the site where the underlying category is the full subcategory of Sch/X such that the structure morphism is an open embedding.

Sites associated to the category of Schemes

We now describe three sites associated to the category of schemes

- The big Zariski site, which we denote Sch_{Zar} , is the site where for a given scheme T , $\text{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective open embeddings.

For a scheme X , we get the induced *big Zariski site of X* , $(\text{Sch}/X)_{\text{Zar}}$.

The *small Zariski site of X* , denoted X_{Zar} , is the site where the underlying category is the full subcategory of Sch/X such that the structure morphism is an open embedding.

- The big Étale site, which we denote $\text{Sch}_{\text{ét}}$,

Sites associated to the category of Schemes

We now describe three sites associated to the category of schemes

- The big Zariski site, which we denote Sch_{Zar} , is the site where for a given scheme T , $\text{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective open embeddings.

For a scheme X , we get the induced *big Zariski site of X* , $(\text{Sch}/X)_{\text{Zar}}$.

The *small Zariski site of X* , denoted X_{Zar} , is the site where the underlying category is the full subcategory of Sch/X such that the structure morphism is an open embedding.

- The big Étale site, which we denote $\text{Sch}_{\text{ét}}$, is the site where for a given scheme T , $\text{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective étale morphisms.

Sites associated to the category of Schemes

We now describe three sites associated to the category of schemes

- The big Zariski site, which we denote Sch_{Zar} , is the site where for a given scheme T , $\mathbf{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective open embeddings.

For a scheme X , we get the induced *big Zariski site of X* , $(\text{Sch}/X)_{\text{Zar}}$.

The *small Zariski site of X* , denoted X_{Zar} , is the site where the underlying category is the full subcategory of Sch/X such that the structure morphism is an open embedding.

- The big Étale site, which we denote $\text{Sch}_{\text{ét}}$, is the site where for a given scheme T , $\mathbf{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective étale morphisms.

For a scheme X , we get the induced big Étale site of X , $(\text{Sch}/X)_{\text{ét}}$.

Sites associated to the category of Schemes

We now describe three sites associated to the category of schemes

- The big Zariski site, which we denote Sch_{Zar} , is the site where for a given scheme T , $\mathbf{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective open embeddings.

For a scheme X , we get the induced *big Zariski site of X* , $(\text{Sch}/X)_{\text{Zar}}$.

The *small Zariski site of X* , denoted X_{Zar} , is the site where the underlying category is the full subcategory of Sch/X such that the structure morphism is an open embedding.

- The big Étale site, which we denote $\text{Sch}_{\text{ét}}$, is the site where for a given scheme T , $\mathbf{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective étale morphisms.

For a scheme X , we get the induced big Étale site of X , $(\text{Sch}/X)_{\text{ét}}$.

The small Étale site of X , denoted $X_{\text{ét}}$, is the site where the underlying category is the full subcategory of Sch/X such that the structure morphism is an étale morphism.

Sites associated to the category of Schemes

We now describe three sites associated to the category of schemes

- The big Zariski site, which we denote Sch_{Zar} , is the site where for a given scheme T , $\mathbf{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective open embeddings.

For a scheme X , we get the induced *big Zariski site of X* , $(\text{Sch}/X)_{\text{Zar}}$.

The *small Zariski site of X* , denoted X_{Zar} , is the site where the underlying category is the full subcategory of Sch/X such that the structure morphism is an open embedding.

- The big Étale site, which we denote $\text{Sch}_{\text{ét}}$, is the site where for a given scheme T , $\mathbf{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective étale morphisms.

For a scheme X , we get the induced big Étale site of X , $(\text{Sch}/X)_{\text{ét}}$.

The small Étale site of X , denoted $X_{\text{ét}}$, is the site where the underlying category is the full subcategory of Sch/X such that the structure morphism is an étale morphism.

When one says the étale topology on X or the étale site of X , one is referring to the small étale site of X .

(...)

Sites associated to the category of Schemes

(...)

- The big fppf site, which we denote Sch_{fppf} is the site where for a given scheme T , $\mathbf{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective fppf morphisms.

Sites associated to the category of Schemes

(...)

- The big fppf site, which we denote Sch_{fppf} is the site where for a given scheme T , $\mathbf{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective fppf morphisms.

For a scheme X , we get the induced big fppf site of X , $(\text{Sch}/X)_{\text{fppf}}$ which we denote X_{fppf} .

Sites associated to the category of Schemes

(...)

- The big fppf site, which we denote Sch_{fppf} is the site where for a given scheme T , $\mathbf{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective fppf morphisms.

For a scheme X , we get the induced big fppf site of X , $(\text{Sch}/X)_{\text{fppf}}$ which we denote X_{fppf} .

When one says the fppf topology on X or the fppf site of X , one is referring to the big fppf site of X .

Sites associated to the category of Schemes

(...)

- The big fppf site, which we denote Sch_{fppf} is the site where for a given scheme T , $\text{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective fppf morphisms.

For a scheme X , we get the induced big fppf site of X , $(\text{Sch}/X)_{\text{fppf}}$ which we denote X_{fppf} .

When one says the fppf topology on X or the fppf site of X , one is referring to the big fppf site of X .

One has that every Zariski covering is an étale covering, and every étale covering is an fppf covering.

Sites associated to the category of Schemes

(...)

- The big fppf site, which we denote Sch_{fppf} is the site where for a given scheme T , $\mathbf{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective fppf morphisms.

For a scheme X , we get the induced big fppf site of X , $(\text{Sch}/X)_{\text{fppf}}$ which we denote X_{fppf} .

When one says the fppf topology on X or the fppf site of X , one is referring to the big fppf site of X .

One has that every Zariski covering is an étale covering, and every étale covering is an fppf covering. Therefore, given a scheme X , the identity map induces the following continuous function of sites

$$X_{\text{fppf}} \rightarrow X_{\text{ét}} \rightarrow X_{\text{Zar}}$$

Sites associated to the category of Schemes

(...)

- The big fppf site, which we denote Sch_{fppf} is the site where for a given scheme T , $\text{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective fppf morphisms.

For a scheme X , we get the induced big fppf site of X , $(\text{Sch}/X)_{\text{fppf}}$ which we denote X_{fppf} .

When one says the fppf topology on X or the fppf site of X , one is referring to the big fppf site of X .

One has that every Zariski covering is an étale covering, and every étale covering is an fppf covering. Therefore, given a scheme X , the identity map induces the following continuous function of sites

$$X_{\text{fppf}} \rightarrow X_{\text{ét}} \rightarrow X_{\text{Zar}}$$

Therefore the fppf topology is finer than the étale topology and Zariski topology, while the étale topology is finer than the Zariski topology.

Sites associated to the category of Schemes

(...)

- The big fppf site, which we denote Sch_{fppf} is the site where for a given scheme T , $\text{cov}(T)$ is the collection of morphisms $\{T_i \rightarrow T\}_i$ that are jointly surjective fppf morphisms.

For a scheme X , we get the induced big fppf site of X , $(\text{Sch}/X)_{\text{fppf}}$ which we denote X_{fppf} .

When one says the fppf topology on X or the fppf site of X , one is referring to the big fppf site of X .

One has that every Zariski covering is an étale covering, and every étale covering is an fppf covering. Therefore, given a scheme X , the identity map induces the following continuous function of sites

$$X_{\text{fppf}} \rightarrow X_{\text{ét}} \rightarrow X_{\text{Zar}}$$

Therefore the fppf topology is finer than the étale topology and Zariski topology, while the étale topology is finer than the Zariski topology. We sometimes work with a finer topology than all of these called the fpqc topology.

Sheaves and Sheaf Cohomology on Sites

× × ×

Presheaves & Sheaves

A *presheaf* of abelian groups on a category \mathcal{C} , or simply a *presheaf*,

Presheaves & Sheaves

A *presheaf of abelian groups on a category \mathcal{C}* , or simply a *presheaf*, is a functor

$$\mathcal{F} : \mathcal{C}^{\text{op}} \rightarrow \text{Ab}.$$

Presheaves & Sheaves

A *presheaf of abelian groups on a category \mathcal{C}* , or simply a *presheaf*, is a functor

$$\mathcal{F} : \mathcal{C}^{\text{op}} \rightarrow \text{Ab}.$$

As before, an element of $\mathcal{F}(U)$ is called a section of \mathcal{F} over U , and a morphism of presheaves is a morphism of functors.

Presheaves & Sheaves

A *presheaf of abelian groups on a category \mathcal{C}* , or simply a *presheaf*, is a functor

$$\mathcal{F} : \mathcal{C}^{\text{op}} \rightarrow \text{Ab}.$$

As before, an element of $\mathcal{F}(U)$ is called a section of \mathcal{F} over U , and a morphism of presheaves is a morphism of functors.

Let \mathcal{T} be a site, then a *sheaf of abelian groups on a site \mathcal{T}* , or simply a *sheaf*,

Presheaves & Sheaves

A *presheaf of abelian groups on a category \mathcal{C}* , or simply a *presheaf*, is a functor

$$\mathcal{F} : \mathcal{C}^{\text{op}} \rightarrow \text{Ab}.$$

As before, an element of $\mathcal{F}(U)$ is called a section of \mathcal{F} over U , and a morphism of presheaves is a morphism of functors.

Let \mathcal{T} be a site, then a *sheaf of abelian groups on a site \mathcal{T}* , or simply a *sheaf*, is a presheaf of abelian group on $\text{Cat}(\mathcal{T})$ such that for any covering $\{U_i \rightarrow U\}_i$ in $\text{Cov}(\mathcal{T})$

Presheaves & Sheaves

A *presheaf of abelian groups on a category \mathcal{C}* , or simply a *presheaf*, is a functor

$$\mathcal{F} : \mathcal{C}^{\text{op}} \rightarrow \text{Ab}.$$

As before, an element of $\mathcal{F}(U)$ is called a section of \mathcal{F} over U , and a morphism of presheaves is a morphism of functors.

Let \mathcal{T} be a site, then a *sheaf of abelian groups on a site \mathcal{T}* , or simply a *sheaf*, is a presheaf of abelian group on $\text{Cat}(\mathcal{T})$ such that for any covering $\{U_i \rightarrow U\}_i$ in $\text{Cov}(\mathcal{T})$ the following is an exact sequence

$$0 \longrightarrow \mathcal{F}(U) \longrightarrow \prod_i \mathcal{F}(U_i) \rightrightarrows \prod_{i,j} \mathcal{F}(U_i \times_U U_j)$$

Presheaves & Sheaves

A *presheaf of abelian groups on a category \mathcal{C}* , or simply a *presheaf*, is a functor

$$\mathcal{F} : \mathcal{C}^{\text{op}} \rightarrow \text{Ab}.$$

As before, an element of $\mathcal{F}(U)$ is called a section of \mathcal{F} over U , and a morphism of presheaves is a morphism of functors.

Let \mathcal{T} be a site, then a *sheaf of abelian groups on a site \mathcal{T}* , or simply a *sheaf*, is a presheaf of abelian group on $\text{Cat}(\mathcal{T})$ such that for any covering $\{U_i \rightarrow U\}_i$ in $\text{Cov}(\mathcal{T})$ the following is an exact sequence

$$0 \longrightarrow \mathcal{F}(U) \longrightarrow \prod_i \mathcal{F}(U_i) \rightrightarrows \prod_{i,j} \mathcal{F}(U_i \times_U U_j)$$

Naturally, relaxing the target category, we obtain the notion of *sheaves of sets, groups, rings, modules, algebras* etc.

Presheaves & Sheaves

A *presheaf of abelian groups on a category \mathcal{C}* , or simply a *presheaf*, is a functor

$$\mathcal{F} : \mathcal{C}^{\text{op}} \rightarrow \text{Ab}.$$

As before, an element of $\mathcal{F}(U)$ is called a section of \mathcal{F} over U , and a morphism of presheaves is a morphism of functors.

Let \mathcal{T} be a site, then a *sheaf of abelian groups on a site \mathcal{T}* , or simply a *sheaf*, is a presheaf of abelian group on $\text{Cat}(\mathcal{T})$ such that for any covering $\{U_i \rightarrow U\}_i$ in $\text{Cov}(\mathcal{T})$ the following is an exact sequence

$$0 \longrightarrow \mathcal{F}(U) \longrightarrow \prod_i \mathcal{F}(U_i) \rightrightarrows \prod_{i,j} \mathcal{F}(U_i \times_U U_j)$$

Naturally, relaxing the target category, we obtain the notion of *sheaves of sets, groups, rings, modules, algebras* etc.

The category of sheaves $\text{Sh}(\mathcal{T})$, as before, is the full subcategory of sheaves of the category of presheaves $[\text{Cat}(\mathcal{T})^{\text{op}}, \text{Ab}]$.

Presheaves & Sheaves

A *presheaf of abelian groups on a category \mathcal{C}* , or simply a *presheaf*, is a functor

$$\mathcal{F} : \mathcal{C}^{\text{op}} \rightarrow \text{Ab}.$$

As before, an element of $\mathcal{F}(U)$ is called a section of \mathcal{F} over U , and a morphism of presheaves is a morphism of functors.

Let \mathcal{T} be a site, then a *sheaf of abelian groups on a site \mathcal{T}* , or simply a *sheaf*, is a presheaf of abelian group on $\text{Cat}(\mathcal{T})$ such that for any covering $\{U_i \rightarrow U\}_i$ in $\text{Cov}(\mathcal{T})$ the following is an exact sequence

$$0 \longrightarrow \mathcal{F}(U) \longrightarrow \prod_i \mathcal{F}(U_i) \rightrightarrows \prod_{i,j} \mathcal{F}(U_i \times_U U_j)$$

Naturally, relaxing the target category, we obtain the notion of *sheaves of sets, groups, rings, modules, algebras* etc.

The category of sheaves $\text{Sh}(\mathcal{T})$, as before, is the full subcategory of sheaves of the category of presheaves $[\text{Cat}(\mathcal{T})^{\text{op}}, \text{Ab}]$.

It's worth pointing out that if we're given a category \mathcal{C} which can be equipped with different Grothendieck topologies then the category of presheaves $[\mathcal{C}^{\text{op}}, \text{Ab}]$ has multiple subcategories of sheaves with respect each of these Grothendieck topologies.

Dictionary

Dictionary

Topological Space X	Site \mathcal{T}

Dictionary

Topological Space X	Site \mathcal{T}
U , an open subset of X , equivalently an object of $\mathbf{Open}(X)$	

Sheaves on Sites

Dictionary

Topological Space X	Site \mathcal{T}
U , an open subset of X , equivalently an object of $\mathbf{Open}(X)$	U , an object of $\mathbf{Cat}(\mathcal{T})$

Dictionary

Topological Space X	Site \mathcal{T}
U , an open subset of X , equivalently an object of $\mathbf{Open}(X)$	U , an object of $\mathbf{Cat}(\mathcal{T})$
$U = \cup_i U_i$, an open cover of U	

Sheaves on Sites

Dictionary

Topological Space X	Site \mathcal{T}
U , an open subset of X , equivalently an object of $\mathbf{Open}(X)$	U , an object of $\mathbf{Cat}(\mathcal{T})$
$U = \cup_i U_i$, an open cover of U	$\{U_i \rightarrow U\}_i$ in $\mathbf{cov}(U)$

Dictionary

Topological Space X	Site \mathcal{T}
U , an open subset of X , equivalently an object of $\mathbf{Open}(X)$	U , an object of $\mathbf{Cat}(\mathcal{T})$
$U = \cup_i U_i$, an open cover of U	$\{U_i \rightarrow U\}_i$ in $\mathbf{cov}(U)$
$f : X \rightarrow Y$ continuous map, $f^{-1} : \mathbf{Open}(Y) \rightarrow \mathbf{Open}(X)$ takes open covers to open covers, commutes with intersections	

Dictionary

Topological Space X	Site \mathcal{T}
U , an open subset of X , equivalently an object of $\mathbf{Open}(X)$	U , an object of $\mathbf{Cat}(\mathcal{T})$
$U = \cup_i U_i$, an open cover of U	$\{U_i \rightarrow U\}_i$ in $\mathbf{cov}(U)$
$f : X \rightarrow Y$ continuous map, $f^{-1} : \mathbf{Open}(Y) \rightarrow \mathbf{Open}(X)$ takes open covers to open covers, commutes with intersections	$f : \mathcal{T} \rightarrow \mathcal{S}$ continuous function, a functor $f^* : \mathbf{Cat}(\mathcal{S}) \rightarrow \mathbf{Cat}(\mathcal{T})$ takes open covers to open covers, commutes with fibered products

Dictionary

Topological Space X	Site \mathcal{T}
U , an open subset of X , equivalently an object of $\text{Open}(X)$	U , an object of $\text{Cat}(\mathcal{T})$
$U = \cup_i U_i$, an open cover of U	$\{U_i \rightarrow U\}_i$ in $\mathbf{cov}(U)$
$f : X \rightarrow Y$ continuous map, $f^{-1} : \text{Open}(Y) \rightarrow \text{Open}(X)$ takes open covers to open covers, commutes with intersections	$f : \mathcal{T} \rightarrow \mathcal{S}$ continuous function, a functor $f^* : \text{Cat}(\mathcal{S}) \rightarrow \text{Cat}(\mathcal{T})$ takes open covers to open covers, commutes with fibered products
Sheaf is a functor $\mathcal{F} : \text{Open}(X)^{\text{op}} \rightarrow \mathbf{Ab}$ satisfying two axioms, equivalently an (equaliser) exact sequence with respect to an open cover	

Dictionary

Topological Space X	Site \mathcal{T}
U , an open subset of X , equivalently an object of $\mathbf{Open}(X)$	U , an object of $\mathbf{Cat}(\mathcal{T})$
$U = \cup_i U_i$, an open cover of U	$\{U_i \rightarrow U\}_i$ in $\mathbf{cov}(U)$
$f : X \rightarrow Y$ continuous map, $f^{-1} : \mathbf{Open}(Y) \rightarrow \mathbf{Open}(X)$ takes open covers to open covers, commutes with intersections	$f : \mathcal{T} \rightarrow \mathcal{S}$ continuous function, a functor $f^* : \mathbf{Cat}(\mathcal{S}) \rightarrow \mathbf{Cat}(\mathcal{T})$ takes open covers to open covers, commutes with fibered products
Sheaf is a functor $\mathcal{F} : \mathbf{Open}(X)^{\mathbf{op}} \rightarrow \mathbf{Ab}$ satisfying two axioms, equivalently an (equaliser) exact sequence with respect to an open cover	Sheaf is a functor $\mathcal{F} : \mathbf{Cat}(\mathcal{T})^{\mathbf{op}} \rightarrow \mathbf{Ab}$ satisfying an (equaliser) exact sequence with respect to an open cover

Representable Sheaves

Let $E \in \{\text{Zar}, \text{ét}, \text{fppf}\}$ and consider the site X_E for a scheme X .

Representable Sheaves

Let $E \in \{\text{Zar}, \text{ét}, \text{fppf}\}$ and consider the site X_E for a scheme X . For any commutative X -group scheme G , we can consider the functor of points

$$h_G : (\text{Sch}/X)^{\text{op}} \rightarrow \text{Ab}, T \mapsto \text{Hom}_S(T, G).$$

Representable Sheaves

Let $E \in \{\text{Zar}, \text{ét}, \text{fppf}\}$ and consider the site X_E for a scheme X . For any commutative X -group scheme G , we can consider the functor of points

$$h_G : (\text{Sch}/X)^{\text{op}} \rightarrow \mathbf{Ab}, T \mapsto \text{Hom}_S(T, G).$$

Then h_G is a sheaf on X_E which we also denote as G .

Representable Sheaves

Let $E \in \{\text{Zar}, \text{ét}, \text{fppf}\}$ and consider the site X_E for a scheme X . For any commutative X -group scheme G , we can consider the functor of points

$$h_G : (\text{Sch}/X)^{\text{op}} \rightarrow \mathbf{Ab}, \quad T \mapsto \text{Hom}_S(T, G).$$

Then h_G is a sheaf on X_E which we also denote as G .

Subcategory of the Category of Sheaves

With E as above, we have by Yoneda a fully faithful embedding

$$\mathcal{Y} : \text{Sch}/X \rightarrow [(\text{Sch}/X)^{\text{op}}, \mathbf{Set}], \quad T \mapsto h_T$$

Representable Sheaves

Let $E \in \{\text{Zar}, \text{ét}, \text{fppf}\}$ and consider the site X_E for a scheme X . For any commutative X -group scheme G , we can consider the functor of points

$$h_G : (\text{Sch}/X)^{\text{op}} \rightarrow \mathbf{Ab}, \quad T \mapsto \text{Hom}_S(T, G).$$

Then h_G is a sheaf on X_E which we also denote as G .

Subcategory of the Category of Sheaves

With E as above, we have by Yoneda a fully faithful embedding

$$\mathcal{Y} : \text{Sch}/X \rightarrow [(\text{Sch}/X)^{\text{op}}, \mathbf{Set}], \quad T \mapsto h_T$$

The previous example generalises and says that the essential image of \mathcal{Y} lies in the subcategory of sheaves of sets on X_E (and is the subcategory of representable sheaves). We denote the sheaf h_T as T as well.

Representable Sheaves

Let $E \in \{\text{Zar}, \text{ét}, \text{fppf}\}$ and consider the site X_E for a scheme X . For any commutative X -group scheme G , we can consider the functor of points

$$h_G : (\text{Sch}/X)^{\text{op}} \rightarrow \mathbf{Ab}, \quad T \mapsto \text{Hom}_S(T, G).$$

Then h_G is a sheaf on X_E which we also denote as G .

Subcategory of the Category of Sheaves

With E as above, we have by Yoneda a fully faithful embedding

$$\mathcal{Y} : \text{Sch}/X \rightarrow [(\text{Sch}/X)^{\text{op}}, \text{Set}], \quad T \mapsto h_T$$

The previous example generalises and says that the essential image of \mathcal{Y} lies in the subcategory of sheaves of sets on X_E (and is the subcategory of representable sheaves). We denote the sheaf h_T as T as well.

This way, we can view Sch/X as a subcategory of $\text{Sh}(X_E)$.

Sheafification

As before, the functor $i : \mathbf{Sh}(\mathcal{T}) \hookrightarrow \mathbf{PSh}(\mathcal{T})$ has a left adjoint

$$(-)^{\mathrm{sh}} : \mathbf{PSh}(\mathcal{T}) \rightarrow \mathbf{Sh}(\mathcal{T})$$

called the sheafification functor.

Sheafification

As before, the functor $i : \text{Sh}(\mathcal{T}) \hookrightarrow \text{PSh}(\mathcal{T})$ has a left adjoint

$$(-)^{\text{sh}} : \text{PSh}(\mathcal{T}) \rightarrow \text{Sh}(\mathcal{T})$$

called the sheafification functor.

Category of Sheaves on Sites is Abelian

The category $\text{Sh}(\mathcal{T})$ is an abelian category where the kernel of a sheaf morphism is defined as before and is a sheaf, while the cokernel (and image) are again taken to be the sheafification of the obvious definitions.

Sheafification

As before, the functor $i : \mathbf{Sh}(\mathcal{T}) \hookrightarrow \mathbf{PSh}(\mathcal{T})$ has a left adjoint

$$(-)^{\mathrm{sh}} : \mathbf{PSh}(\mathcal{T}) \rightarrow \mathbf{Sh}(\mathcal{T})$$

called the sheafification functor.

Category of Sheaves on Sites is Abelian

The category $\mathbf{Sh}(\mathcal{T})$ is an abelian category where the kernel of a sheaf morphism is defined as before and is a sheaf, while the cokernel (and image) are again taken to be the sheafification of the obvious definitions.

The category $\mathbf{Sh}(\mathcal{T})$ has enough injectives, this is a consequence of it being a Grothendieck category (an AB5 category with a family of generators).

Sheaf Cohomology on Sites

Fix a scheme X and an $E \in \{\text{Zar}, \text{ét}, \text{fppf}\}$, and consider the site X_E .

Sheaf Cohomology on Sites

Fix a scheme X and an $E \in \{\text{Zar}, \text{ét}, \text{fppf}\}$, and consider the site X_E . The *global section functor*

$$\Gamma(X, -) : \text{Sh}(X_E) \rightarrow \text{Ab}$$

is left exact.

Sheaf Cohomology on Sites

Fix a scheme X and an $E \in \{\text{Zar}, \text{ét}, \text{fppf}\}$, and consider the site X_E . The *global section functor*

$$\Gamma(X, -) : \text{Sh}(X_E) \rightarrow \text{Ab}$$

is left exact. Since $\text{Sh}(X_E)$ has enough injectives, we define the i^{th} E -cohomology group of X with coefficients in \mathcal{F} as

$$H^i(X_E, \mathcal{F}) = H_E^i(X, \mathcal{F}) := R^i\Gamma(X, \mathcal{F}),$$

that is, the right derived functor of $\Gamma(X, -)$.

Sheaf Cohomology on Sites

Fix a scheme X and an $E \in \{\text{Zar}, \text{ét}, \text{fppf}\}$, and consider the site X_E . The *global section functor*

$$\Gamma(X, -) : \text{Sh}(X_E) \rightarrow \text{Ab}$$

is left exact. Since $\text{Sh}(X_E)$ has enough injectives, we define the i^{th} E -cohomology group of X with coefficients in \mathcal{F} as

$$H^i(X_E, \mathcal{F}) = H_E^i(X, \mathcal{F}) := R^i\Gamma(X, \mathcal{F}),$$

that is, the right derived functor of $\Gamma(X, -)$.

Therefore, given any short exact sequence of sheaves in $\text{Sh}(X_E)$

$$0 \longrightarrow \mathcal{F} \xrightarrow{\phi} \mathcal{G} \xrightarrow{\psi} \mathcal{H} \longrightarrow 0$$

Sheaf Cohomology on Sites

Fix a scheme X and an $E \in \{\text{Zar}, \text{ét}, \text{fppf}\}$, and consider the site X_E . The *global section functor*

$$\Gamma(X, -) : \text{Sh}(X_E) \rightarrow \text{Ab}$$

is left exact. Since $\text{Sh}(X_E)$ has enough injectives, we define the i^{th} E -cohomology group of X with coefficients in \mathcal{F} as

$$H^i(X_E, \mathcal{F}) = H_E^i(X, \mathcal{F}) := R^i\Gamma(X, \mathcal{F}),$$

that is, the right derived functor of $\Gamma(X, -)$.

Therefore, given any short exact sequence of sheaves in $\text{Sh}(X_E)$

$$0 \longrightarrow \mathcal{F} \xrightarrow{\phi} \mathcal{G} \xrightarrow{\psi} \mathcal{H} \longrightarrow 0$$

we obtain a long exact sequence of abelian groups

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Gamma_E(X, \mathcal{F}) & \longrightarrow & \Gamma_E(X, \mathcal{G}) & \longrightarrow & \Gamma_E(X, \mathcal{H}) \\ & & & & & & \searrow \\ & & & & & & H_E^1(X, \mathcal{F}) \longrightarrow H_E^1(X, \mathcal{G}) \longrightarrow H_E^1(X, \mathcal{H}) \longrightarrow H_E^2(X, \mathcal{F}) \longrightarrow \dots \end{array}$$

Hilbert's Theorem 90

Let X be any scheme, then

$$H_{\text{fppf}}^1(X, \mathbb{G}_m) \cong H_{\text{ét}}^1(X, \mathbb{G}_m) \cong H_{\text{Zar}}^1(X, \mathcal{O}_X^*) \cong \text{Pic}(X)$$

where \mathbb{G}_m is the sheaf given as $\mathbb{G}_m(U) = \Gamma(U, \mathcal{O}_U)^*$.

Hilbert's Theorem 90

Let X be any scheme, then

$$H_{\text{fppf}}^1(X, \mathbb{G}_m) \cong H_{\text{ét}}^1(X, \mathbb{G}_m) \cong H_{\text{Zar}}^1(X, \mathcal{O}_X^*) \cong \text{Pic}(X)$$

where \mathbb{G}_m is the sheaf given as $\mathbb{G}_m(U) = \Gamma(U, \mathcal{O}_U)^*$. It's a representable sheaf, represented by the scheme $\text{Spec } \mathbb{Z}[T, T^{-1}]$.

Some Examples

Hilbert's Theorem 90

Let X be any scheme, then

$$H_{\text{fppf}}^1(X, \mathbb{G}_m) \cong H_{\text{ét}}^1(X, \mathbb{G}_m) \cong H_{\text{Zar}}^1(X, \mathcal{O}_X^*) \cong \text{Pic}(X)$$

where \mathbb{G}_m is the sheaf given as $\mathbb{G}_m(U) = \Gamma(U, \mathcal{O}_U)^*$. It's a representable sheaf, represented by the scheme $\text{Spec } \mathbb{Z}[T, T^{-1}]$.

Kummer Sequence

Consider the subsheaf μ_n of \mathbb{G}_m by setting $\mu_n(U)$ as the group of n^{th} roots of unity in $\Gamma(U, \mathcal{O}_U)$.

Some Examples

Hilbert's Theorem 90

Let X be any scheme, then

$$H_{\text{fppf}}^1(X, \mathbb{G}_m) \cong H_{\text{ét}}^1(X, \mathbb{G}_m) \cong H_{\text{Zar}}^1(X, \mathcal{O}_X^*) \cong \text{Pic}(X)$$

where \mathbb{G}_m is the sheaf given as $\mathbb{G}_m(U) = \Gamma(U, \mathcal{O}_U)^*$. It's a representable sheaf, represented by the scheme $\text{Spec } \mathbb{Z}[T, T^{-1}]$.

Kummer Sequence

Consider the subsheaf μ_n of \mathbb{G}_m by setting $\mu_n(U)$ as the group of n^{th} roots of unity in $\Gamma(U, \mathcal{O}_U)$. It's a representable sheaf, represented by the scheme $\text{Spec } \mathbb{Z}[T]/(T^n - 1)$.

Some Examples

Hilbert's Theorem 90

Let X be any scheme, then

$$H_{\text{fppf}}^1(X, \mathbb{G}_m) \cong H_{\text{ét}}^1(X, \mathbb{G}_m) \cong H_{\text{Zar}}^1(X, \mathcal{O}_X^*) \cong \text{Pic}(X)$$

where \mathbb{G}_m is the sheaf given as $\mathbb{G}_m(U) = \Gamma(U, \mathcal{O}_U)^*$. It's a representable sheaf, represented by the scheme $\text{Spec } \mathbb{Z}[T, T^{-1}]$.

Kummer Sequence

Consider the subsheaf μ_n of \mathbb{G}_m by setting $\mu_n(U)$ as the group of n^{th} roots of unity in $\Gamma(U, \mathcal{O}_U)$. It's a representable sheaf, represented by the scheme $\text{Spec } \mathbb{Z}[T]/(T^n - 1)$. Then we have the exact sequence in $\text{Sh}(X_{\text{fppf}})$

$$0 \longrightarrow \mu_n \longrightarrow \mathbb{G}_m \xrightarrow{n} \mathbb{G}_m \longrightarrow 0$$

Some Examples

Hilbert's Theorem 90

Let X be any scheme, then

$$H_{\text{fppf}}^1(X, \mathbb{G}_m) \cong H_{\text{ét}}^1(X, \mathbb{G}_m) \cong H_{\text{Zar}}^1(X, \mathcal{O}_X^*) \cong \text{Pic}(X)$$

where \mathbb{G}_m is the sheaf given as $\mathbb{G}_m(U) = \Gamma(U, \mathcal{O}_U)^*$. It's a representable sheaf, represented by the scheme $\text{Spec } \mathbb{Z}[T, T^{-1}]$.

Kummer Sequence

Consider the subsheaf μ_n of \mathbb{G}_m by setting $\mu_n(U)$ as the group of n^{th} roots of unity in $\Gamma(U, \mathcal{O}_U)$. It's a representable sheaf, represented by the scheme $\text{Spec } \mathbb{Z}[T]/(T^n - 1)$. Then we have the exact sequence in $\text{Sh}(X_{\text{fppf}})$

$$0 \longrightarrow \mu_n \longrightarrow \mathbb{G}_m \xrightarrow{n} \mathbb{G}_m \longrightarrow 0$$

(this sequence is exact in $\text{Sh}(X_{\text{ét}})$ provided n is coprime to the characteristic of X .) (...)

Kummer Sequence

(...) This gives rise to an exact cohomology sequence

$$0 \longrightarrow \mathcal{M}_n(X) \longrightarrow \Gamma(X, \mathcal{O}_X)^* \xrightarrow{n} \Gamma(X, \mathcal{O}_X)^* \longrightarrow H^1(X, \mathcal{M}_n) \longrightarrow \text{Pic}(X) \xrightarrow{n} \text{Pic}(X).$$

Kummer Sequence

(...) This gives rise to an exact cohomology sequence

$$0 \longrightarrow \mathcal{H}_n(X) \longrightarrow \Gamma(X, \mathcal{O}_X)^* \xrightarrow{n} \Gamma(X, \mathcal{O}_X)^* \longrightarrow H^1(X, \mathcal{H}_n) \longrightarrow \text{Pic}(X) \xrightarrow{n} \text{Pic}(X).$$

This sequence has the following explicit interpretation:

Kummer Sequence

(...) This gives rise to an exact cohomology sequence

$$0 \longrightarrow \mathcal{H}_n(X) \longrightarrow \Gamma(X, \mathcal{O}_X)^* \xrightarrow{n} \Gamma(X, \mathcal{O}_X)^* \longrightarrow H^1(X, \mathcal{H}_n) \longrightarrow \text{Pic}(X) \xrightarrow{n} \text{Pic}(X).$$

This sequence has the following explicit interpretation:

$H^1(X, \mathcal{H}_n)$ is the set (modulo isomorphism) of pairs (\mathcal{L}, ϕ) where \mathcal{L} is a line bundle on X and ϕ is an isomorphism $\mathcal{O}_X \rightarrow \mathcal{L}^{\otimes n}$.

Kummer Sequence

(...) This gives rise to an exact cohomology sequence

$$0 \longrightarrow \mathcal{H}^1(X) \longrightarrow \Gamma(X, \mathcal{O}_X)^* \xrightarrow{n} \Gamma(X, \mathcal{O}_X)^* \longrightarrow H^1(X, \mathcal{H}^1) \longrightarrow \text{Pic}(X) \xrightarrow{n} \text{Pic}(X).$$

This sequence has the following explicit interpretation:

$H^1(X, \mathcal{H}^1)$ is the set (modulo isomorphism) of pairs (\mathcal{L}, ϕ) where \mathcal{L} is a line bundle on X and ϕ is an isomorphism $\mathcal{O}_X \rightarrow \mathcal{L}^{\otimes n}$.

The image of (\mathcal{L}, ϕ) in $\text{Pic}(X)[n] = \ker(\text{Pic}(X) \xrightarrow{n} \text{Pic}(X))$ is the class of \mathcal{L} .

Kummer Sequence

(...) This gives rise to an exact cohomology sequence

$$0 \longrightarrow \mathcal{H}_n(X) \longrightarrow \Gamma(X, \mathcal{O}_X)^* \xrightarrow{n} \Gamma(X, \mathcal{O}_X)^* \longrightarrow H^1(X, \mathcal{H}_n) \longrightarrow \text{Pic}(X) \xrightarrow{n} \text{Pic}(X).$$

This sequence has the following explicit interpretation:

$H^1(X, \mathcal{H}_n)$ is the set (modulo isomorphism) of pairs (\mathcal{L}, ϕ) where \mathcal{L} is a line bundle on X and ϕ is an isomorphism $\mathcal{O}_X \rightarrow \mathcal{L}^{\otimes n}$.

The image of (\mathcal{L}, ϕ) in $\text{Pic}(X)[n] = \ker(\text{Pic}(X) \xrightarrow{n} \text{Pic}(X))$ is the class of \mathcal{L} .

If there exists an isomorphism $\psi : \mathcal{O}_X \rightarrow \mathcal{L}$, that is, if (\mathcal{L}, ϕ) maps to zero in $\text{Pic}(X)$,

Kummer Sequence

(...) This gives rise to an exact cohomology sequence

$$0 \longrightarrow \mathcal{H}_n(X) \longrightarrow \Gamma(X, \mathcal{O}_X)^* \xrightarrow{n} \Gamma(X, \mathcal{O}_X)^* \longrightarrow H^1(X, \mathcal{H}_n) \longrightarrow \text{Pic}(X) \xrightarrow{n} \text{Pic}(X).$$

This sequence has the following explicit interpretation:

$H^1(X, \mathcal{H}_n)$ is the set (modulo isomorphism) of pairs (\mathcal{L}, ϕ) where \mathcal{L} is a line bundle on X and ϕ is an isomorphism $\mathcal{O}_X \rightarrow \mathcal{L}^{\otimes n}$.

The image of (\mathcal{L}, ϕ) in $\text{Pic}(X)[n] = \ker(\text{Pic}(X) \xrightarrow{n} \text{Pic}(X))$ is the class of \mathcal{L} .

If there exists an isomorphism $\psi : \mathcal{O}_X \rightarrow \mathcal{L}$, that is, if (\mathcal{L}, ϕ) maps to zero in $\text{Pic}(X)$, then (\mathcal{L}, ϕ) is the image of $a \in \Gamma(X, \mathcal{O}_X)^*$,

Kummer Sequence

(...) This gives rise to an exact cohomology sequence

$$0 \longrightarrow \mathcal{H}^1(X) \longrightarrow \Gamma(X, \mathcal{O}_X)^* \xrightarrow{n} \Gamma(X, \mathcal{O}_X)^* \longrightarrow H^1(X, \mathcal{H}^1) \longrightarrow \text{Pic}(X) \xrightarrow{n} \text{Pic}(X).$$

This sequence has the following explicit interpretation:

$H^1(X, \mathcal{H}^1)$ is the set (modulo isomorphism) of pairs (\mathcal{L}, ϕ) where \mathcal{L} is a line bundle on X and ϕ is an isomorphism $\mathcal{O}_X \rightarrow \mathcal{L}^{\otimes n}$.

The image of (\mathcal{L}, ϕ) in $\text{Pic}(X)[n] = \ker(\text{Pic}(X) \xrightarrow{n} \text{Pic}(X))$ is the class of \mathcal{L} .

If there exists an isomorphism $\psi : \mathcal{O}_X \rightarrow \mathcal{L}$, that is, if (\mathcal{L}, ϕ) maps to zero in $\text{Pic}(X)$, then (\mathcal{L}, ϕ) is the image of $a \in \Gamma(X, \mathcal{O}_X)^*$, where

$$\mathcal{O}_X \xrightarrow{\phi} \mathcal{L}^{\otimes n} \xrightarrow{(\psi^{-1})^{\otimes n}} \mathcal{O}_X^{\otimes n} \xrightarrow{\text{can}} \mathcal{O}_X$$

is multiplication by a^{-1} .

Étale Cohomology of $\text{Spec } k$

Fix a field k and a separable closure k_{sep} , denote by $G_k := \text{Gal}(k_{\text{sep}}/k)$ the absolute Galois group of k .

Étale Cohomology of $\mathrm{Spec} k$

Fix a field k and a separable closure k_{sep} , denote by $G_k := \mathrm{Gal}(k_{\mathrm{sep}}/k)$ the absolute Galois group of k .

Let \mathcal{F} be a sheaf on $(\mathrm{Spec} k)_{\mathrm{\acute{e}t}}$,

Étale Cohomology of $\mathrm{Spec} k$

Fix a field k and a separable closure k_{sep} , denote by $G_k := \mathrm{Gal}(k_{\mathrm{sep}}/k)$ the absolute Galois group of k .

Let \mathcal{F} be a sheaf on $(\mathrm{Spec} k)_{\mathrm{\acute{e}t}}$, define

$$M_{\mathcal{F}} := \varinjlim \mathcal{F}(\mathrm{Spec} L),$$

where L are finite extensions of k contained in k_{sep} (that is, they're separable).

Étale Cohomology of $\mathrm{Spec} k$

Fix a field k and a separable closure k_{sep} , denote by $G_k := \mathrm{Gal}(k_{\mathrm{sep}}/k)$ the absolute Galois group of k .

Let \mathcal{F} be a sheaf on $(\mathrm{Spec} k)_{\mathrm{\acute{e}t}}$, define

$$M_{\mathcal{F}} := \varinjlim \mathcal{F}(\mathrm{Spec} L),$$

where L are finite extensions of k contained in k_{sep} (that is, they're separable). Since G_k acts on L (say as $\alpha_g : L \rightarrow L$), it acts on $\mathcal{F}(\mathrm{Spec} L)$ in the following manner

$$\mathcal{F}(\mathrm{Spec} \alpha_g) : \mathcal{F}(\mathrm{Spec} L) \rightarrow \mathcal{F}(\mathrm{Spec} L),$$

Étale Cohomology of $\mathrm{Spec} k$

Fix a field k and a separable closure k_{sep} , denote by $G_k := \mathrm{Gal}(k_{\mathrm{sep}}/k)$ the absolute Galois group of k .

Let \mathcal{F} be a sheaf on $(\mathrm{Spec} k)_{\mathrm{\acute{e}t}}$, define

$$M_{\mathcal{F}} := \varinjlim \mathcal{F}(\mathrm{Spec} L),$$

where L are finite extensions of k contained in k_{sep} (that is, they're separable). Since G_k acts on L (say as $\alpha_g : L \rightarrow L$), it acts on $\mathcal{F}(\mathrm{Spec} L)$ in the following manner

$$\mathcal{F}(\mathrm{Spec} \alpha_g) : \mathcal{F}(\mathrm{Spec} L) \rightarrow \mathcal{F}(\mathrm{Spec} L),$$

and therefore acts on $M_{\mathcal{F}}$ making it a G_k -module.

Étale Cohomology of $\mathrm{Spec} k$

Fix a field k and a separable closure k_{sep} , denote by $G_k := \mathrm{Gal}(k_{\mathrm{sep}}/k)$ the absolute Galois group of k .

Let \mathcal{F} be a sheaf on $(\mathrm{Spec} k)_{\mathrm{\acute{e}t}}$, define

$$M_{\mathcal{F}} := \varinjlim \mathcal{F}(\mathrm{Spec} L),$$

where L are finite extensions of k contained in k_{sep} (that is, they're separable). Since G_k acts on L (say as $\alpha_g : L \rightarrow L$), it acts on $\mathcal{F}(\mathrm{Spec} L)$ in the following manner

$$\mathcal{F}(\mathrm{Spec} \alpha_g) : \mathcal{F}(\mathrm{Spec} L) \rightarrow \mathcal{F}(\mathrm{Spec} L),$$

and therefore acts on $M_{\mathcal{F}}$ making it a G_k -module. In fact, we have a functor

$$\mathrm{Sh}_{\mathrm{\acute{e}t}}(\mathrm{Spec} k) \rightarrow G_k\text{-Mod} : \mathcal{F} \mapsto M_{\mathcal{F}}$$

Étale Cohomology of $\mathrm{Spec} k$

Fix a field k and a separable closure k_{sep} , denote by $G_k := \mathrm{Gal}(k_{\mathrm{sep}}/k)$ the absolute Galois group of k .

Let \mathcal{F} be a sheaf on $(\mathrm{Spec} k)_{\mathrm{\acute{e}t}}$, define

$$M_{\mathcal{F}} := \varinjlim \mathcal{F}(\mathrm{Spec} L),$$

where L are finite extensions of k contained in k_{sep} (that is, they're separable). Since G_k acts on L (say as $\alpha_g : L \rightarrow L$), it acts on $\mathcal{F}(\mathrm{Spec} L)$ in the following manner

$$\mathcal{F}(\mathrm{Spec} \alpha_g) : \mathcal{F}(\mathrm{Spec} L) \rightarrow \mathcal{F}(\mathrm{Spec} L),$$

and therefore acts on $M_{\mathcal{F}}$ making it a G_k -module. In fact, we have a functor

$$\mathrm{Sh}_{\mathrm{\acute{e}t}}(\mathrm{Spec} k) \rightarrow G_k\text{-Mod} : \mathcal{F} \mapsto M_{\mathcal{F}}$$

The above functor is an equivalence of categories,

Étale Cohomology of $\mathrm{Spec} k$

Fix a field k and a separable closure k_{sep} , denote by $G_k := \mathrm{Gal}(k_{\mathrm{sep}}/k)$ the absolute Galois group of k .

Let \mathcal{F} be a sheaf on $(\mathrm{Spec} k)_{\mathrm{\acute{e}t}}$, define

$$M_{\mathcal{F}} := \varinjlim \mathcal{F}(\mathrm{Spec} L),$$

where L are finite extensions of k contained in k_{sep} (that is, they're separable). Since G_k acts on L (say as $\alpha_g : L \rightarrow L$), it acts on $\mathcal{F}(\mathrm{Spec} L)$ in the following manner

$$\mathcal{F}(\mathrm{Spec} \alpha_g) : \mathcal{F}(\mathrm{Spec} L) \rightarrow \mathcal{F}(\mathrm{Spec} L),$$

and therefore acts on $M_{\mathcal{F}}$ making it a G_k -module. In fact, we have a functor

$$\mathrm{Sh}_{\mathrm{\acute{e}t}}(\mathrm{Spec} k) \rightarrow G_k\text{-Mod} : \mathcal{F} \mapsto M_{\mathcal{F}}$$

The above functor is an equivalence of categories, under this identification, taking global sections $\mathcal{F} \mapsto \Gamma(\mathrm{Spec} k, \mathcal{F})$ corresponds to the fixed points functor $M \mapsto M^{G_k}$.

Étale Cohomology of $\text{Spec } k$

Fix a field k and a separable closure k_{sep} , denote by $G_k := \text{Gal}(k_{\text{sep}}/k)$ the absolute Galois group of k .

Let \mathcal{F} be a sheaf on $(\text{Spec } k)_{\text{ét}}$, define

$$M_{\mathcal{F}} := \varinjlim \mathcal{F}(\text{Spec } L),$$

where L are finite extensions of k contained in k_{sep} (that is, they're separable). Since G_k acts on L (say as $\alpha_g : L \rightarrow L$), it acts on $\mathcal{F}(\text{Spec } L)$ in the following manner

$$\mathcal{F}(\text{Spec } \alpha_g) : \mathcal{F}(\text{Spec } L) \rightarrow \mathcal{F}(\text{Spec } L),$$

and therefore acts on $M_{\mathcal{F}}$ making it a G_k -module. In fact, we have a functor

$$\text{Sh}_{\text{ét}}(\text{Spec } k) \rightarrow G_k\text{-Mod} : \mathcal{F} \mapsto M_{\mathcal{F}}$$

The above functor is an equivalence of categories, under this identification, taking global sections $\mathcal{F} \mapsto \Gamma(\text{Spec } k, \mathcal{F})$ corresponds to the fixed points functor $M \mapsto M^{G_k}$. This ends up telling us that

$$H_{\text{ét}}^i(\text{Spec } k, \mathcal{F}) \cong H^i(G_k, M_{\mathcal{F}}),$$

where the latter is the i^{th} Galois cohomology group (that is, group cohomology of G_k) with coefficients in $M_{\mathcal{F}}$.

(...)

Étale Cohomology of $\operatorname{Spec} k$

(...) In particular if $\mathcal{F} = \mathbb{G}_m$, then

$$H_{\text{ét}}^i(\operatorname{Spec} k, \mathbb{G}_m) \cong H^i(G_k, k_{\text{sep}}^\times)$$

Étale Cohomology of $\operatorname{Spec} k$

(...) In particular if $\mathcal{F} = \mathbb{G}_m$, then

$$H_{\text{ét}}^i(\operatorname{Spec} k, \mathbb{G}_m) \cong H^i(G_k, k_{\text{sep}}^\times)$$

The classical Hilbert's Theorem 90 tells us that the latter cohomology group is trivial, while the one stated above tells us that the former cohomology group is $\operatorname{Pic}(\operatorname{Spec} k)$. This is trivial, since the only line bundles on $\operatorname{Spec} k$ are the trivial ones.

Étale Cohomology of $\operatorname{Spec} k$

(...) In particular if $\mathcal{F} = \mathbb{G}_m$, then

$$H_{\text{ét}}^i(\operatorname{Spec} k, \mathbb{G}_m) \cong H^i(G_k, k_{\text{sep}}^\times)$$

The classical Hilbert's Theorem 90 tells us that the latter cohomology group is trivial, while the one stated above tells us that the former cohomology group is $\operatorname{Pic}(\operatorname{Spec} k)$. This is trivial, since the only line bundles on $\operatorname{Spec} k$ are the trivial ones.

Brauer Groups

Since we have $H^2(G_k, k_{\text{sep}}^\times) \cong \operatorname{Br}(k)$, the Brauer group of k which can be defined as the group of Morita equivalent central simple algebras over k ,

Étale Cohomology of $\mathrm{Spec} k$

(...) In particular if $\mathcal{F} = \mathbb{G}_m$, then

$$H_{\text{ét}}^i(\mathrm{Spec} k, \mathbb{G}_m) \cong H^i(G_k, k_{\text{sep}}^\times)$$

The classical Hilbert's Theorem 90 tells us that the latter cohomology group is trivial, while the one stated above tells us that the former cohomology group is $\mathrm{Pic}(\mathrm{Spec} k)$. This is trivial, since the only line bundles on $\mathrm{Spec} k$ are the trivial ones.

Brauer Groups

Since we have $H^2(G_k, k_{\text{sep}}^\times) \cong \mathrm{Br}(k)$, the Brauer group of k which can be defined as the group of Morita equivalent central simple algebras over k , we call, for any scheme X , the group $H_{\text{ét}}^2(X, \mathbb{G}_m)$ the *cohomological Brauer group* of X .

Étale Cohomology of $\operatorname{Spec} k$

(...) In particular if $\mathcal{F} = \mathbb{G}_m$, then

$$H_{\text{ét}}^i(\operatorname{Spec} k, \mathbb{G}_m) \cong H^i(G_k, k_{\text{sep}}^\times)$$

The classical Hilbert's Theorem 90 tells us that the latter cohomology group is trivial, while the one stated above tells us that the former cohomology group is $\operatorname{Pic}(\operatorname{Spec} k)$. This is trivial, since the only line bundles on $\operatorname{Spec} k$ are the trivial ones.

Brauer Groups

Since we have $H^2(G_k, k_{\text{sep}}^\times) \cong \operatorname{Br}(k)$, the Brauer group of k which can be defined as the group of Morita equivalent central simple algebras over k , we call, for any scheme X , the group $H_{\text{ét}}^2(X, \mathbb{G}_m)$ the *cohomological Brauer group* of X .

The above group was denoted by Grothendieck as $\operatorname{Br}'(X)$.

Étale Cohomology of $\operatorname{Spec} k$

(...) In particular if $\mathcal{F} = \mathbb{G}_m$, then

$$H_{\text{ét}}^i(\operatorname{Spec} k, \mathbb{G}_m) \cong H^i(G_k, k_{\text{sep}}^\times)$$

The classical Hilbert's Theorem 90 tells us that the latter cohomology group is trivial, while the one stated above tells us that the former cohomology group is $\operatorname{Pic}(\operatorname{Spec} k)$. This is trivial, since the only line bundles on $\operatorname{Spec} k$ are the trivial ones.

Brauer Groups

Since we have $H^2(G_k, k_{\text{sep}}^\times) \cong \operatorname{Br}(k)$, the Brauer group of k which can be defined as the group of Morita equivalent central simple algebras over k , we call, for any scheme X , the group $H_{\text{ét}}^2(X, \mathbb{G}_m)$ the *cohomological Brauer group* of X .

The above group was denoted by Grothendieck as $\operatorname{Br}'(X)$. He reserved $\operatorname{Br}(X)$ for the group of Morita equivalent *Azumaya algebras* (a generalisation of central simple algebras).

Étale Cohomology of $\operatorname{Spec} k$

(...) In particular if $\mathcal{F} = \mathbb{G}_m$, then

$$H_{\text{ét}}^i(\operatorname{Spec} k, \mathbb{G}_m) \cong H^i(G_k, k_{\text{sep}}^\times)$$

The classical Hilbert's Theorem 90 tells us that the latter cohomology group is trivial, while the one stated above tells us that the former cohomology group is $\operatorname{Pic}(\operatorname{Spec} k)$. This is trivial, since the only line bundles on $\operatorname{Spec} k$ are the trivial ones.

Brauer Groups

Since we have $H^2(G_k, k_{\text{sep}}^\times) \cong \operatorname{Br}(k)$, the Brauer group of k which can be defined as the group of Morita equivalent central simple algebras over k , we call, for any scheme X , the group $H_{\text{ét}}^2(X, \mathbb{G}_m)$ the *cohomological Brauer group* of X .

The above group was denoted by Grothendieck as $\operatorname{Br}'(X)$. He reserved $\operatorname{Br}(X)$ for the group of Morita equivalent *Azumaya algebras* (a generalisation of central simple algebras). One could mean either groups when one says the Brauer group of a scheme.

Étale Cohomology of $\operatorname{Spec} k$

(...) In particular if $\mathcal{F} = \mathbb{G}_m$, then

$$H_{\text{ét}}^i(\operatorname{Spec} k, \mathbb{G}_m) \cong H^i(G_k, k_{\text{sep}}^\times)$$

The classical Hilbert's Theorem 90 tells us that the latter cohomology group is trivial, while the one stated above tells us that the former cohomology group is $\operatorname{Pic}(\operatorname{Spec} k)$. This is trivial, since the only line bundles on $\operatorname{Spec} k$ are the trivial ones.

Brauer Groups

Since we have $H^2(G_k, k_{\text{sep}}^\times) \cong \operatorname{Br}(k)$, the Brauer group of k which can be defined as the group of Morita equivalent central simple algebras over k , we call, for any scheme X , the group $H_{\text{ét}}^2(X, \mathbb{G}_m)$ the *cohomological Brauer group* of X .

The above group was denoted by Grothendieck as $\operatorname{Br}'(X)$. He reserved $\operatorname{Br}(X)$ for the group of Morita equivalent *Azumaya algebras* (a generalisation of central simple algebras). One could mean either groups when one says the Brauer group of a scheme. For some nice schemes (an affine scheme, for example), we have $\operatorname{Br}(X) \cong \operatorname{Br}'(X)_{\text{tors}}$.

Fin.

- [1] Artin, M. (1962), *Grothendieck Topologies. Notes on a Seminar by M. Artin. Spring, 1962*. Harvard University, Department of Mathematics.
- [2] Balmer, P. (2013), *Stacks of group representations*. arXiv:1302.6290.
- [3] Griffiths, P. & Harris, J. (1978), *Principles of Algebraic Geometry*. Wiley.
- [4] Milne, J.S. (1980), *Étale Cohomology (PMS-33)*. Princeton University Press.
- [5] Tamme, G. (1994), *Introduction to Étale Cohomology*. Springer-Verlag.
- [6] Vakil, R., *The Rising Sea: Foundations Of Algebraic Geometry Notes*. November 18, 2017 version.