

# Asymptotic bounds for the combinatorial diameter of random polytopes


**Sophie Huiberts** (Columbia University)

joint work with Gilles Bonnet, Daniel Dadush, Uri Grupel, Galyna Livshyts

Cargese 2022


<https://sophie.huiberts.me>

# Outline




# Dramatis Personae


Diameter



Random polytopes




Polar duality



# Diameter: a Question About Geometry


Given a polyhedron, how many edges do we need to traverse to go from any vertex to any other?



- ▶  $P \subset \mathbb{R}^n$
- ▶  $m$  facets

# Diameter: a Question About Optimization

Given a polyhedron, how many edges do we need to traverse to go from any vertex to any other?



- ▶  $P \subset \mathbb{R}^n$
- ▶  $m$  facets

Given a vertex of the set of points  $x$  satisfying

$$Ax \leq b,$$

how many steps do you need to find a vertex maximizing  $c^T x$ ?

- ▶  $n$  variables
- ▶  $m$  constraints

## Diameter Backstory: The Hirsch conjecture

If  $P \subset \mathbb{R}^n$  has  $m$  facets, is the diameter at most  $m - n$ ?

# Diameter Backstory: The Hirsch conjecture

If  $P \subset \mathbb{R}^n$  has  $m$  facets, is the diameter at most  $m - n$ ?

True for

- ▶ polytopes in  $\mathbb{R}^3$
- ▶ network flow polytopes
- ▶ fractional stable set polytopes
- ▶ polytopes with vertices in  $\{0, 1\}^n$
- ▶ and many more

But false in general.

# Diameter Backstory: The Hirsch conjecture

If  $P \subset \mathbb{R}^n$  has  $m$  facets, is the diameter at most  $m - n$ ?

True for

- ▶ polytopes in  $\mathbb{R}^3$
- ▶ network flow polytopes
- ▶ fractional stable set polytopes
- ▶ polytopes with vertices in  $\{0, 1\}^n$
- ▶ and many more

But false in general.

B69, L70, B74:

$$\text{Diameter}(P) \leq 2^{n-2}m.$$

KK92, T14, S19:

$$\text{Diameter}(P) \leq (m - n)^{\log_2 O(n/\log n)}.$$



## Polar Duality

Given a convex set  $P \subset \mathbb{R}^n$ , define

$$P^\circ = \{x \in \mathbb{R}^n : y^\top x \leq 1 \quad \forall y \in P\}.$$


# Polar Duality

Given a convex set  $P \subset \mathbb{R}^n$ , define

$$P^\circ = \{x \in \mathbb{R}^n : y^\top x \leq 1 \quad \forall y \in P\}.$$

If that  $P$  is a bounded polytope and contains the origin in its interior, then:

- ▶  $P^\circ$  is a bounded polytope and contains the origin in its interior
- ▶  $(P^\circ)^\circ = P$
- ▶ if  $P$  is simple then  $P^\circ$  is simplicial
- ▶ vertices of  $P$  correspond to facets of  $P^\circ$



# Random Models

Let

$$P = \{x \in \mathbb{R}^n : \langle a, x \rangle \leq 1 \quad \forall a \in A\}, \quad Q = \text{conv}(A),$$

where  $A \subset \mathbb{S}^{n-1}$ ,  $|A| \sim \text{Poisson}(m)$  is sampled iid.




Figure:  $P \subset \mathbb{R}^3$  might look like this

## Call to Adventure

DF94, BdSEHN14, DH16, NSS22:

If  $S = \{x \in \mathbb{R}^n : Ax \leq b\}$  where  $A \in \mathbb{Z}^{n \times m}$  and every absolute square subdeterminant of  $A$  is at most  $\Delta$  then


$$\text{Diameter}(S) \leq O(n^3 \Delta^2 \log(\Delta)).$$

# Call to Adventure

DF94, BdSEHN14, DH16, NSS22:

If  $S = \{x \in \mathbb{R}^n : Ax \leq b\}$  where  $A \in \mathbb{Z}^{n \times m}$  and every absolute square subdeterminant of  $A$  is at most  $\Delta$  then

$$\text{Diameter}(S) \leq O(n^3 \Delta^2 \log(\Delta)).$$



Borgwardt 1977,1982,1987,1999:


consider the projection  $\pi$  onto a fixed two-dimensional subspace  $W \subset \mathbb{R}^n$ . In expectation, the random polygon  $\pi(P)$  has  $\Theta(n^2 m^{\frac{1}{n-1}})$  vertices.

# Call to Adventure

DF94, BdSEHN14, DH16, NSS22:

If  $S = \{x \in \mathbb{R}^n : Ax \leq b\}$  where  $A \in \mathbb{Z}^{n \times m}$  and every absolute square subdeterminant of  $A$  is at most  $\Delta$  then

$$\text{Diameter}(S) \leq O(n^3 \Delta^2 \log(\Delta)).$$



Borgwardt 1977,1982,1987,1999:

consider the projection  $\pi$  onto a fixed two-dimensional subspace  $W \subset \mathbb{R}^n$ . In expectation, the random polygon  $\pi(P)$  has  $\Theta(n^2 m^{\frac{1}{n-1}})$  vertices.

In particular, any two vertices that project onto the boundary of  $\pi(P)$  are connected by a path along the boundary, so of length  $O(n^2 m^{\frac{1}{n-1}})$ .

## Quest: Prove the Following Results

When  $A \subset \mathbb{S}^{n-1}$  follows a Poisson point process with  $\mathbb{E}[|A|] = m > 2^{\Omega(n)}$ , then for

$$P = \{x \in \mathbb{R}^n : \langle a, x \rangle \leq 1 \quad \forall a \in A\}, \quad Q = \text{conv}(A),$$

with probability  $1 - O(1/m)$  we have

$$\Omega(nm^{\frac{1}{n-1}}) \leq \text{diam}(P) \leq O(n^2 m^{\frac{1}{n-1}} + 17^n)$$

$$\Omega(m^{\frac{1}{n-1}}) \leq \text{diam}(Q) \leq O(nm^{\frac{1}{n-1}} + 17^n).$$

## Mentor: Relating the Diameters

Let  $P \subset \mathbb{R}^n$  be a bounded simple polytope containing the origin in its interior. Let  $P^\circ = \{x \in \mathbb{R}^n : y^\top x \leq 1 \ \forall y \in P\}$ .

Lemma: We have  $\text{diam}(P) \geq (n - 1)\text{diam}(P^\circ) - 2$ .



## Mentor: Relating the Diameters

Let  $P \subset \mathbb{R}^n$  be a bounded simple polytope containing the origin in its interior. Let  $P^\circ = \{x \in \mathbb{R}^n : y^\top x \leq 1 \ \forall y \in P\}$ .

Lemma: We have  $\text{diam}(P) \geq (n - 1)\text{diam}(P^\circ) - 2$ .

Proof: consider two vertices of  $P^\circ$ , which correspond to two facets  $F, F'$  of  $P$ . Pick a path  $v_1, v_2, \dots, v_k \in P$  of vertices, with length  $k - 1 \leq \text{diam}(P)$  and  $v_1 \in F, v_k \in F'$ .


## Mentor: Relating the Diameters

Let  $P \subset \mathbb{R}^n$  be a bounded simple polytope containing the origin in its interior. Let  $P^\circ = \{x \in \mathbb{R}^n : y^\top x \leq 1 \ \forall y \in P\}$ .

Lemma: We have  $\text{diam}(P) \geq (n - 1)\text{diam}(P^\circ) - 2$ .

Proof: consider two vertices of  $P^\circ$ , which correspond to two facets  $F, F'$  of  $P$ . Pick a path  $v_1, v_2, \dots, v_k \in P$  of vertices, with length  $k - 1 \leq \text{diam}(P)$  and  $v_1 \in F, v_k \in F'$ .

Construct a sequence  $F_1, \dots, F_p$  of facets of  $P$ , each consecutive pair having non-empty intersection, such that  $v_1 \in F_1, v_k \in F_p$  and  $v_2, \dots, v_{k-1} \in F_1 \cup \dots \cup F_p$ . We can get  $p \leq (k - 1)/(n - 1)$ .




## Mentor: Relating the Diameters

Let  $P \subset \mathbb{R}^n$  be a bounded simple polytope containing the origin in its interior. Let  $P^\circ = \{x \in \mathbb{R}^n : y^\top x \leq 1 \ \forall y \in P\}$ .

Lemma: We have  $\text{diam}(P) \geq (n - 1)\text{diam}(P^\circ) - 2$ .

Proof: consider two vertices of  $P^\circ$ , which correspond to two facets  $F, F'$  of  $P$ . Pick a path  $v_1, v_2, \dots, v_k \in P$  of vertices, with length  $k - 1 \leq \text{diam}(P)$  and  $v_1 \in F, v_k \in F'$ .

Construct a sequence  $F_1, \dots, F_p$  of facets of  $P$ , each consecutive pair having non-empty intersection, such that  $v_1 \in F_1, v_k \in F_p$  and  $v_2, \dots, v_{k-1} \in F_1 \cup \dots \cup F_p$ . We can get  $p \leq (k - 1)/(n - 1)$ .



Since  $P$  is simple, any two facets with non-empty intersection share a  $(n - 2)$ -dimensional face, meaning that the corresponding vertices of  $P^\circ$  are adjacent.


## Mentor: Relating the Diameters

Let  $P \subset \mathbb{R}^n$  be a bounded simple polytope containing the origin in its interior. Let  $P^\circ = \{x \in \mathbb{R}^n : y^\top x \leq 1 \ \forall y \in P\}$ .

Lemma: We have  $\text{diam}(P) \geq (n-1)\text{diam}(P^\circ) - 2$ .

Proof: consider two vertices of  $P^\circ$ , which correspond to two facets  $F, F'$  of  $P$ . Pick a path  $v_1, v_2, \dots, v_k \in P$  of vertices, with length  $k-1 \leq \text{diam}(P)$  and  $v_1 \in F, v_k \in F'$ .

Construct a sequence  $F_1, \dots, F_p$  of facets of  $P$ , each consecutive pair having non-empty intersection, such that  $v_1 \in F_1, v_k \in F_p$  and  $v_2, \dots, v_{k-1} \in F_1 \cup \dots \cup F_p$ . We can get  $p \leq (k-1)/(n-1)$ .



Since  $P$  is simple, any two facets with non-empty intersection share a  $(n-2)$ -dimensional face, meaning that the corresponding vertices of  $P^\circ$  are adjacent.

Hence the vertices of  $P^\circ$  corresponding to  $F, F_1, \dots, F_p, F'$  form a path of length at most

$$p+2 \leq (k-1)/(n-1) + 2 \leq \text{diam}(P)/(n-1) + 2.$$

## Revelation: A Simple Lower Bound

A set  $B \subset \mathbb{S}^{n-1}$  is called  $\varepsilon$ -dense if every spherical cap of radius  $\varepsilon$  contains a point from  $B$ .

## Revelation: A Simple Lower Bound

A set  $B \subset \mathbb{S}^{n-1}$  is called  $\varepsilon$ -dense if every spherical cap of radius  $\varepsilon$  contains a point from  $B$ .

Our set  $A$  is  $\varepsilon = O((\log(m)/m)^{\frac{1}{n-1}})$ -dense with probability  $1 - n^{-n}m^{-2}$ .

## Revelation: A Simple Lower Bound

A set  $B \subset \mathbb{S}^{n-1}$  is called  $\varepsilon$ -dense if every spherical cap of radius  $\varepsilon$  contains a point from  $B$ .

Our set  $A$  is  $\varepsilon = O((\log(m)/m)^{\frac{1}{n-1}})$ -dense with probability  $1 - n^{-n}m^{-2}$ .

Every facet of  $Q$  is encapsulated in a spherical cap of radius  $\varepsilon$ .

## Revelation: A Simple Lower Bound

A set  $B \subset \mathbb{S}^{n-1}$  is called  $\varepsilon$ -dense if every spherical cap of radius  $\varepsilon$  contains a point from  $B$ .

Our set  $A$  is  $\varepsilon = O((\log(m)/m)^{\frac{1}{n-1}})$ -dense with probability  $1 - n^{-n}m^{-2}$ .

Every facet of  $Q$  is encapsulated in a spherical cap of radius  $\varepsilon$ .

Every edge of  $Q$  has length at most  $2\varepsilon$ , but  $Q$  has vertices at distance  $\geq 1$  from each other.



## Revelation: A Simple Lower Bound

A set  $B \subset \mathbb{S}^{n-1}$  is called  $\varepsilon$ -dense if every spherical cap of radius  $\varepsilon$  contains a point from  $B$ .

Our set  $A$  is  $\varepsilon = O((\log(m)/m)^{\frac{1}{n-1}})$ -dense with probability  $1 - n^{-n}m^{-2}$ .

Every facet of  $Q$  is encapsulated in a spherical cap of radius  $\varepsilon$ .

Every edge of  $Q$  has length at most  $2\varepsilon$ , but  $Q$  has vertices at distance  $\geq 1$  from each other.

With high probability  $\text{Diameter}(Q) \geq 1/2\varepsilon = \Omega(m/\log m)^{\frac{1}{n-1}}$ .

## Transformation: Sketch of the Upper Bound

- ▶ Since every facet of  $Q$  is encapsulated in a spherical cap of radius  $\varepsilon$  whp, events taking place at distance  $20\varepsilon$  from each other are nearly independent.

## Transformation: Sketch of the Upper Bound

- ▶ Since every facet of  $Q$  is encapsulated in a spherical cap of radius  $\varepsilon$  whp, events taking place at distance  $20\varepsilon$  from each other are nearly independent.
- ▶ Borgwardt's result tells us the expected number of vertices of  $\pi(P)$  in each arc segment of length  $2\varepsilon$ .

## Transformation: Sketch of the Upper Bound

- ▶ Since every facet of  $Q$  is encapsulated in a spherical cap of radius  $\varepsilon$  whp, events taking place at distance  $20\varepsilon$  from each other are nearly independent.
- ▶ Borgwardt's result tells us the expected number of vertices of  $\pi(P)$  in each arc segment of length  $2\varepsilon$ .
- ▶ Independence + upper and lower bounds on expectation  $\implies$  concentration

## Transformation: Sketch of the Upper Bound

- ▶ Since every facet of  $Q$  is encapsulated in a spherical cap of radius  $\varepsilon$  whp, events taking place at distance  $20\varepsilon$  from each other are nearly independent.
- ▶ Borgwardt's result tells us the expected number of vertices of  $\pi(P)$  in each arc segment of length  $2\varepsilon$ .
- ▶ Independence + upper and lower bounds on expectation  $\implies$  concentration
- ▶ The polygon  $\pi(P)$  has  $O(n^2 m^{\frac{1}{n-1}} + 2^{O(n)})$  vertices with probability  $1 - n^{-n} m^{-2}$ .

## Transformation: Sketch of the Upper Bound

- ▶ Since every facet of  $Q$  is encapsulated in a spherical cap of radius  $\varepsilon$  whp, events taking place at distance  $20\varepsilon$  from each other are nearly independent.
- ▶ Borgwardt's result tells us the expected number of vertices of  $\pi(P)$  in each arc segment of length  $2\varepsilon$ .
- ▶ Independence + upper and lower bounds on expectation  $\implies$  concentration
- ▶ The polygon  $\pi(P)$  has  $O(n^2 m^{\frac{1}{n-1}} + 2^{O(n)})$  vertices with probability  $1 - n^{-n} m^{-2}$ .
- ▶ Taking a union bound over many different projections  $\pi$ , this can be extended to a diameter bound.

## Atonement: Sketch of the True Lower Bound

- ▶ Every path along  $Q$  can be described by a curve of objectives

# Atonement: Sketch of the True Lower Bound

- ▶ Every path along  $Q$  can be described by a curve of objectives
- ▶ Each such curve connecting antipodal points meets  $\Omega(m^{\frac{1}{n-1}})$  distinct vertices of  $Q$  with high probability.



## Atonement: Sketch of the True Lower Bound

- ▶ Every path along  $Q$  can be described by a curve of objectives
- ▶ Each such curve connecting antipodal points meets  $\Omega(m^{\frac{1}{n-1}})$  distinct vertices of  $Q$  with high probability.
- ▶ Do the rest of the proof.

# Return

- ▶ Close the gap between upper and lower bound.
- ▶ Other distributions, such as Gaussian?
- ▶ What if  $m \leq \text{poly}(n)$ ?