

Asymptotic bounds for the combinatorial diameter of random polytopes

Sophie Huiberts (CWI → Columbia University)


joint work with Gilles Bonnet, Daniel Dadush, Uri Grupel, Galyna Livshyts

CGWeek 2022

<https://sophie.huiberts.me>

A question about geometry


Given a polyhedron, how many edges do we need to traverse to go from any vertex to any other?



- ▶ $P \subset \mathbb{R}^n$
- ▶ m facets

A question about geometry


Given a polyhedron, how many edges do we need to traverse to go from any vertex to any other?



- ▶ $P \subset \mathbb{R}^n$
- ▶ m facets

A question about optimization

Given a polyhedron, how many edges do we need to traverse to go from any vertex to any other?



- ▶ $P \subset \mathbb{R}^n$
- ▶ m facets

Given $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$, $c \in \mathbb{R}^n$ and a vertex of the polyhedron

$$\{x \in \mathbb{R}^n \mid Ax \leq b\},$$

how many steps do you need to find a vertex maximizing $c^T x$?

- ▶ n variables
- ▶ m constraints

The polynomial Hirsch conjecture

If $P \subset \mathbb{R}^n$ has m facets, how large can the diameter be?

The polynomial Hirsch conjecture

If $P \subset \mathbb{R}^n$ has m facets, how large can the diameter be?

Conjecture: at most $\text{poly}(n, m)$.

The polynomial Hirsch conjecture

If $P \subset \mathbb{R}^n$ has m facets, how large can the diameter be?

Conjecture: at most $\text{poly}(n, m)$.

At most $m - n$ for:

- ▶ polytopes in \mathbb{R}^3
- ▶ network flow polytopes
- ▶ fractional stable set polytopes
- ▶ polytopes with vertices in $\{0, 1\}^n$
- ▶ and many more

The polynomial Hirsch conjecture

If $P \subset \mathbb{R}^n$ has m facets, how large can the diameter be?

Conjecture: at most $\text{poly}(n, m)$.

At most $m - n$ for:

- ▶ polytopes in \mathbb{R}^3
- ▶ network flow polytopes
- ▶ fractional stable set polytopes
- ▶ polytopes with vertices in $\{0, 1\}^n$
- ▶ and many more

Best constructions:

KW67, unbounded polyhedra with diameter $\geq 1.25(m - n)$.

S12, bounded polyhedra with diameter $\geq 1.05(m - n)$.

Best available diameter bounds

B69, L70, B74:

$$\text{Diameter}(P) \leq 2^{n-2}m.$$

KK92, T14, S19:

$$\text{Diameter}(P) \leq (m - n)^{\log_2 O(n/\log n)}.$$

Best available diameter bounds

B69, L70, B74:

$$\text{Diameter}(P) \leq 2^{n-2}m.$$

KK92, T14, S19:

$$\text{Diameter}(P) \leq (m - n)^{\log_2 O(n/\log n)}.$$

DF94, BdSEHN14, DH16, NSS22:

If $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ where $A \in \mathbb{Z}^{n \times m}$ and every absolute square subdeterminant of A is at most Δ then

$$\text{Diameter}(P) \leq O(n^3 \Delta^2 \log(\Delta)).$$

Best available diameter bounds

B69, L70, B74:

$$\text{Diameter}(P) \leq 2^{n-2}m.$$

KK92, T14, S19:

$$\text{Diameter}(P) \leq (m - n)^{\log_2 O(n/\log n)}.$$

DF94, BdSEHN14, DH16, NSS22:

If $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ where $A \in \mathbb{Z}^{n \times m}$ and every absolute square subdeterminant of A is at most Δ then

$$\text{Diameter}(P) \leq O(n^3 \Delta^2 \log(\Delta)).$$


Random polytopes are “well-conditioned on average”. Do they have small diameter?

Tantalizing evidence from the simplex method

Let

$$P = \{x \in \mathbb{R}^n : \langle a, x \rangle \leq 1 \ \forall a \in A\}$$

where $A \subset \mathbb{S}^{n-1}$, $|A| = m$ is sampled iid.




Tantalizing evidence from the simplex method

Let

$$P = \{x \in \mathbb{R}^n : \langle a, x \rangle \leq 1 \ \forall a \in A\}$$

where $A \subset \mathbb{S}^{n-1}$, $|A| = m$ is sampled iid.




Borgwardt 1977,1982,1987,1999:
consider a fixed two-dimensional
subspace $W \subset \mathbb{R}^n$. On expectation,
 $O(n^2 m^{\frac{1}{n-1}})$ distinct vertices are
optimized by the objectives in W .

Tantalizing evidence from the simplex method

Let

$$P = \{x \in \mathbb{R}^n : \langle a, x \rangle \leq 1 \ \forall a \in A\}$$

where $A \subset \mathbb{S}^{n-1}$, $|A| = m$ is sampled iid.




Borgwardt 1977,1982,1987,1999:
consider a fixed two-dimensional
subspace $W \subset \mathbb{R}^n$. On expectation,
 $O(n^2 m^{\frac{1}{n-1}})$ distinct vertices are
optimized by the objectives in W .

Tantalizing evidence from the simplex method

Let

$$P = \{x \in \mathbb{R}^n : \langle a, x \rangle \leq 1 \ \forall a \in A\}$$

where $A \subset \mathbb{S}^{n-1}$, $|A| = m$ is sampled iid.



Borgwardt 1977,1982,1987,1999:
consider a fixed two-dimensional
subspace $W \subset \mathbb{R}^n$. On expectation,
 $O(n^2 m^{\frac{1}{n-1}})$ distinct vertices are
optimized by the objectives in W .

\implies two vertices optimized by
randomly chosen objective vectors
are at distance $O(n^2 m^{\frac{1}{n-1}})$ in
expectation.

Our results

Let

$$P = \{x \in \mathbb{R}^n : \langle a, x \rangle \leq 1 \ \forall a \in A\}$$

where $A \subset \mathbb{S}^{n-1}$ follows a Poisson point process with $\mathbb{E}[|A|] = m > 2^{\Omega(n)}$.

Our results

Let

$$P = \{x \in \mathbb{R}^n : \langle a, x \rangle \leq 1 \ \forall a \in A\}$$

where $A \subset \mathbb{S}^{n-1}$ follows a Poisson point process with $\mathbb{E}[|A|] = m > 2^{\Omega(n)}$.

Then with probability $1 - O(1/m)$ we have

$$\Omega(nm^{\frac{1}{n-1}}) \leq \text{Diameter}(P) \leq O(n^2 m^{\frac{1}{n-1}} + 17^n)$$

$$\Omega(m^{\frac{1}{n-1}}) \leq \text{Diameter}(\text{Conv}(A)) \leq O(nm^{\frac{1}{n-1}} + 17^n).$$

A Weak but Simple Lower bound

A set $B \subset \mathbb{S}^{n-1}$ is called ε -dense if for every $x \in \mathbb{S}^{n-1}$ there exists $b \in B$ with $\|x - b\| \leq \varepsilon$.

A Weak but Simple Lower bound

A set $B \subset \mathbb{S}^{n-1}$ is called ε -dense if for every $x \in \mathbb{S}^{n-1}$ there exists $b \in B$ with $\|x - b\| \leq \varepsilon$.

Constraint set A is $O((\log(m)/m)^{\frac{1}{n-1}})$ -dense with probability $1 - 1/m$.

A Weak but Simple Lower bound

A set $B \subset \mathbb{S}^{n-1}$ is called ε -dense if for every $x \in \mathbb{S}^{n-1}$ there exists $b \in B$ with $\|x - b\| \leq \varepsilon$.

Constraint set A is $O((\log(m)/m)^{\frac{1}{n-1}})$ -dense with probability $1 - 1/m$.

If A is ε -dense then the boundary of P lies inside $(1 - \varepsilon^2/2)^{-1} \mathbb{B}^n \setminus \mathbb{B}^n$.

A Weak but Simple Lower bound

A set $B \subset \mathbb{S}^{n-1}$ is called ε -dense if for every $x \in \mathbb{S}^{n-1}$ there exists $b \in B$ with $\|x - b\| \leq \varepsilon$.

Constraint set A is $O((\log(m)/m)^{\frac{1}{n-1}})$ -dense with probability $1 - 1/m$.

If A is ε -dense then the boundary of P lies inside $(1 - \varepsilon^2/2)^{-1}\mathbb{B}^n \setminus \mathbb{B}^n$.

If a line segment lies inside $(1 - \varepsilon^2/2)^{-1}\mathbb{B}^n \setminus \mathbb{B}^n$ and $\varepsilon \leq 1$ then it has length $\leq 2\varepsilon$


A Weak but Simple Lower bound

A set $B \subset \mathbb{S}^{n-1}$ is called ε -dense if for every $x \in \mathbb{S}^{n-1}$ there exists $b \in B$ with $\|x - b\| \leq \varepsilon$.

Constraint set A is $O((\log(m)/m)^{\frac{1}{n-1}})$ -dense with probability $1 - 1/m$.

If A is ε -dense then the boundary of P lies inside $(1 - \varepsilon^2/2)^{-1}\mathbb{B}^n \setminus \mathbb{B}^n$.

If a line segment lies inside $(1 - \varepsilon^2/2)^{-1}\mathbb{B}^n \setminus \mathbb{B}^n$ and $\varepsilon \leq 1$ then it has length $\leq 2\varepsilon$



A Weak but Simple Lower bound

A set $B \subset \mathbb{S}^{n-1}$ is called ε -dense if for every $x \in \mathbb{S}^{n-1}$ there exists $b \in B$ with $\|x - b\| \leq \varepsilon$.

Constraint set A is $O((\log(m)/m)^{\frac{1}{n-1}})$ -dense with probability $1 - 1/m$.

If A is ε -dense then the boundary of P lies inside $(1 - \varepsilon^2/2)^{-1}\mathbb{B}^n \setminus \mathbb{B}^n$.

If a line segment lies inside $(1 - \varepsilon^2/2)^{-1}\mathbb{B}^n \setminus \mathbb{B}^n$ and $\varepsilon \leq 1$ then it has length $\leq 2\varepsilon$

\implies If A is ε -dense then $\text{Diameter}(P) \geq \Omega(1/\varepsilon)$.

A Weak but Simple Lower bound

A set $B \subset \mathbb{S}^{n-1}$ is called ε -dense if for every $x \in \mathbb{S}^{n-1}$ there exists $b \in B$ with $\|x - b\| \leq \varepsilon$.

Constraint set A is $O((\log(m)/m)^{\frac{1}{n-1}})$ -dense with probability $1 - 1/m$.

If A is ε -dense then the boundary of P lies inside $(1 - \varepsilon^2/2)^{-1}\mathbb{B}^n \setminus \mathbb{B}^n$.

If a line segment lies inside $(1 - \varepsilon^2/2)^{-1}\mathbb{B}^n \setminus \mathbb{B}^n$ and $\varepsilon \leq 1$ then it has length $\leq 2\varepsilon$

\implies If A is ε -dense then $\text{Diameter}(P) \geq \Omega(1/\varepsilon)$.

\implies With high probability $\text{Diameter}(P) \geq \Omega(m/\log m)^{\frac{1}{n-1}}$.

Implications

- ▶ If all vertices of P are induced by “nearby” constraint vectors, then “far-apart” pieces of P are nearly independent.

Implications

- ▶ If all vertices of P are induced by “nearby” constraint vectors, then “far-apart” pieces of P are nearly independent.
- ▶ This happens with high probability \implies shadow size concentrates around its mean.

Implications

- ▶ If all vertices of P are induced by “nearby” constraint vectors, then “far-apart” pieces of P are nearly independent.
- ▶ This happens with high probability \implies shadow size concentrates around its mean.
- ▶ Many 2D planes simultaneously have small shadow sizes.

Implications

- ▶ If all vertices of P are induced by “nearby” constraint vectors, then “far-apart” pieces of P are nearly independent.
- ▶ This happens with high probability \implies shadow size concentrates around its mean.
- ▶ Many 2D planes simultaneously have small shadow sizes.
- ▶ Can be leveraged to upper bound the diameter (!).

Future directions

- ▶ Close the gap between upper and lower bound.
- ▶ Other distributions, such as Gaussian?
- ▶ What if $m \leq \text{poly}(n)$?