Journal of Meaning-Centered Psychology
PUBLISHED BY THE INSTITUTE FOR MEANING-CENTERED PSYCHOLOGY

Volume 1 · Issue 1 · April 2026
pp. 11–13 · Open access · CC BY 4.0

THEORETICAL ARTICLE

Paradoxical Intention and Modern Exposure Procedures

A Theoretical Comparison of Assumed Mechanisms, Procedure, and Limitations

Samuel R. Ashford, Eleanor M. Whitcombe and Sora Pazer
Institute for Meaning-Centered Psychology

ABSTRACT

Paradoxical intention is among the best-known techniques of logotherapy. Viktor E. Frankl described it as early as the period between the 1930s and the 1950s as a method for interrupting anticipatory anxiety in phobias and obsessive-compulsive symptoms. This paper compares paradoxical intention at a theoretical level with contemporary exposure-based procedures in cognitive behavioral therapy, in particular with emotional processing theory and the inhibitory learning approach. First, substantial similarities are identified: both approaches aim at relinquishing avoidance and control behaviors, encourage confrontation with feared experiences, and enable corrective learning. At the same time, there are differences in their anthropological rationale, in the role of humor and self-distancing, and in how the procedure is structured. The paper then discusses to what extent concepts such as ironic processes of mental control, metacognitive models, and cognitive defusion may serve as bridges between the two traditions. It concludes with a discussion of the limits of indication and with suggestions for a more precise theoretical placement of paradoxical intention within current psychotherapy research, without abandoning its distinct existential-analytic foundation.

Keywords: paradoxical intention · exposure therapy · inhibitory learning · self-distancing · anxiety disorders

1 Introduction

Among the techniques of logotherapy, paradoxical intention has achieved the widest dissemination beyond its own school. In a much-cited paper, Frankl (1960) described it as a procedure in which patients are instructed to wish for, or deliberately bring about, precisely what they fear instead of strenuously avoiding it. In his account of the theory of neuroses, he systematically embedded this approach in a model of anticipatory anxiety (Frankl, 1956).

Since then, the treatment of anxiety and obsessive-compulsive disorders has developed considerably. Exposure-based procedures are today among the best-researched psychotherapeutic interventions of all (Abramowitz et al., 2019). It therefore seems natural to compare paradoxical intention with these procedures. The present contribution does so at the theoretical level. It asks which assumptions about mechanisms of action underlie the two approaches, where they converge, and where they differ.

The comparison is also worthwhile because paradoxical intention is occasionally referred to in the literature as a forerunner of behavioral techniques. Such a classification is historically understandable but does only partial justice to Frankl's self-understanding. It is therefore necessary to take seriously both the functional kinship and the independent rationale.

2 Paradoxical Intention as Understood in Logotherapy

Frankl's starting point is a circular model: a symptom triggers the fear that it might recur; this anticipatory anxiety in turn promotes the renewed occurrence of the symptom, which further reinforces the fear (Frankl, 1956). According to Frankl, typical patterns of reaction to this vicious circle are flight from fear in phobic complaints and fighting against the obsessive idea in obsessive-compulsive symptoms. Both patterns contribute to the entrenchment of the problem.

Paradoxical intention intervenes at this point. It asks the affected person to wish consciously for what they fear – for instance blushing, trembling, or sweating – and, where possible, to do so in an exaggerated, humorous form. Frankl (1960) emphasized that this technique is based on the specifically human capacity for self-distancing, which he attributed to the noetic dimension. Humor is not a decorative addition here but a central means of gaining distance from the symptom and depriving the fear of its power.

It is noteworthy that Frankl did not understand paradoxical intention as a mere technique. It was intended to enable the person not to identify with their fear but to adopt a stance toward it. This expresses the basic anthropological assumption of logotherapy that human beings are not entirely at the mercy of their psychological conditions but can take a stand toward them.

Frankl also distinguished paradoxical intention from dereflection. While paradoxical intention targets anticipatory anxiety, dereflection addresses excessive self-observation, which Frankl called hyperreflection and described, for example, in sleep disorders and sexual dysfunctions (Frankl, 1956). Common to both procedures is the assumption that a certain form of strained attention to oneself maintains the very problem it seeks to solve.

3 Exposure Procedures: From Emotional Processing to Inhibitory Learning

The theoretical rationale for exposure has changed over the past decades. With emotional processing theory, Foa and Kozak (1986) formulated a model according to which fear is represented in memory structures that can be modified through activation and the incorporation of corrective information. In this model, habituation within and between practice sessions was regarded as an important indicator of successful processing.

Craske et al. (2014), by contrast, proposed an inhibitory learning model. According to this model, the original fear association is not erased by exposure but supplemented by newly learned, inhibitory associations that compete with the original one. For practice, it follows that the focus is less on reducing fear during the exercise than on violating expectancies, varying practice contexts, and refraining from safety behaviors. Tolerating fear thus becomes a goal in itself.

Also significant for practice is that modern exposure procedures are usually based on careful case conceptualization and a convincing rationale that makes it comprehensible to patients why confronting fear is worthwhile (Abramowitz et al., 2019). Here, too, a parallel emerges: Frankl attached importance to ensuring that those affected understand the mechanism of anticipatory anxiety before they apply paradoxical intention.

4 Convergences

From the perspective of inhibitory learning, paradoxical intention appears in a new light. By wishing for what they fear, patients give up avoidance and control strategies and expose themselves to the feared experience. The expectation that the symptom will escalate catastrophically is frequently not confirmed in the process. Viewed functionally, paradoxical intention thus contains central elements of an exposure exercise: confrontation, abandonment of safety behaviors, and expectancy violation.

Research on thought control offers a further bridge. Wegner (1994) showed that attempting to suppress certain thoughts can, under certain conditions, lead precisely to their increased occurrence. This concept of ironic processes provides an experimentally grounded explanation for the observation, described clinically by Frankl, that fighting a symptom can intensify it. The metacognitive model of Wells (2009), which focuses on problematic beliefs about the controllability and dangerousness of one's own thoughts, also shows points of contact here.

Finally, there are parallels with cognitive defusion in Acceptance and Commitment Therapy, in which thoughts are regarded as passing mental events rather than being treated as literal truths (Hayes et al., 2012). Some defusion exercises that work with exaggeration, repetition, or playful alienation resemble, in their structure, the humorous distancing that Frankl described.

5 Differences

Despite these convergences, it would be reductive to interpret paradoxical intention simply as an early form of exposure. A first difference concerns the rationale: while exposure procedures are grounded in the psychology of learning and memory, Frankl invokes an anthropological capacity for self-distancing. A second difference lies in the degree of structuring. Modern exposure protocols are usually carefully planned, with hierarchies, expectancy checks, and debriefings. Paradoxical intention, by contrast, was often conveyed as a compact attitude to be applied in everyday life.

A third difference concerns the role of humor. In exposure procedures, humor is at most a facilitative contextual factor, whereas in paradoxical intention it is constitutive. Whether and to what extent humor itself contributes to efficacy is an open empirical question. In addition, paradoxical interventions have been studied in the literature under very different names and procedural formats. An early meta-analysis of paradoxical interventions (Shoham-Salomon & Rosenthal, 1987) does point to a certain efficacy but at the same time indicates considerable heterogeneity, so that conclusions about the logotherapeutic technique in the narrower sense can be drawn only with caution.

A fourth difference concerns the goal. Exposure procedures are directed primarily at reducing fear and avoidance or at improving fear tolerance. For Frankl, by contrast, overcoming anticipatory anxiety was not an end in itself but a precondition for the person being able to turn once again to the tasks and values of their life. In this respect, paradoxical intention is close to Acceptance and Commitment Therapy, which likewise does not regard fear reduction as the primary goal but places value-oriented action at the center (Hayes et al., 2012).

6 Limits of Indication and Outlook

Frankl (1956) himself pointed out that paradoxical intention is not indicated for all clinical presentations. In depressive states with suicidality, for example, an invitation to wish for what one fears would obviously be contraindicated. In the case of obsessive thoughts with aggressive content, too, careful psychoeducation is required so that the technique is not experienced as trivialization. A sound working alliance and a clear rationale for the intervention are therefore indispensable.

For future research, the inhibitory learning model offers a framework in which hypotheses about how paradoxical intention works can be formulated precisely, for instance regarding the role of expectancy violation or the contribution of humor and self-distancing. Such an integration would not deprive paradoxical intention of its existential-analytic rationale but would supplement it with a compatible mechanism perspective.

Methodologically, component analyses lend themselves to testing such hypotheses, in which, for example, an exposure condition with and without a humorous-paradoxical instruction is compared. In this way, it could be clarified whether the paradoxical formulation makes an independent contribution beyond the confrontation. It would also be worth investigating whether people with a pronounced tendency toward thought suppression or with metacognitive beliefs about the dangerousness of their own thoughts benefit particularly from a paradoxical stance (Wegner, 1994; Wells, 2009).

References

Abramowitz, J. S., Deacon, B. J., & Whiteside, S. P. H. (2019). Exposure therapy for anxiety: Principles and practice (2nd ed.). Guilford Press.

Craske, M. G., Treanor, M., Conway, C. C., Zbozinek, T., & Vervliet, B. (2014). Maximizing exposure therapy: An inhibitory learning approach. Behaviour Research and Therapy, 58, 10–23.

Foa, E. B., & Kozak, M. J. (1986). Emotional processing of fear: Exposure to corrective information. Psychological Bulletin, 99(1), 20–35.

Frankl, V. E. (1956). Theorie und Therapie der Neurosen: Einführung in Logotherapie und Existenzanalyse. Urban & Schwarzenberg.

Frankl, V. E. (1960). Paradoxical intention: A logotherapeutic technique. American Journal of Psychotherapy, 14(3), 520–535.

Hayes, S. C., Strosahl, K. D., & Wilson, K. G. (2012). Acceptance and commitment therapy: The process and practice of mindful change (2nd ed.). Guilford Press.

Shoham-Salomon, V., & Rosenthal, R. (1987). Paradoxical interventions: A meta-analysis. Journal of Consulting and Clinical Psychology, 55(1), 22–28.

Wegner, D. M. (1994). Ironic processes of mental control. Psychological Review, 101(1), 34–52.

Wells, A. (2009). Metacognitive therapy for anxiety and depression. Guilford Press.

HOW TO CITE

Ashford, S. R., Whitcombe, E. M., & Pazer, S. (2026). Paradoxical Intention and Modern Exposure Procedures: A Theoretical Comparison of Assumed Mechanisms, Procedure, and Limitations. Journal of Meaning-Centered Psychology, 1(1), 11–13.

LICENSE, CORRESPONDENCE AND USE OF AI

Published open access under the Creative Commons Attribution 4.0 International license (CC BY 4.0). Correspondence: Editorial office, Journal of Meaning-Centered Psychology. This article was written with the assistance of generative AI and reviewed by the authors, who take full responsibility for its content.