

ALGEBRAIC GEOMETRY EXERCISES

1. AFFINE VARIETIES

- (1) Draw the following algebraic sets in $\mathbb{A}_{\mathbb{R}}^2$ (with coordinates x, y):
 - (a) $Z(xy - 1)$
 - (b) $Z(x^3 - xy^2)$
 - (c) $Z(x^2 + y^2 - 1)$
- (2) Show that the set of points $\{(t, \sin t) \mid t \in \mathbb{R}\} \subset \mathbb{A}_{\mathbb{R}}^2$ is not an algebraic set.
- (3) Let $X = Z(I) \subset \mathbb{A}^n$ for some ideal $I \subset k[x_1, \dots, x_n]$. Let $p \in \mathbb{A}^n \setminus X$. Show that there is a polynomial $f \in I$ such that $f(p) = 1$.
- (4) Show that $f = x^2 + y^2 + z^2$ is irreducible as an element of $\mathbb{R}[x, y, z]$. What is the dimension of $Z(f) \subset \mathbb{A}_{\mathbb{R}}^3$?
- (5) Let $X \subset \mathbb{A}^3$ be the algebraic set defined by $X = Z(x^3 + y^2 + z^2, xy - xz)$. Describe the irreducible components of X .
- (6) Show that the algebraic set $Z(xy, xz) \subset \mathbb{A}^3$ has two components of different dimension.
- (7) (a) Work in the affine plane $\mathbb{A}_{\mathbb{R}}^2$, with coordinates x, y . Let $L = Z(y)$ be the x -axis, choose a number t and let $Q_t = Z(x^2 + t - y)$ be the parabola. Determine the subsets U_0, U_1, U_2 of $\mathbb{R}$ such that:

$$L \cap Q_t = \begin{cases} \emptyset & \text{if and only if } t \in U_0 \\ \text{a point} & \text{if and only if } t \in U_1 \\ \text{two points} & \text{if and only if } t \in U_2 \end{cases}$$

- (b) Which of U_0, U_1, U_2 are quasi-affine varieties?
 - (c) Do (a) and (b) again, but with $\mathbb{C}$ in place of $\mathbb{R}$.
- (8) (a) Suppose that L is a line in $\mathbb{A}^2$, and let $C \subset \mathbb{A}^2$ be a curve of degree $d > 1$. Show that $L \cap C$ consists of at most d points.
 - (b) Construct an example to show that it is possible for $L \cap C$ to be empty, even over an algebraically closed field.
- (9) (a) Let $p = (a_1, \dots, a_n) \in \mathbb{A}^n$ be a point. Show that $I(\{p\}) = (x_1 - a_1, \dots, x_2 - a_2, \dots, x_n - a_n)$.
 - (b) Show that $I(\{p\})$ is a maximal ideal of $k[x_1, \dots, x_n]$.

- (c) Show that, if k is algebraically closed, every maximal ideal of $k[x_1, \dots, x_n]$ has the form $I(\{p\})$.
- (d) Give an example of a maximal ideal of $\mathbb{R}[x]$ that is *not* of the form $I(\{p\})$.
- (10) For every polynomial $f \in k[x_1, \dots, x_n]$, let $U(f) = \mathbb{A}^n \setminus Z(f)$. Show that the sets $U(f)$ form a *subbasis* for the Zariski topology on $\mathbb{A}^n$. (This means: Any open set is a union of finite intersections of sets on the form $U(f)$.)
- (11) *The twisted cubic curve.* Consider the set of points $C = \{(t, t^2, t^3) | t \in k\} \subset \mathbb{A}^3$. Find generators of the ideal $I(C)$, and show that C is an affine variety. Show that there is an isomorphism $k[x] \cong \mathcal{O}(C)$.
- (12) Think of $\mathbb{A}^2$ as $\mathbb{A}^1 \times \mathbb{A}^1$ in the straightforward way. Show that the Zariski topology on $\mathbb{A}^2$ does not equal the product of the Zariski topologies on the two copies of $\mathbb{A}^1$.
- (13) Show that a nonempty open subset U of an irreducible topological space X is irreducible and *dense*. (U is dense if every open subset $V \subset X$ satisfies $U \cap V \neq \emptyset$.)
- (14) Let B be a finitely-generated k -algebra. Show that there is some algebraic set V for which $B = \mathcal{O}(V)$ if and only if B has no nilpotent elements. (A *nilpotent* element of a ring B is an element $a \in B$ such that $a \neq 0$, and there is some $k > 0$ such that $a^k = 0$.)

2. PROJECTIVE VARIETIES

- (1) Consider the projective algebraic set $C = Z(x^2z^2 - y^4 - xy^2z) \subset \mathbb{P}^2$. Before drawing them, in which two of the affine charts $C \cap D(x), C \cap D(y), C \cap D(z)$ will you get the same image? Draw each of the charts to verify. (Use a computer if you like!)
- (2) (a) Show that $X = Z(xy - z^2) \subset \mathbb{P}^2$ is the projective closure of the hyperbola $X' = Z(xy - 1) \subset \mathbb{A}^2$.
 (b) Let $L'_1, L'_2 \subset \mathbb{A}^2$ be the asymptotic lines to X' . Describe the projective closures L_1, L_2 of the two lines. Show that $X \cap L_1$ and $X \cap L_2$ are nonempty.
- (3) Count the number of points in $\mathbb{P}_{\mathbb{F}_q}^n$, where $\mathbb{F}_q$ is the finite field with q elements. Conclude that the only n, m for which $\mathbb{P}_{\mathbb{F}_q}^n = \mathbb{A}_{\mathbb{F}_q}^m$ are $n = m = 0$.
- (4) Show that every quasiprojective variety can be covered by quasiaffine varieties.
- (5) If $f \in k[x_0, \dots, x_n]$ is homogeneous of degree 1, we call $H = Z(f) \subset \mathbb{P}^n$ a *hyperplane*. Show that the intersection of n hyperplanes in $\mathbb{P}^n$ is nonempty. Show that if the intersection is a finite set, then it is precisely one point.
- (6) Show that if $H \subset \mathbb{P}^n$ is a hyperplane, there is a homeomorphism $H \cong \mathbb{P}^{n-1}$.
- (7) (a) Show that if k is algebraically closed, the intersection of a curve and a line in $\mathbb{P}^2$ is nonempty.


- (b) If $f \in k[x_0, \dots, x_n]$ is irreducible and homogeneous of degree d , we call $X = Z(f) \subset \mathbb{P}^n$ a *projective hypersurface* of degree d . Show that if k is algebraically closed, the intersection of X with $n - 1$ hyperplanes is nonempty. (Hint: Use induction, with (a) as the base case.)
- (8) Consider the function $\phi: \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^3$ given by
- $$\phi: ((a_1 : a_2), (b_1 : b_2)) \mapsto (a_1 b_1 : a_1 b_2 : a_2 b_1 : a_2 b_2).$$
- (a) Show that ϕ is injective, and that the image of ϕ is the projective variety $X = Z(x_0 x_3 - x_1 x_2)$.
- (b) Show that for any $p \in \mathbb{P}^1$, both $\phi(p \times \mathbb{P}^1)$ and $\phi(\mathbb{P}^1 \times p)$ are lines. Show that every line in X can be written as either $\phi(p \times \mathbb{P}^1)$ or $\phi(\mathbb{P}^1 \times p)$.
- (c) Show that, if l_1, l_2 are two lines in X , then it is possible that $l_1 \cap l_2 = \emptyset$. When does this happen?
- (9) Let $X = Z(x_0^3 + x_1^3 + x_2^3 + x_3^3) \subset \mathbb{P}_{\mathbb{C}}^3$. Show that there are at least 27 lines in X . (In fact, there are precisely 27, but we won't show this.)
- (10) Choose an integer $n > 0$ and consider the function $\phi: \mathbb{P}^1 \rightarrow \mathbb{P}^n$ given by

$$(a_0 : a_1) \mapsto (a_0^n : a_0^{n-1} a_1 : \dots : a_0 a_1^{n-1} : a_1^n).$$

- (a) Show that ϕ is injective.
- (b) Show that the image of ϕ is a projective variety, and find generators for $I(\text{im}(\phi))$.
- (c) Show that, for $n = 3$, $\phi(\mathbb{P}^1)$ is the projective closure of the *twisted cubic curve* from exercise (10) in the *Affine varieties* section.

3. CURVES AND SINGULARITIES

- (1) Draw the following curves in the affine plane $\mathbb{A}_{\mathbb{R}}^2$. If they are singular, find their singular points.
- (a) $Z(x^3 - y^2 - 1)$
- (b) $Z(x^2 - y^4)$
- (c) $Z(x^3 - y^3 - yx)$. (This curve is called the *folium of Descartes*.) Find the equation of the unique line that does not intersect the curve.
- (d) $Z((x^2 + 1)y - 1)$. (This curve is called the *witch of Agnesi*.)



- (2) Show that the degree of this curve:
 is at least 6. (Hint: Find a line that intersects the curve 6 times.) (The equation is $(x^2 + y^2)^3 - (x^2 - y^2)^2$.)
- (3) Draw the curve $C = Z((x^2 + y^2 - 1)^3 - x^2 y^3) \subset \mathbb{A}_{\mathbb{R}}^2$. (Isn't it nice?) Where are the singular points of C ?
- (4) Let X be a (projective or affine) variety. Let $\text{Sing}(X)$ be the set of singular points on X . Show that $\text{Sing}(X)$ is a closed proper subset of X . We call $\text{Sing}(X)$ the *singular locus* of X .
- (5) Find the singular locus of the projective variety

$$Z(x_0 x_1 x_2 + x_0 x_2 x_3 + x_0 x_1 x_3 + x_1 x_2 x_3) \subset \mathbb{P}^3.$$

(This variety is called the *Cayley nodal cubic surface*.)

- (6) Choose an integer n .
- Show that the surface $S = Z(x_1 x_2 - x_3^n) \subset \mathbb{A}^3$ is singular for every $n > 1$.
 - Identify $\mathbb{A}^3$ with $D^+(x_0) \subset \mathbb{P}^3$, and find the equations for the projective closure $\bar{S}$ of S in $\mathbb{P}^3$. Where is $\bar{S}$ singular?
- (7) Let $C = Z(x^2 + yz) \subset \mathbb{P}_k^2$, where k is a field of characteristic 2. Show that, for any point $p \in C$, the tangent line $T_p C$ passes through the point $(1 : 0 : 0)$. (We say that C is a *strange curve*.)

4. MORPHISMS

- (1) *Projection from a point.* Let $p = (1, 0, 0, \dots, 0) \in \mathbb{A}^n$, and let $H = Z(x_1)$ be the hyperplane. Define a function $\pi: \mathbb{A}^n \setminus \{p\} \rightarrow H$ by sending any point $q = (a_1, a_2, \dots, a_n)$ to $(0, a_2, \dots, a_n) \in H$. Show that π is a morphism of varieties.
- (2) Let p be a prime number, and let k be a field of characteristic p .
- Show that there is a ring homomorphism $\phi: k[x_1, \dots, x_n] \rightarrow k[x_1, \dots, x_n]$ determined by $f \mapsto f^p$ for every $f \in k[x_1, \dots, x_n]$.

- (b) Let $X \subset \mathbb{A}_k^n$ be an affine variety. Show that ϕ induces a morphism $\phi': X \rightarrow X$.
(This is called the *Frobenius endomorphism*.)
- (3) (a) Let $f \in k[x_1, \dots, x_n]$ be a nonzero polynomial. Let $U = \mathbb{A}^n \setminus Z(f)$. Show that U is isomorphic to the affine variety

$$Y = Z(x_{n+1}f - 1) \subset \mathbb{A}^{n+1}.$$

(Hint: You can define a map $\phi: U \rightarrow Y$ by

$$(a_1, \dots, a_n) \mapsto (a_1, \dots, a_n, \frac{1}{f(a_1, \dots, a_n)}).$$

Show that ϕ is the required isomorphism.)

- (b) Use (a) to show that, if $X \subset \mathbb{A}^n$ is an affine variety, then $X \setminus Z(f)$ is isomorphic to an affine variety.
- (4) Let C be a projective curve, let $U \subset C$ be open, and let $f: U \rightarrow \mathbb{A}^1$ be a regular map. Show that there is another open subset $V \subset C$ and a regular map $g: V \rightarrow \mathbb{A}^1$ with the properties that:
- (a) $V \cup U = C$
- (b) On $V \cap U$, $f = \frac{1}{g}$.
- Use f and g to define a morphism $C \rightarrow \mathbb{P}^1$. (Hint: Write $\mathbb{P}^1$ as a union of $D^+(x_0)$ and $D^+(x_1)$, and consider f and g as morphisms to $D^+(x_0), D^+(x_1)$, respectively.)
- Conclude that for every curve C , it is always possible to find a morphism $C \rightarrow \mathbb{P}^1$.
- (5) Let $f \in k[x_0, \dots, x_n]$ be irreducible and homogeneous of degree $d > 1$. Let $X = Z(f) \subset \mathbb{A}^{n+1}$, and let $Y = Z(f) \subset \mathbb{P}^n$. Show that $\mathbf{0} \in X$, and that there is a morphism $X \rightarrow Y$ given by $X \ni (a_0, \dots, a_n) \mapsto (a_0 : \dots : a_n) \in Y$. We call X the *cone* over Y .
- (6) *The function field of a variety.* Let X be a variety, and consider the set $\text{Reg}(X)$ of all pairs (V, f) where V is an open subset of X and $f: V \rightarrow k$ is a regular function. We impose an equivalence relation $\sim$ on $\text{Reg}(X)$ by saying:

$$(V, f) \sim (U, g) \Leftrightarrow f|_{U \cap V} = g|_{U \cap V}.$$

Let $K(X)$ be the set of $\sim$ -equivalence classes of $\text{Reg}(X)$. For any $(V, f) \in \text{Reg}(X)$, we write its $\sim$ -equivalence class as $\overline{(V, f)}$. In this exercise, we will show that $K(X)$ is a field.

- (a) Let $(V, f), (U, g)$ be two elements of $\text{Reg}(X)$. Show that $V \cap U$ is nonempty.
- Then:
- (b) Show that $\overline{(V \cap U, f + g)} \in K(X)$.
- (c) Show that $\overline{(V \cap U, fg)} \in K(X)$.

- (d) For any $(\overline{V}, f) \in K(X)$, show that $(\overline{V}, -f) \in K(X)$.
- (e) Let $(\overline{V}, f) \in K(X)$, and suppose that we can write $f = g/h$ where g, h are polynomials. Show that $(\overline{V \setminus Z(g)}, \frac{1}{f}) \in K(X)$.
- (f) Use all the previous points to define a field structure on $K(X)$.
- (g) Show that, if $U \subset X$ is open, then there is an isomorphism of fields $K(U) \rightarrow K(X)$.
- (h) Show that $K(\mathbb{A}^1) = k(x)$, the field of all rational functions in one variable.
Conclude that $K(\mathbb{P}^1) = k(x)$.

The field $K(X)$ is called the *function field* of X .