

# Worksheet SOLUTIONS: Propositional Logic & First-Order Logic

Foundations of Artificial Intelligence – SS 2026

University of Bremen, Institute for Artificial Intelligence

## Part 1 – Entailment & Satisfiability

**1.1 True.**  $A \models A \vee B$  because every model that makes  $A$  true also makes  $A \vee B$  true (disjunction is weaker). There is no model satisfying  $A$  that falsifies  $A \vee B$ .

**1.2 Yes, tautology.** Truth table:

| A | B | $A \rightarrow B$ | $B \rightarrow A$ | $(A \rightarrow B) \vee (B \rightarrow A)$ |
|---|---|-------------------|-------------------|--------------------------------------------|
| T | T | T                 | T                 | T                                          |
| T | F | F                 | T                 | T                                          |
| F | T | T                 | F                 | T                                          |
| F | F | T                 | T                 | T                                          |

At least one of  $A \rightarrow B$  or  $B \rightarrow A$  is always true, so the disjunction is always true.

**1.3**

| #   | Formula                                      | Satisfiable?         | Unsatisfiable? | Tautology? |
|-----|----------------------------------------------|----------------------|----------------|------------|
| i   | $A \wedge \neg A$                            | ✗                    | ✓              | ✗          |
| ii  | $A \vee \neg A$                              | ✓                    | ✗              | ✓          |
| iii | $(A \rightarrow B) \wedge (B \rightarrow A)$ | ✓ (e.g. $A=T, B=T$ ) | ✗              | ✗          |
| iv  | $\neg(A \vee B) \wedge A$                    | ✗                    | ✓              | ✗          |

**1.4 Yes.**  $A \rightarrow B$  means  $(A \rightarrow B) \wedge (B \rightarrow A)$ . The left conjunct is exactly  $A \rightarrow B$ , so  $A \rightarrow B \models A \rightarrow B$ .

**1.5 (a) only.** From  $A \wedge B$  we can derive  $A$  and  $B$  individually, and by disjunction introduction  $A \vee B \vee C$ . We cannot derive  $A \rightarrow B$  in general without knowing the relationship between  $A$  and  $B$  —  $A \wedge B$  is consistent with any truth value combination, but  $A \rightarrow B$  requires that they always match, which is not entailed by  $A \wedge B$  being true in one model.

## Part 2 – Counting Models

**Vocabulary:**  $\{A, B, C, D\}$ , 16 total models.

(a)  $(A \wedge B) \vee (B \wedge C)$

Factor out B:  $B \wedge (A \vee C)$ . B must be true (8 models), and at least one of A, C must be true (3 of 4 combinations of A,C are valid). D is free ( $\times 2$ ).

**Answer: 6 models** — wait, let's count carefully.

$B=T, A=T, C=T \rightarrow$  valid;  $B=T, A=T, C=F \rightarrow$  valid;  $B=T, A=F, C=T \rightarrow$  valid;  $B=T, A=F, C=F \rightarrow$  invalid.

So 3 combinations  $\times$  2 values of D = **6 models**.

(b)  $A \vee B$

Complement: neither A nor B =  $A=F, B=F \rightarrow$  4 models (C and D free). Valid models =  $16 - 4 =$  **12 models**.

(c)  $(A \rightarrow B) \rightarrow D$

$A \rightarrow B$  is true in 2 out of 4 (A,B) combinations (both T or both F). The outer biconditional is true when both sides match.

- $(A \rightarrow B) = T, D=T \rightarrow 2 \times 1 = 2$  (C free  $\times 2 = 4$ )
- $(A \rightarrow B) = F, D=F \rightarrow 2 \times 1 = 2$  (C free  $\times 2 = 4$ )

**Answer: 8 models.**

## Part 3 – Validity and Unsatisfiability

|   | Formula                                                                                                                                                     | Answer  | Reason                                                                                                                                                                                                                                                                                                                                                                                            |
|---|-------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| a | $\text{Smoke} \rightarrow \text{Smoke}$                                                                                                                     | Valid   | $P \rightarrow P$ is always true                                                                                                                                                                                                                                                                                                                                                                  |
| b | $(\text{Smoke} \rightarrow \text{Fire}) \rightarrow (\neg \text{Smoke} \rightarrow \neg \text{Fire})$                                                       | Neither | Contrapositive confusion — the converse is not equivalent. Counterexample: Smoke=F, Fire=T makes the premise true ( $F \rightarrow T$ ), but conclusion $F \rightarrow F = T$ . Actually try Smoke=F, Fire=T: premise T, $\neg \text{Smoke}=T$ , $\neg \text{Fire}=F$ , so conclusion $T \rightarrow F = F$ . Formula is false. Not valid; satisfiable when Smoke=T. $\rightarrow$ <b>Neither</b> |
| c | $\text{Smoke} \vee \text{Fire} \vee \neg \text{Fire}$                                                                                                       | Valid   | Contains $\text{Fire} \vee \neg \text{Fire}$ which is always true                                                                                                                                                                                                                                                                                                                                 |
| d | $((\text{Smoke} \wedge \text{Heat}) \rightarrow \text{Fire}) \rightarrow (\text{Smoke} \rightarrow \text{Fire}) \vee (\text{Heat} \rightarrow \text{Fire})$ | Valid   | Both sides are logically equivalent — if either single condition implies Fire, then surely both together do                                                                                                                                                                                                                                                                                       |
| e | $(\text{Smoke} \rightarrow \text{Fire}) \rightarrow ((\text{Smoke} \wedge \text{Heat}) \rightarrow \text{Fire})$                                            | Valid   | Adding more antecedents to an implication preserves it. If Smoke alone implies Fire, then $\text{Smoke} \wedge \text{Heat}$ also implies Fire                                                                                                                                                                                                                                                     |

|   | Formula                                                                                        | Answer               | Reason                                                                                                               |
|---|------------------------------------------------------------------------------------------------|----------------------|----------------------------------------------------------------------------------------------------------------------|
| f | $(\text{Fire} \rightarrow \text{Smoke})$<br>$\wedge \text{Fire} \wedge$<br>$\neg \text{Smoke}$ | <b>Unsatisfiable</b> | Fire=T forces Smoke=T (by first conjunct + modus ponens), but $\neg \text{Smoke}=T$ is also required — contradiction |

## Part 4 – Normal Forms

### 4.1 CNF conversions:

(i)  $\neg(s \rightarrow \neg p) \wedge (p \rightarrow \neg r)$

Step 1 — eliminate  $\rightarrow$ :  $\neg(\neg s \vee \neg p) \wedge (\neg p \vee \neg r)$

Step 2 — push  $\neg$  inward:  $(s \wedge p) \wedge (\neg p \vee \neg r)$

Step 3 — already CNF:  $\boxed{s \wedge p \wedge (\neg p \vee \neg r)}$

(ii)  $p \rightarrow (\neg q \vee (q \rightarrow \neg p))$

Simplify inner:  $q \rightarrow \neg p = \neg q \vee \neg p$ , so the right side becomes  $\neg q \vee \neg q \vee \neg p = \neg q \vee \neg p$ .

Formula:  $p \rightarrow (\neg q \vee \neg p)$

Expand biconditional:  $(p \rightarrow (\neg q \vee \neg p)) \wedge ((\neg q \vee \neg p) \rightarrow p)$

$= (\neg p \vee \neg q \vee \neg p) \wedge (q \wedge p \vee p)$

$= (\neg p \vee \neg q) \wedge (p \vee (q \wedge p))$  ... simplify:  $\neg p \vee \neg p = \neg p$

$= (\neg p \vee \neg q) \wedge p$

$\boxed{(\neg p \vee \neg q) \wedge p}$

(iii)  $\neg(p \rightarrow (\neg q \vee (r \rightarrow \neg p)))$

$\neg(\neg p \vee (\neg q \vee (r \rightarrow \neg p)))$

$= p \wedge \neg(\neg q \vee (r \rightarrow \neg p))$

$= p \wedge q \wedge \neg(r \rightarrow \neg p)$

$\neg(r \rightarrow \neg p) = (r \wedge p) \vee (\neg r \wedge \neg p)$  ... wait,  $\neg(X \rightarrow Y) = (X \wedge \neg Y) \vee (\neg X \wedge Y)$

$= (r \wedge \neg(\neg p)) \vee (\neg r \wedge \neg p) = (r \wedge p) \vee (\neg r \wedge \neg p)$

Distribute into CNF:  $(r \vee \neg r) \wedge (r \vee \neg p) \wedge (p \vee \neg r) \wedge (p \vee \neg p)$

Simplify tautological clauses:  $(r \vee \neg r) \wedge (p \vee \neg r)$

Full formula:  $\boxed{p \wedge q \wedge (r \vee \neg p) \wedge (p \vee \neg r)}$  — simplifies further to  $p \wedge q \wedge \neg r$  (since  $r \vee \neg p$  with  $p$  gives  $r$ ... but we also need  $\neg r$ , contradiction — formula is unsatisfiable if we trace through carefully).

4.2 DNF:  $(s \rightarrow \neg(\neg r \rightarrow q)) \rightarrow p$

Inner:  $\neg r \rightarrow q = r \vee q$ , so  $\neg(\neg r \rightarrow q) = \neg r \wedge \neg q$

$s \rightarrow (\neg r \wedge \neg q) = \neg s \vee (\neg r \wedge \neg q)$

Full:  $\neg(\neg s \vee (\neg r \wedge \neg q)) \vee p = (s \wedge (r \vee q)) \vee p$

Distribute:  $\boxed{(s \wedge r) \vee (s \wedge q) \vee p}$

## Part 5 – Python Truth Tables (Solution)

```

from itertools import product

def truth_table(variables, formula_fn):
    print(" | ".join(variables) + " | result")
    print("-" * (4 * len(variables) + 10))
    count = 0
    for vals in product([False, True], repeat=len(variables)):
        assignment = dict(zip(variables, vals))
        result = formula_fn(assignment)
        row = " | ".join("T" if v else "F" for v in vals)
        print(f"{row} | {'T' if result else 'F'}")
        if result:
            count += 1
    print(f"\nSatisfying models: {count}")

# Example: tautology from 1.2
truth_table(['A', 'B'], lambda m: (not m['A'] or m['B']) or (not m['B'] or m['A']))

# Part 2a: (A∧B)∨(B∧C) – with D free
truth_table(['A', 'B', 'C', 'D'],
    lambda m: (m['A'] and m['B']) or (m['B'] and m['C']))

```

## Part 6 – Python Resolution (Solution)

```

def resolve(c1, c2):
    """Return the resolvent of two clauses, or None if not resolvable."""
    for lit in c1:
        neg = lit[1:] if lit.startswith('¬') else '¬' + lit
        if neg in c2:
            resolvent = (c1 - {lit}) | (c2 - {neg})
            return resolvent
    return None

def pl_resolution(kb_clauses, goal_clause):
    """
    Try to prove goal_clause from kb_clauses via resolution refutation.
    Adds the negation of each literal in goal_clause and attempts to derive {}.
    """
    # Negate the goal: negate each literal and add as unit clauses
    negated = []
    for lit in goal_clause:
        neg = lit[1:] if lit.startswith('¬') else '¬' + lit
        negated.append(frozenset({neg}))

    clauses = set(frozenset(c) for c in kb_clauses) | set(negated)
    new = set()

    while True:
        pairs = [(c1, c2) for c1 in clauses for c2 in clauses if c1 != c2]
        for c1, c2 in pairs:
            resolvent = resolve(set(c1), set(c2))

```

```

    if resolvent is not None:
        if len(resolvent) == 0:
            return True # contradiction found → goal is proved
        new.add(frozenset(resolvent))
    if new.issubset(clauses):
        return False # nothing new → cannot prove goal
    clauses |= new

```

### Unicorn problem in CNF:

|                                             |                                                    |
|---------------------------------------------|----------------------------------------------------|
| Mystical $\rightarrow$ Immortal             | = { $\neg$ Mystical, Immortal}                     |
| $\neg$ Mystical $\rightarrow$ Mammal        | = {Mystical, Mammal}                               |
| (Immortal $\vee$ Mammal) $\rightarrow$ Horn | = { $\neg$ Immortal, Horn}, { $\neg$ Mammal, Horn} |
| Horn $\rightarrow$ Magical                  | = { $\neg$ Horn, Magical}                          |

Goal: {Magical} — resolution derives the empty clause, proving Magical. ✓

## Part 7 – FOL Formula Matching

| # | Formula                                              | Answer                                                                                                                                                                                                |
|---|------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1 | $\exists x. (\text{big}(x) \wedge \text{cat}(x))$    | (F) There exists a big cat                                                                                                                                                                            |
| 2 | $\exists x. (\text{big}(x) \wedge \text{cat}(x))$    | (B) There exists a small thing or a cat                                                                                                                                                               |
| 3 | $\exists x. (\text{big}(x) \vee \text{cat}(x))$      | (H) There exists a big thing or a cat                                                                                                                                                                 |
| 4 | $\forall x. (\text{big}(x) \wedge \text{cat}(x))$    | (D) All things are big cats                                                                                                                                                                           |
| 5 | $\forall x. (\text{big}(x) \wedge \text{cat}(x))$    | (C) All big things are cats                                                                                                                                                                           |
| 6 | $\forall x. (\text{big}(x) \vee \text{cat}(x))$      | (A) All cats are big ← Note: this is actually "everything is big or a cat." Closest is (A) by elimination                                                                                             |
| 7 | $\forall x. (\text{big}(x) \vee \neg \text{cat}(x))$ | (E) All cats are small ← $\neg \text{cat}(x) \vee \text{big}(x) = \text{cat}(x) \rightarrow \text{big}(x)$ , so (A); #7 = $\text{cat} \rightarrow \text{big} = \text{A}$ , #6 = everything big or cat |

### Careful note on 6 and 7:

- Formula 6:  $\forall x. (\text{big}(x) \vee \text{cat}(x))$  — everything is either big or a cat
- Formula 7:  $\forall x. (\text{big}(x) \vee \neg \text{cat}(x)) = \forall x. (\text{cat}(x) \rightarrow \text{big}(x)) = \text{(A)}$   
**All cats are big**
- Formula 6 then maps to the remaining sentence: "everything is big or a cat" — not listed precisely, closest is none but by elimination maps to (G) is wrong. The sentence set has (E) **All cats are small** =  $\forall x. (\text{cat}(x) \rightarrow \neg \text{big}(x))$  which matches no formula above. Formula 6  $\rightarrow$  (none)

**perfectly**); in exam context: **7 → A, 6 → unlisted** (exercise has 7 formulas, 8 sentences — one sentence is a distractor).

---

## Part 8 – FOL Translation Critique

---

**(a) Incorrect.** The inner implication  $\text{In}(x, \text{Starnberg}) \rightarrow \text{Rent}(x) < \text{Rent}(y)$  uses  $x$  instead of  $y$  for Starnberg. Also, using  $\rightarrow$  inside  $\exists y$  is too weak — it is satisfied vacuously if  $y$  is not in Starnberg. The correct form needs  $\wedge$ :

$\forall x. (\text{App}(x) \wedge \text{In}(x, \text{Munich}) \rightarrow \exists y. (\text{App}(y) \wedge \text{In}(y, \text{Starnberg}) \wedge \text{Rent}(x) < \text{Rent}(y)))$

**(b) Correct.** The  $\exists x$  picks one apartment, and the  $\forall y$  says any other apartment in Starnberg below 500€ must be the same one. This correctly encodes uniqueness (the standard "exactly one" pattern).

**(c) Incorrect.** The parenthesization is wrong — the  $\rightarrow \text{In}(x, \text{Starnberg})$  is inside the  $\forall y$  scope rather than being the conclusion of the outer  $\forall x$ . The correct reading of the formula is ambiguous/broken. The intended formula should be:

$\forall x. (\text{App}(x) \wedge (\forall y. (\text{App}(y) \wedge \text{In}(y, \text{Munich}) \rightarrow \text{Rent}(x) > \text{Rent}(y)) \rightarrow \text{In}(x, \text{Starnberg})))$

---

End of solutions. 🎓