Hong Kong Exchanges and Clearing Limited and The Stock Exchange of Hong Kong Limited take no responsibility for the contents of this announcement, make no representation as to its accuracy or completeness and expressly disclaim any liability whatsoever for any loss howsoever arising from or in reliance upon the whole or any part of the contents of this announcement.

SHANGHAI JUNSHI BIOSCIENCES CO., LTD.*
上海君實生物醫藥科技股份有限公司

(a joint stock company incorporated in the People's Republic of China with limited liability)

(Stock code: 1877)

VOLUNTARY ANNOUNCEMENT – APPROVAL OF THE SUPPLEMENTAL NEW DRUG APPLICATION FOR ONGERICIMAB INJECTION

This announcement is made by Shanghai Junshi Biosciences Co., Ltd.* (上海君實生物醫藥科技股份有限公司) (the "**Company**") on a voluntary basis. Reference is also made to the overseas regulatory announcement of the Company dated 27 May 2025.

The board (the "**Board**") of directors (the "**Directors**") of the Company is pleased to announce that the Company has received the Drug Registration Certificate* (《藥品註冊證書》) issued by the National Medical Products Administration (the "**NMPA**"). The two supplemental new drug applications for the ongericimab injection (a recombinant humanized anti-PCSK9 monoclonal antibody injection, trade name: JUNSHIDA (君適達®)) for: 1) adult patients with heterozygous familial hypercholesterolemia ("**HeFH**"); 2) alone or in combination with ezetimibe, in adult patients with non-familial hypercholesterolemia and mixed dyslipidemia who are statin-intolerant or statins contraindicated, have been approved. Ongericimab has become the first domestic PCSK9-targeted drug approved for statin-intolerant patients.

ABOUT ONGERICIMAB

Drug name: Ongericimab injection*(昂戈瑞西單抗注射液)
Application matter: Registration of Pharmaceutical Product (Domestic Production)
Acceptance Nos.: CXSS2400028, CXSS2400029, CXSS2400030, CXSS2400031
Certificate Nos.: 2025S01445, 2025S01446
Marketing Authorization Holder: Shanghai Junshi Biosciences Co., Ltd.*
Specification: 150 mg (1 ml) in a single dose (pre-filled syringe), 150 mg (1 ml) in a single dose (pre-filled autosyringe)
Indications: On the basis of dietary control, in combination with statins, or statins and other lipid-lowering therapies, for the treatment of adult patients with primary hypercholesterolemia (including heterozygous familial and non-familial hypercholesterolemia) and mixed dyslipidemia, who fail to achieve low-density lipoprotein cholesterol (LDL-C) goals after receiving moderate or higher doses of statins treatment; or alone or in combination with ezetimibe, in adult patients with non-familial hypercholesterolemia and mixed dyslipidemia who are statin-intolerant or statins contraindicated, to reduce low-density lipoprotein cholesterol (LDL-C), total cholesterol (TC) and apolipoprotein B (ApoB) levels.

According to the Chinese Guidelines for Lipid Management (2023), cardiovascular disease is the leading cause of death among both urban and rural residents in China, with atherosclerotic cardiovascular disease ("**ASCVD**") being the predominant subtype. The rise of low-density lipoprotein cholesterol ("**LDL-C**") levels is a major cause of ASCVD. Lowering LDL-C levels can significantly decrease the incidence of ASCVD and risk of death.

HeFH is a common type of familial hypercholesterolemia with an estimated prevalence of 1:250 – 1:200, and its main clinical features are significantly elevated LDL-C and early onset of coronary artery disease. Compared to patients with non-familial hypercholesterolemia, patients with HeFH exhibit higher baseline LDL-C levels and lower target levels for control recommended by the guidelines. Failure to achieve target LDL-C levels after treatments such as statins will result in patients being at high cardiovascular risk.

In addition, despite statins currently being the cornerstone of lipid lowering treatment, approximately 9.1% of patients clinically exhibit statin intolerance, with a higher proportion observed in Asian populations. Discontinuation of statins or the use of only tolerable doses in patients with statin intolerance may lead to suboptimal LDL-C levels, thereby failing to effectively reduce the patient's ASCVD risk.

As a new lipid-lowering drug that effectively reduces LDL-C levels, PCSK9 inhibitor has been recommended in the guidelines for the management of lipids both in China and overseas and is widely recognized by clinicians. The significant lipid-lowering effects of ongericimab have been demonstrated in multiple phase III clinical studies, and ongericimab was approved by the NMPA for the treatment of adult patients with primary hypercholesterolemia (non-familial) and mixed dyslipidemia in October 2024.

The approval of the two supplemental new drug applications are mainly based on two registered clinical trials – JS002-005 (NCT05325203) and JS002-007 (NCT05621070).

JS002-005 is a randomized, double-blind, placebo-controlled Phase III clinical study in adult patients with HeFH. JS002-005 was led by Professor Ma Changsheng (馬長生) and Professor Lin Jie (藺潔) from Beijing Anzhen Hospital affiliated to Capital Medical University* (首都醫科大學附屬北京安貞醫院) as the principal investigators. A total of 135 patients with HeFH were enrolled. This study is the first Phase III clinical study evaluating an anti-PCSK9 monoclonal antibody in Chinese patients diagnosed with HeFH (DLCN>8). The latest data on JS002-005 has been published in full in *Atherosclerosis*. The results showed that, compared with placebo, the ongericimab subcutaneous injection (150 mg every 2 weeks (Q2W) or 450 mg every 4 weeks (Q4W)) significantly reduced LDL-C levels by 69.4% and 80.6% ($p<0.0001$), respectively, for a 24-week treatment, with steady reduction during the treatment. At the same time, ongericimab also demonstrated significant improvements in other lipid parameters. Non-high density lipoprotein cholesterol (non-HDL-C), apolipoprotein B (ApoB), total cholesterol (TC), and lipoprotein (a) ("**Lp(a)**") levels all markedly decreased from baseline. Notably, Lp(a) levels decreased by 50% from baseline. The overall safety profile was favorable, with the incidence of treatment-emergent adverse events (TEAEs) being comparable to that of the placebo group.

JS002-007 is a randomized, double-blind, placebo-controlled Phase III clinical study conducted in adult patients with primary hypercholesterolemia and mixed hyperlipidemia who are statin-intolerant. JS002-007 was led by Professor Tang Yida (唐熠達) from Peking University Third Hospital* (北京大學第三醫院) as the principal investigator. This is the first Phase III clinical study of anti-PCSK9 monoclonal antibody in Chinese patients with primary hypercholesterolemia and mixed dyslipidemia who are statin-intolerant. The results will be published in an international academic journal shortly.

Ongericimab injection is a recombinant humanized anti-PCSK9 monoclonal antibody injection independently developed by the Company. It was approved for marketing by the NMPA in October 2024. As of the date of this announcement, ongericimab injection has been approved for three indications in Chinese mainland for the treatment of: 1) adult patients with primary hypercholesterolemia (non-familial) and mixed dyslipidemia; 2) adult patients with HeFH; 3) alone or in combination with ezetimibe, in adult patients with non-familial hypercholesterolemia and mixed dyslipidemia who are statin-intolerant or statins contraindicated. The approved specifications are 150 mg (1 ml) in a single dose (pre-filled syringe), 150 mg (1 ml) in a single dose (pre-filled autosyringe).

In October 2023, the Company signed an agreement with Chongqing Bochuang Pharmaceuticals Co., Ltd.* (重慶博創醫藥有限公司) ("**Bochuang Pharmaceuticals**"), pursuant to which the Company granted Bochuang Pharmaceuticals an exclusive license to conduct research and development on, manufacture and commercialize, ongericimab for the licensed purposes and within Chinese Mainland. Bochuang Pharmaceuticals will be responsible for the subsequent commercialization of ongericimab in Chinese Mainland and will make corresponding milestone payments and sales commissions to the Company.

RISK WARNING

Due to the high-tech, high-risk and high-value-added characteristics of pharmaceutical products, there are substantial risks and uncertainties in the process of commercialization of pharmaceutical products. Investors are advised to make cautious decisions and pay careful attention to investment risks. The Company will actively pursue the described project and fulfill its information disclosure obligations in a timely manner for subsequent progress in strict accordance with relevant regulations.

By order of the Board
Shanghai Junshi Biosciences Co., Ltd.*
Mr. Xiong Jun
Chairman

Shanghai, the PRC, 27 May 2025

As at the date of this announcement, the Board of Directors of the Company comprises Mr. Xiong Jun, Dr. Li Ning, Dr. Zou Jianjun, Mr. Li Cong, Mr. Zhang Zhuobing, Dr. Yao Sheng, Dr. Wang Gang and Dr. Li Xin as executive Directors; Mr. Tang Yi as a non-executive Director; and Mr. Zhang Chun, Dr. Feng Xiaoyuan, Dr. Yang Yue, Mr. Li Zhongxian and Ms. Lu Kun as independent non-executive Directors.

* *For identification purpose only*