



Conference Abstract

Building Essential Biodiversity Variable netCDFs with the ebvcube R Package

Luise Quoß^{‡,§}, Néstor Fernández^{‡,§}, Christian Langer^{‡,§}, Jose Valdez^{‡,§}, Miguel Alejandro Fernández ^{‡,§}, Henrique M. Pereira^{‡,§}

- ‡ German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany
- § Institute of Biology, Martin Luther University Halle Wittenberg, Halle, Germany

Corresponding author: Luise Quoß (luise.quoss@idiv.de)

Received: 03 Aug 2022 | Published: 23 Aug 2022

Citation: Quoß L, Fernández N, Langer C, Valdez J, Fernández MA, Pereira HM (2022) Building Essential Biodiversity Variable netCDFs with the ebvcube R Package. Biodiversity Information Science and Standards 6:

e91215. https://doi.org/10.3897/biss.6.91215

Abstract

The concept of <u>Essential Biodiversity Variables</u> (EBVs) was conceived to study, report, and manage biodiversity change. The EBV netCDF structure was developed in order to support publication and interoperability of biodiversity data. This standard is based on the <u>Network Common Data Format</u> (netCDF). Additionally, it follows the <u>Climate and Forecast Conventions</u> (CF, version 1.8) and the <u>Attribute Convention for Data Discovery</u> (ACDD, version 1.3).

The standard allows several datacubes per netCDF file (see Fig. 1). These cubes have four dimensions: longitude, latitude, time and entity, whereby the last dimension can, for example, encompass different species or groups of species, ecosystem types or other aspects. The usage of hierarchical groups enables the coexistence of multiple EBV cubes (see Fig. 2). The first level (netCDF group) are scenarios, e.g., the modelling for different Shared Socioeconomic Pathways (SSP) scenarios. The second level (netCDF group) are metrics, e.g., the percentage of protected area per pixel and its proportional loss over a certain time span per pixel. All metrics are repeated per scenario, if any are present. The result is a rather complex raster dataset (see example dataset in Fig. 3).

2 Quoß L et al




Figure 1.

Abstract visualization of the EBV cube (Mahecha 2017, License CC BY 4.0).

```
ebv-dataset
                                                ebv-dataset
 _global attributes
                                                  _global attributes
                                                  _dimensions [lon, lat, time, entity]
  _dimensions [lon, lat, time, entity]
                                                  (coordinate) variables
  _(coordinate) variables
                                                   _scenario_1
  _metric_1
                                                   group attributes
   group attributes
                                                    _metric_1
   _ebvcube [lon, lat, time, entity]
                                                      group attributes
     __datacube attributes
                                                      ∟ebvcube [lon, lat, time, entity]
                                                        __datacube attributes
                                                     metric_2
                                                   scenario_2
(a)
                                                 (b)
```

Figure 2.

EBV netCDF hierarchical data structure: (a) shows the structure of a minimal dataset and (b) shows the structure of an exhaustive dataset.

```
mammals dataset
__title: InSiGHTS_LUH2_2015_2055
__summary: Data on Area Of Habitat (AOH) for 100
  mammals from 2015 to 2055, in 5 year intervals
  _ebv_class: Species populations
 __ebv_name: Species distributions
  _dimensions [lon, lat, time, entity]
  Llat attributes
  └_lon attributes
  L time attributes
  entities
  __entities attributes
  __crs attributes
  scenario_1
   standard name: Sustainability
   long_name: SSP1-RCP2.6
   __metric_1
      __standard_name: Habitat availability
       _long_name: absolute values per 5 years and
      species - in km2
       _ebvcube [720,1 440, 9, 100]
        _units: land-use of mammals calculated in km2
        grid_mapping: crs
   __standard_name: Middle of the Road
   long_name: SSP2-RCP4.5
   __metric_1
     group attributes
  scenario_3
  scenario 4
  ⊥...
  scenario_5
```

Figure 3.

EBV netCDF Structure of the Global habitat availability for mammals <u>dataset</u> by Daniele Baisero (License CC BY 4.0).

Blue elements are variables, green elements represent groups, attributes are displayed in black and dimension are red.

This is where the <u>ebvcube R package</u> comes into play. This R package enables scientists to create their own netCDFs in the EBV cube standard. Its functionality covers the creation, opening/reading and visualizing the EBV netCDFs. The ebvcube package is part of the overall EBV infrastructure and works together with the <u>EBV Data Portal</u>. Users can work with the downloaded EBV netCDFs or upload their own EBV netCDFs to the portal.

Generally, the package aims to condense the output for the users and assist in the understanding of the file structure to overcome the complexity. The output is reduced to the necessary information, e.g., not displaying coordinate variables or any technical attributes. Moreover, functionality for a quick data exploration is implemented.

Keywords

GEO BON, data standard, EBV, monitoring, interoperability, FAIR, data portal

4 Quoß L et al

Presenting author

Luise Quoß

Presented at

TDWG 2022

References

 Mahecha M (2017) Earth System Data Cube. figshare. Figure. https://doi.org/10.6084/ m9.figshare.4822930.v2 © 2022. This work is published under http://creativecommons.org/licenses/by/4.0/ (the "License"). Notwithstanding the ProQuest Terms and Conditions, you may use this content in accordance with the terms of the License.