A temporally-constrained latent space reveals the stochastic dynamics of convective organization

Sophie Abramian,[1] Olivier Pauluis,[2] L. Ruby Leung,[3] and Pierre Gentine[1]

[1] *Department of Earth and Environmental Engineering, Columbia University, New York, NY, USA*
[2] *Courant Institute of Mathematical Sciences, New York University, New York, NY, USA*
[3] *Atmospheric Sciences and Global Change Division, Pacific Northwest National Laboratory, Richland, WA, USA*

(Dated: July 27, 2026)

Atmospheric convection is intrinsically stochastic: under identical large-scale forcing, storms can follow widely different life cycles. We introduce an unsupervised framework that learns a low-dimensional representation of cloud-resolving squall-line simulations by constraining the latent space to evolve according to a linear stochastic operator. The learned representation partitions convection into five physically interpretable states whose transitions define a Markov chain. Its spectrum reveals a dominant slow mode separating organized from unorganized convection and governing the system's life cycle. The remaining departures from Markovianity are partly explained by cold pools, identifying a physical source of memory absent from the reduced description. The framework provides a physically interpretable stochastic coarse-graining of convection and offers an alternative to existing multi-cloud parameterizations.

Discovering interpretable, low-dimensional representations of high-dimensional stochastic ~~dynamics~~ systems is a foundational challenge across the physical sciences. Deep convection ~~in the atmosphere~~ is a prime example. ~~Deep~~ : deep convective systems—organized cloud structures larger and longer-lived than individual cells—dominate the tropical water and energy cycle [1, 2]; the largest ~~, mesoscale convective systems,~~ contribute over 50% of tropical precipitation and a disproportionate share of extremes [3–5].

In climate models they occur below the grid scale and must be ~~parameterized a~~ parameterized, a persistent source of uncertainty ~~, especially~~ under warming [6, 7]. The difficulty is that a convective system is ~~a~~ high-dimensional~~dynamical object;~~ : compressing it into a few prognostic variables without discarding what matters is hard.

Compounding this, ~~the convective response to a~~ under *fixed* large-scale forcing the convective response is not a single outcome but a broad distribution of storm evolutions [8–11]—an *internal* variability intrinsic to ~~the convection, not inherited from the environment~~convection. Capturing it requires a reduced description ~~that is~~ both low-dimensional and faithful to the dynamics.

Stochastic multi-cloud models [12–14] offer such a reduction, but ~~a strongly simplified~~ one whose states are prescribed *a priori*: a lattice of sites, each in one of a few hand-chosen cloud states—clear sky, shallow, congestus, or deep—whose area fractions evolve by a Markov chain. This captures the changing cloud cover but not the vertical mass-flux profiles that also govern extreme precipitation [15–19].

Koopman operator theory [20, 21] offers a principled alternative. Instead of the state $\mathbf{x}_t$ of a high-dimensional nonlinear system, one follows *observables* $g(\mathbf{x})$ that recast it in new coordinates; the operator advancing them—$\mathcal{K}$, the Koopman operator, defined by $(\mathcal{K}g)(\mathbf{x}_t) = g(\mathbf{x}_{t+1})$—is linear by construction, whatever the nonlinear dynamics. The challenge is ~~thus~~ then not to resolve the nonlinear evolution ~~,~~ but to find an observable that simplifies ~~it ideally one on which $\mathcal{K}$ acts as a simple multiplication while~~ it while retaining most of the system's information. Such an observable is generally infinite-dimensional [21]—the long-standing obstacle. Only recently have machine-learning encoder–decoder architectures [22, 23] overcome it, learning from data the best finite-dimensional observable for a given latent dimension together with its linear operator.

We depart from existing Koopman autoencoders in two ways. First, we start from a physically reduced representation~~of the system~~, casting the flow in isentropic coordinates [24]—a thermodynamically consistent reduction that filters reversible adiabatic motions—rather than ~~using the raw , noisy~~ raw three-dimensional fields. Second, rather than enforcing *deterministic* linear dynamics, we *partition* the latent space into discrete states by joint clustering [25, 26] and model their transitions *probabilistically*. The result is a Markov chain ~~on data-discovered states finite-dimensional, stochastic, and shaped by the dynamics it represents~~whose states are discovered from the data rather than prescribed. Though demonstrated on convection, the framework applies to any high-dimensional system where regime discovery and transition modeling are needed.

We analyze cloud-resolving simulations of squall lines (SAM [27]) under fixed vertical wind shear, in statistical equilibrium yet with strong internal variability. ~~Convection is cast in isentropic coordinates: at~~ At each time the three-dimensional vertical mass flux is projected onto height z and moist static energy h, yielding a two-dimensional diagram $M(h, z)$ that ~~cleanly~~ sep-




FIG. 1. ~~Three contrasting snapshots of the simulated convective field.~~ Three contrasting snapshots of the convective field. **Left (a, c, e):** isovolume of non-precipitating condensate QN = 0.02 $\mathrm{kg\,kg^{-1}}$ over the surface temperature anomaly; white patches mark cold pools. **Right (b, d, f):** ~~the corresponding~~ isentropic mass-flux diagram $M(h, z, t)$, red (blue) for upward (downward) transport. Rows: **(a–b)** suppressed, **(c–d)** organized deep, ~~and~~ **(e–f)** unorganized active~~convection~~.

arates high-h updrafts from the low-h evaporative downdrafts that build cold pools [24]. Figure 1 shows three representative ~~snapshots—suppressed, organized-deep, and unorganized-active convection—alongside~~ snapshots alongside their mass-flux diagrams; this time series ~~of diagrams~~ is the input to the model.

Applied to these simulations, the construction asks: (i) what mass-flux states a given forcing makes accessible; (ii) whether their transitions form a Markov chain, and how organized and weakly organized pathways differ; (iii) where internal feedbacks—cold pools in particular—introduce memory beyond the discrete state. Five interpretable states emerge without supervision, and their transitions form a Markov chain with a slow (∼3 h) eigenmode alternating two regimes, organized and unorganized; two long-lived states decay more slowly than their Markov prediction, and conditioning on cold-pool intensity closes ∼30% of that gap.

CONSTRUCTION AND PHYSICAL INTERPRETATION OF THE LATENT SPACE

The squall-line evolution is seen through its time series of isentropic mass-flux fields X_t. ~~The model compresses each field into a latent vector $\mathbf{z}_t \in \mathbb{R}^8$ and groups it into five discrete states. Specifically, an~~ An encoder maps X_t to a Gaussian posterior over ~~$\mathbf{z}_t$~~ a latent vector $\mathbf{z}_t \in \mathbb{R}^8$ (a variational autoencoder with reconstruction and Kullback–Leibler terms); a softmax head turns $\mathbf{z}_t$ into a probability vector s_t over ~~the~~ $K = 5$ states; and a row-stochastic matrix $\mathbf{T}$ propagates it one step, $s_{t+1} \approx \mathbf{T}\, s_t$. ~~These three components~~ The three are trained jointly on consecutive pairs $(X_t, X_{t+\Delta t})$, so the latent space is at once reconstructive, cluster-separable, and dynamically predictive (Fig. 2). A contrastive ~~loss based on the structural~~ (Dice) ~~similarity of~~ loss on the mass-flux fields prevents state collapse, keeping all five states populated. Full ~~architecture, losses, and training are given~~ details are in the Supplemental Material.




FIG. 2. **Model architecture.** A variational autoencoder ~~(VAE)~~ is coupled to a Markovian state-assignment and transition framework ~~. An encoder maps the input to a latent representation z_t, reconstructed by a decoder; a small network assigns each z_t a probability distribution over $K = 5$ discrete states;~~ and ~~a linear transition operator T predicts~~ trained jointly (components defined in the ~~state distribution at $t + 1$ from that at t~~text).

No physical constraint is imposed during ~~training—the~~ training: the VAE is regularized only toward an isotropic Gaussian ~~,~~ and states are assigned without reference to ~~intensity, organization, or any diagnostic—so whether the~~ any physical diagnostic, so a physically meaningful latent space is ~~physically meaningful is~~ not guaranteed. Applying PCA to $\mathbf{z}(t)$, the first two components explain 22% and 17% of the variance (Fig. 3); a three-dimensional projection (Supplemental Material, Fig. **??**) shows that states overlapping in 2D separate along higher-order components.

The dominant latent directions are physically interpretable (Fig. 3). The first principal component




FIG. 3. **Latent-space visualization and physical interpretation.** PCA projection of the latent space (each point is one time step), coloured by **(a)** state assignment, with two example trajectories over 5 h (dashed: strongly organized; dotted: weakly organized~~; time series in the Supplemental Material~~); **(b)** domain-mean precipitation, a proxy for convective intensity; and **(c)** the variance of precipitable water (the PW index), a proxy for spatial organization (higher values indicate more clustered convection). ~~The first component (PCA 1) aligns with intensity, the second (PCA 2) with organization; joint~~ Joint distributions ~~of the two components against these diagnostics~~ are given in the Supplemental Material (Fig. **??**).

correlates strongly with domain-mean precipitation—convective intensity and life-cycle stage, from suppressed to vigorously precipitating. The second is nearly uncorrelated with precipitation but strongly anticorrelated with the variance of precipitable water, a proxy for spatial organization. Intensity and organization thus emerge as two largely independent axes of the learned representation.

The five states are also dynamically and thermodynamically distinct. Averaging the isentropic mass flux within each state as anomalies from the simulation-wide mean (definitions and full profiles in the Supplemental Material) shows that the organized-deep and stratiform-like states carry deep, high–moist-static-energy ~~ascent the stratiform-like state additionally showing a mid-level mass-flux signature suggestive of stratiform detrainment while~~ ascent, while the active-unorganized state peaks at lower altitude and lower moist static energy, and the suppressed and shallow-congestus states carry only weak, shallow anomalies. Table I summarizes each state.

MARKOV TRANSITION MATRIX, EIGENMODES, AND REGIMES

The second output of the model is a Markov transition matrix $\mathbf{T}$, whose elements

$$T_{ij} = \mathbb{P}\left(X_{t+\Delta t} = j \mid X_t = i\right) \tag{1}$$

denote the probability of transitioning from state i to state j. The resulting transition matrix is shown in Fig. 4.

The matrix $\mathbf{T}$ is strongly diagonally ~~dominated, indicating that the system tends to persist within a given~~ dominant and sparse: the system persists within a state over several ~~time steps. Moreover, the matrix is sparse,~~ steps, with only a ~~limited number of~~ few preferred transitions between states~~, suggesting constrained pathways in state space~~.

To ~~further~~ characterize the dynamics, we ~~perform a spectral decomposition of~~ spectrally decompose $\mathbf{T}$. ~~Because $\mathbf{T}$ is~~ Being stochastic, it admits a unit eigenvalue $\lambda_1 = 1$ ~~. The corresponding~~ whose right eigenvector $\boldsymbol{\pi}$ satisfies $\mathbf{T}\boldsymbol{\pi} = \boldsymbol{\pi}$~~and defines~~, the stationary distribution~~of the Markov chain~~. We obtain

$$\boldsymbol{\pi} = \begin{bmatrix} 12.9\% & 12.2\% & 37.0\% & 22.7\% & 15.2\% \end{bmatrix}. \tag{2}$$

The stationary distribution is dominated by state 3 (37%~~, suppressed convection~~), the remaining weight spread across the four active ~~states all~~ states, all sufficiently populated to constrain the ~~inferred~~ transition statistics.

~~We next examine the subdominant eigenvalues and eigenvectors of $\mathbf{T}$. All~~ The non-stationary eigenvalues ~~have imaginary parts at the level of numerical noise~~ are effectively real ($|\mathrm{Im}(\lambda_k)| < 10^{-3}$)~~; the spectrum is effectively real. This indicates that the dominant dynamics cannot be described as simple linear~~, so the dominant dynamics are not simple oscillations in state space.

State	Name	π (%)	Physical signature
1	*Organized deep*	12.9	sustained deep ascent in a moist environment; reduced low-level detrainment; highest precipitable-water variance
2	*Stratiform-like*	12.2	strongly active convection with large upper-tropospheric mass flux, consistent with a mesoscale stratiform region; highest precipitation
3	*Suppressed convection*	37.0	weak up- and down-drafts; dry background atmosphere; dominant state
4	*Active unorganized*	22.7	active but unorganized convection, with mass flux peaking at lower altitudes than in the organized regime; drier environment
5	*Shallow congestus*	15.2	suppressed deep convection with residual shallow convective activity; dry environment

TABLE I. **Summary of the five latent convective states.** ~~State identities~~ Identities are inferred from the latent-space axes ~~of variability~~ (Fig. 3) and confirmed by ~~the~~ per-state ~~vertical and moist-static-energy~~ mass-flux profiles (Supplemental Material). π_i is the stationary probability of state i [Eq. (2)].




FIG. 4. **Markov state dynamics in latent space. (a)** ~~Latent space~~ Latent-space visualization~~with the inferred Markov chain overlaid. The size of~~ ; each state node is ~~proportional to~~ sized by its ~~asymptotic~~ (stationary ~~)~~ probability~~, and arrows indicate transition probabilities between states~~. **(b)** Transition matrix ~~describing the dynamics of convection~~ for the given shear. Two coupled blocks, separated by state 3, form the organized (states 1–2) and unorganized (states 4–5) regimes.

The leading nontrivial eigenvalue ~~of the transition matrix~~ is $\lambda_2 \approx 0.94$, ~~indicating~~ a highly persistent ~~dynamical~~ mode. The corresponding eigenvector

$$\mathbf{v}_2 = \begin{bmatrix} -0.56 & -0.30 & -0.25 & +0.65 & +0.34 \end{bmatrix} \qquad (3)$$

is unit-norm (the overall sign is arbitrary). A positive $(\mathbf{v}_2)_i$ means state i is enhanced relative to π_i in one phase ~~of this mode~~ and suppressed in the other~~;~~ , with $|(\mathbf{v}_2)_i|$ ~~sets the relative size of that deviation across states. Components of opposite sign therefore~~ its relative amplitude; opposite signs vary in anti-phase.

The components of $\mathbf{v}_2$ split the chain into two opposite-sign groups, $\{1, 2\}$ $(-0.56, -0.30)$ and $\{4, 5\}$

(+0.65, +0.34); state 3 has the smallest amplitude in this mode (−0.25). We interpret these two groups as two convective regimes—*organized* ($\{1,2\}$) and *unorganized* ($\{4,5\}$)—that the chain alternates between on the eigenmode decay time $\tau_2 = -\Delta t/\log|\lambda_2| \approx 172$ min. State 3, despite its small amplitude in $\mathbf{v}_2$, carries the dominant stationary occupancy ($\pi_3 = 37\%$), and nearly every transition between the other four states passes through it (Fig. 4b); it therefore , so it functions as the chain's intermediate state, visited routinely between successive episodes in the other states.

The internal dynamics of the two regimes differ. The unorganized regime is a closed oscillation: states 4 and 5 exchange directly and frequently ($T_{4\to5} = 0.09$, $T_{5\to4} = 0.14$). The organized regime is not. States: states 1 and 2 do not exchange directly ($T_{1\to2} = T_{2\to1} = 0.01$) , and they are also separated in time: and are separated in time—200 min after a state-1 timestep, the conditional probability of being in state 2 is 0.04, well below the stationary value $\pi_2 = 0.12$. States 1 (deep moist ascent, reduced low-level detrainment) andstateand 2 (mesoscale stratiform-like deep convection) therefore appear as independent organized episodes, each returning to state 3 before the next can begin. Time series of individual Individual realizations (Supplemental Material, Fig. **??**) reflect this: organized intervals are dominated by a $3 \leftrightarrow 1$ oscillationwith occasional state-2 excursions, while unorganized intervals follow , unorganized intervals by the $3 \leftrightarrow 5 \leftrightarrow 4$ cycle.

The eigenvalue spectrum just identified spectrum makes a quantitative prediction: for a true Markov chain, the autocorrelation of every each state's occupancy must be is a sum of exponentials with the decay times $\tau_k = -\Delta t/\log|\lambda_k|$. We next test this prediction directly , which we now test against the empirical autocorrelations, identifying which states deviate from Markovian behavior and how those deviations reveal internal feedbacks such as cold-pool memory.

MARKOVIANITY ASSESSMENT AND INTERNAL FEEDBACKS

We now ask whether the five-state chain is self-contained—whether self-contained: whether the state at time t alone determines the distribution at $t+\tau$. Departures from this Markov property identify slow internal processes (moisture processes—moisture memory, cold pools, mesoscale organization) that organization—that the partition does not resolve. The question is long-standing: classical convective parameterizations assume instantaneous quasi-equilibrium adjustment [28, 29], while more recent schemes introduce recent schemes add explicit memory variables to capture for lagged responses [30–33].

Indicator autocorrelation.—Defining the occupancy indicator $I_i(t) = \mathbf{1}[X_t = i]$ and its normalized autocorrelation

$$C_i(\tau) = \frac{\mathbb{E}[(I_i(t)-\pi_i)(I_i(t+\tau)-\pi_i)]}{\mathbb{E}[(I_i(t)-\pi_i)^2]}, \tag{4}$$

a discrete-time Markov chain with transition matrix $\mathbf{T}$ and stationary distribution $\boldsymbol{\pi}$ admits the closed form [26]

$$C_i(\tau) \;=\; \frac{T^{\tau}_{ii}-\pi_i}{1-\pi_i} \;=\; \sum_{\alpha=2}^{K} A^{(i)}_{\alpha}\,\lambda^{\tau}_{\alpha}, \tag{5}$$

a superposition of exponentially decaying modes set by the eigenvalues λ_α of $\mathbf{T}$, with state-dependent amplitudes $A^{(i)}_\alpha = (\mathbf{v}_\alpha)_i(\mathbf{w}_\alpha)_i/(1-\pi_i)$, where ($\mathbf{v}_\alpha$and , $\mathbf{w}_\alpha$ are the right and left eigenvectorsof $\mathbf{T}$ (; derivation in the Supplemental Material). Departures from this prediction it quantify residual memory not encoded by the partition.

Memory beyond Markov in the convection-active states.—In Fig. 5a, the empirical autocorrelations decrease and fall below 0.2 within the lag window, in line with the Markov prediction of Eq. (5)—except for the organized state 1 (blue) and the active-unorganized state 4 (yellow), which stay elevated, well above their predictions; state 3 (suppressed convection) tracks its prediction most closely. This distinction matters: a state can be *sustained* (long dwell times) and yet still *Markovian* if its diagonal T_{ii} alone explains the persistence; states 1 and 4 are sustained persist *beyond* what T_{ii} predicts, indicating that so an internal degree of freedom not represented in absent from the partition is keeping them alive.

Cold pools partly close the gap.—Cold pools—masses of cold, dense air that form formed below convective clouds through by rain evaporation and are a key control on squall-line organization [34–37]—are a natural candidate carrier of this memory: they outlast the downdrafts that generate them and trigger or suppress new convection at their gust fronts [38], so they can sustain sustaining an organized state beyond what the instantaneous partition records. The hypothesis is testable: if cold pools carry the memory, then knowing the cold-pool field at the starting time should make the subsequent state 1 dynamics more Markovian—faster autocorrelation decay, closer to the Markov predictionMarkovian. To test this we introduce an indicator that picks out state 1 timesteps with a high cold pool,

$$J^{\star}_1(t) \;=\; \mathbf{1}[X_t = 1 \text{ and } \mathrm{CP}(t) > \mathrm{CP}_{\star}], \tag{6}$$

where $\mathrm{CP}(t)$ is a cold-pool index—the total area covered by cold pools at time t, normalized by its temporal mean (Supplemental Material)—and $\mathrm{CP}_\star$ is its 75th percentile. The cold-pool-conditioned state 1 survival is then the expected state 1 occupancy a lag τ later, averaged over

the timesteps selected by $J_1^\star$:

$$S_1^{(\mathrm{CP})}(\tau) \;=\; \mathbb{E}\big[\, I_1(t+\tau) \,\big|\, J_1^\star(t) = 1 \,\big]\,. \qquad (7)$$

We then rescale to autocorrelation form via $C_1^{(\mathrm{CP})}(\tau) = (S_1^{(\mathrm{CP})}(\tau) - \pi_1)/(1-\pi_1)$ for comparison with panel (a). The condition is applied *only at the starting time*; the trajectory is free ~~thereafter—holding it throughout would truncate the survival artificially~~ thereafter (Supplemental Material). The conditioned curve (Fig. 5b, dashed) sits between the unconditional empirical curve (solid) and the Markov reference (dotted), closing roughly 30% of the gap at a 200 min lag: more Markovian, but not fully.

Cold pools are part—but not all—of the hidden memory.—The unclosed portion of the gap implicates other slow ~~internal variables: mesoscale~~ variables—mesoscale moisture pooling [32], cloud-cluster geometry [33], ~~and~~ free-tropospheric ~~humidity all evolve on the relevant~~ humidity—evolving on similar time scales. Two caveats apply: CP(t) collapses spatial structure into a scalar, and stratifying by ~~CP(t)~~ it inherits some lifecycle-stage dependence; Supplemental Material sensitivity tests ~~(per-cold-pool diagnostic and entry-conditional survival anchored at state 1 entries)~~ are consistent with the closure reported here. A definitive identification ~~of the dominant carrier~~ would require re-training the encoder with explicit cold-pool inputs, ~~which we leave~~ left to future work.

DISCUSSION

Convection varies widely even under a fixed forcing, yet this variability is structured, not random. ~~We represented the flow with isentropic analysis and encoded~~ Casting the flow in isentropic coordinates and encoding it into a low-dimensional latent space, ~~where its evolution reduces~~ we reduced its evolution to a Markov chain over ~~a few discrete states . Five states proved enough to~~ five discrete states that recover physically meaningful ~~convective regimesand to follow the trajectories convection takes between them. This~~ regimes. The chain is nearly memoryless~~, with~~ : a single slow eigenmode ($\lambda_2 \approx 0.94$, $\tau_2 \approx 172$ min) ~~that alternates two regimes,~~ alternates organized and unorganized ~~,~~ regimes around a dominant suppressed background ($\pi_3 = 37\%$)~~: two competing routes emerge, a~~ , with two competing routes—a direct ascent into deep convection and a slower drift through weakly organized phases. ~~The framework's~~ Its distinctive feature is ~~that~~ diagnostic: departures from Markovianity ~~*diagnose* what the discrete partition omits: states 1 and 4 persist beyond their Markov prediction~~reveal what the partition omits, and conditioning on cold-pool intensity closes ∼30% of ~~that~~ the gap at a 200 min lag, identifying cold pools as one measurable carrier of the hidden memory.

These results describe a single ~~, idealized forcing—fixed vertical shear in radiative-convective equilibrium—and~~ idealized forcing, and the simplicity of the latent space may partly reflect that control~~: a more variable or transient forcing could populate a higher-dimensional space with more states and stronger non-Markovianity. Testing this is the natural next step.~~ . Because the transition matrix ~~is a function of~~ depends on the forcing, repeating the construction across shear regimes—weak (popcorn), moderate, strong (supercritical squall lines)~~—turns—~~would turn convective organization into a measurable dynamical observable~~: under stronger shear the~~ , with organized-state persistence T_{ii} and eigenmode time scales ~~should lengthen, and the direct route into deep convection should dominate. Sharper internal descriptors—cold-pool age, gust-front strength, mesoscale moisture, or cluster geometry—should each close more of the memory gap, every increment naming another slow variable a memory-aware parameterization must track~~lengthening under stronger shear.

DATA AND CODE AVAILABILITY

The idealized squall-line SAM simulations (27 days at 2-minute resolution, with 1D, 2D, and 3D outputs) are available via Globus (link to be provided). All code is openly available at `https://github.com/sososquirrel/troposwhere`, including the model implementation and the scripts to reproduce all figures. Full methods—simulation setup, isentropic preprocessing, model architecture, training, and sensitivity tests—are given in the Supplemental Material.

ACKNOWLEDGMENTS

We are grateful to Juan Nathanael and Carla Roesch for discussions on Koopman operators, and to Fabrizio Falasca and Matthieu Blanke for recommendations on the dynamical systems literature. We also thank Sara Shamekh, Zhe Feng, Xingchao Chen, and the WACCEM project team for valuable discussions throughout the course of this study. We thank Koichi Sakaguchi and Huancui Hu for their internal review and constructive feedback. Portions of the manuscript were rephrased with the assistance of ChatGPT (OpenAI) for language editing, and Claude (Anthropic) assisted with aspects of code development and cleaning; all scientific ideas, analysis, and conclusions were developed by the authors.

This study is supported by Office of Science, U.S. Department of Energy (DOE) Biological and Environmental Research as part of the Regional and Global Model Analysis program area through the Water Cycle: Modeling of Circulation, Convection, and Earth System Mechanisms (WACCEM) scientific focus area. S.A. designed




FIG. 5. **Markovianity assessment.** **(a)** Per-state indicator autocorrelations (solid) and ~~their~~ Markov predictions (dashed, same colour); states 1 and 4 sit ~~visibly~~ above their predictions, the others track theirs~~closely~~. **(b)** State-1 autocorrelation: empirical (solid~~blue~~), Markov prediction (dotted~~black~~), and empirical conditioned on $\mathrm{CP}(t) > \mathrm{CP}_\star$ at the starting time (dashed~~blue~~), ~~which closes~~ closing ∼30% of the ~~empirical–Markov~~ gap at a 200 min lag.

the research, performed the simulations, developed the methods and code, and wrote the manuscript. O.P. contributed to the methodological development and the code and participated in the analysis and discussions. L.R.L. and P.G. contributed to improving the analysis and interpretation.

[1] R. A. Houze, Jr., Rev. Geophys. **42**, RG4003 (2004).
[2] S. Chakraborty, S. C. Sullivan, and Z. Feng, Clouds and Their Climatic Impacts: Radiation, Circulation, and Precipitation , 195 (2023).
[3] R. Roca and T. Fiolleau, Communications Earth & Environment **1**, 1 (2020).
[4] Z. Feng, A. F. Prein, J. Kukulies, T. Fiolleau, W. K. Jones, B. Maybee, Z. Moon, K. M. N. Ocasio, W. Dong, M. J. Molina, *et al.*, Authorea Preprints (2024).
[5] Z. Feng, L. R. Leung, J. Hardin, C. R. Terai, F. Song, and P. Caldwell, Geophysical Research Letters **50**, e2022GL102603 (2023).
[6] Q. Yang, L. R. Leung, Z. Feng, and X. Chen, Journal of Climate **36**, 4597 (2023).
[7] M. Bolot, R. Roca, T. Fiolleau, and C. Muller, npj Climate and Atmospheric Science (2025).
[8] S. Abramian, C. Muller, C. Risi, T. Fiolleau, and R. Roca, npj climate and atmospheric science **8**, 258 (2025).
[9] Z. Feng, A. F. Prein, J. Kukulies, T. Fiolleau, W. K. Jones, B. Maybee, Z. L. Moon, K. M. Núñez Ocasio, W. Dong, M. J. Molina, *et al.*, Journal of Geophysical Research: Atmospheres **130**, e2024JD042204 (2025).
[10] H. Hu, L. R. Leung, and Z. Feng, Geophysical Research Letters **47**, e2019GL086783 (2020).
[11] T. Fiolleau and R. Roca, Earth System Science Data **16**, 4021 (2024).
[12] B. Khouider and A. J. Majda, Theoretical and Computational Fluid Dynamics **20**, 351 (2006).
[13] A. J. Majda and S. N. Stechmann, Proceedings of the National Academy of Sciences **105**, 17614 (2008).
[14] B. Khouider, J. Biello, and A. J. Majda, Commun. Math. Sci. **8**, 187 (2010).
[15] C. J. Muller and Y. Takayabu, Environ. Res. Lett. (2020).
[16] M. S. Singh and P. A. O'Gorman, J. Climate (2012).
[17] S. Abramian, C. Muller, and C. Risi, Journal of Advances in Modeling Earth Systems **15**, e2022MS003477 (2023).
[18] D. Wang, A. F. Prein, S. E. Giangrande, A. Ramos-Valle, M. Ge, and M. P. Jensen, Journal of Geophysical Research: Atmospheres **127**, e2022JD036746 (2022).
[19] Y.-X. Li, M. S. Singh, H. Masunaga, and J.-Y. Yu, Geophysical Research Letters **52**, e2025GL116056 (2025).
[20] B. O. Koopman, Proceedings of the National Academy of Sciences **17**, 315 (1931).
[21] S. L. Brunton, M. Budišić, E. Kaiser, and J. N. Kutz, arXiv preprint arXiv:2102.12086 (2021).
[22] B. Lusch, J. N. Kutz, and S. L. Brunton, Nature communications **9**, 4950 (2018).
[23] J. Nathaniel and P. Gentine, Computer Methods in Applied Mechanics and Engineering **448**, 118410 (2026).
[24] O. M. Pauluis and A. A. Mrowiec, Journal of the Atmospheric Sciences **70**, 3673 (2013).
[25] E. Kaiser, B. R. Noack, L. Cordier, A. Spohn, M. Segond, M. Abel, G. Daviller, J. Östh, S. Krajnović, and R. K. Niven, Journal of Fluid Mechanics **754**, 365 (2014).
[26] A. N. Souza, Journal of Fluid Mechanics **997**, A1 (2024).
[27] M. F. Khairoutdinov and D. A. Randall, J. Atmos. Sci. **60**, 607 (2003).
[28] G. J. Zhang, Journal of Geophysical Research: Atmospheres **108**, 10.1029/2003JD003520 (2003).
[29] P. Bechtold, N. Semane, P. Lopez, J.-P. Chaboureau, A. Beljaars, and N. Bormann, Journal of the Atmospheric Sciences **71**, 734 (2014).
[30] J.-M. Piriou, J.-L. Redelsperger, J.-F. Geleyn, J.-P. Lafore, and F. Guichard, Journal of the Atmospheric Sciences **64**, 4127 (2007).

[31] L. Gerard, J.-M. Piriou, R. Brožková, J.-F. Geleyn, and D. Banciu, Monthly Weather Review **137**, 3960 (2009).
[32] B. Mapes and R. Neale, J. Adv. Model. Earth Syst. **3** (2011).
[33] Y.-L. Hwong, M. Colin, P. Aglas-Leitner, C. J. Muller, and S. C. Sherwood, Journal of Advances in Modeling Earth Systems **15**, e2023MS003726 (2023).
[34] R. Rotunno, J. B. Klemp, and M. L. Weisman, J. Atmos. Sci. **45**, 463 (1988).
[35] R. A. Houze, Monthly Weather Review **105**, 1540 (1977).
[36] A. M. Tompkins, J. Atmos. Sci. **58**, 1650ï¿½1672 (2001).
[37] S. Abramian, C. Muller, and C. Risi, Geophysical Research Letters **49**, e2021GL095184 (2022).
[38] G. Torri, Z. Kuang, and Y. Tian, Geophys. Res. Lett. **42**, 1943 (2015).