

Causal Discovery from Complex Survey Data

A Covariate-Adjusted and Weighted Framework

To-Do List

- (this is not just a survey data problem)
- Image of a Directed Acyclic Graph representing causal discovery
- (maybe make that upstream/downstream comment about discovery and inference)
- Overall I am struggling with the fact that we are showing causal model for ordinal but no applied dataset for the other classes. Maybe this needs a little more thought.
- The Non-Identifiability Problem for Categorical Data (not sure if this really is background since Yang's paper already adjusts for this)

1 Introduction

Survey questionnaires are among the most widely used instruments in psychological, behavioral, and health services research. They provide population-level data on latent constructs such as depression, well-being, social functioning, health-related quality of life that cannot be measured through clinical observation or administrative records alone. Yet, despite their prevalence, survey data present a fundamental analytic challenge: understanding not merely which variables are associated, but which variables causally drive others **(this is not just a survey data problem)**. This distinction is not semantic. Knowing that financial hardship is correlated with poor mental health tells us little about whether interventions targeting financial need will improve mental health outcomes, whether the relationship operates through intermediate pathways, or whether the apparent association reflects a common upstream cause. Answering these questions requires causal knowledge, not associational knowledge.

Causal discovery methods — data-driven algorithms that infer directed causal structure from observational data — offer a powerful and increasingly practical approach to this challenge (Spirtes et al. (2001), Pearl (2009), Glymour et al. (2019)).

Image of a Directed Acyclic Graph representing causal discovery

Unlike structural equation modeling (SEM), which requires investigators to fully specify the causal structure before fitting a model, causal discovery algorithms learn the structure objectively from data, subject to identifiability assumptions (Zanga et al. (2022)) **(maybe make that upstream/downstream comment about discovery and inference)**. This property makes them especially valuable in domains where causal relationships are plausible but their precise ordering is unknown — a situation that describes the vast majority of psychological and social science research. Applications of causal discovery in psychology have grown substantially in recent years, spanning psychopathology (Briganti et al. (2024), Zhu et al. (2026)), academic achievement (Quintana (2023)), eating disorders (Anderson et al. (2023)), and most recently, obsessive-compulsive disorder and depression (Ni et al. (2025)).

Despite this growth, a critical assumption underlies every existing causal discovery algorithm: that observations are independent and identically distributed (i.i.d.). In practice, this assumption is routinely violated by the very data sources most commonly used in psychological and social research. Large-scale survey datasets — the National Health Interview Survey, the Behavioral Risk Factor Surveillance System, the Consumer Assessment of Healthcare Providers and Systems (CAHPS), the Health Behavior in School-Aged Children study — are collected using complex probability sampling designs that deliberately oversample certain demographic or clinical subgroups to ensure adequate representation of minority populations and clinical subpopulations (Groves et al. (2009)).

[Image of complex probability survey sampling design]

These designs assign unequal selection probabilities to respondents, and the resulting data are not i.i.d. When survey weights, which encode the inverse of each respondent’s selection probability, are ignored, the observed joint distribution of variables in the sample does not represent the joint distribution in the target population. This discrepancy is not merely a precision problem — it is a systematic bias that distorts conditional independence relationships, the very relationships that causal discovery algorithms rely upon to identify causal structure.

To understand why this matters concretely, consider a health survey that oversamples Veterans with both high basic needs burden and poor health outcomes. In such a sample, the correlation between basic needs and health outcomes will be inflated relative to the true population relationship, because the joint distribution has been distorted by the sampling design. An unweighted causal discovery algorithm, presented with this distorted distribution, will infer spurious direct edges between variables that are in fact conditionally independent in the population once sampling design is accounted for. The result is a causal graph that reflects the sampling mechanism rather than the true data-generating process. Sampling-induced bias of this kind has long been recognized in the causal inference literature focused on treatment effect estimation (Zanutto et al. (2005), Wang et al. (2009), Benedetti et al. (2024), Vo et al. (2024)), and recent work on causal mediation analysis confirms that ignoring survey weights in even moderately complex designs produces substantial bias in estimated causal effects (Coffman et al. (2024)). Yet to our knowledge, no prior work has addressed survey weighting in the context of causal structure learning — that is, in recovering the causal graph itself rather than estimating effects along a pre-specified graph.

This omission is particularly consequential for ordinal survey data. The dominant class of causal discovery algorithms for categorical data relies on standard multinomial Bayesian networks (BNs), which are well-known to be non-identifiable: multiple causal graphs can fit the observed data equally well, leaving the direction of many causal relationships undetermined (Spirtes et al. (2001), Chickering (2002)). This non-identifiability problem is separate from and compounds the survey weighting problem — even after correcting for sampling design, an analyst using a standard categorical BN cannot uniquely determine causal direction from observational ordinal data. Ni and Mallick (2022) resolved this identifiability problem for ordinal data by showing that ordinal Bayesian networks (oBNs), which model conditional distributions using cumulative ordinal regression, are fully identifiable under mild regularity conditions: the ordinal structure of the data breaks the distributional equivalence that renders nominal categorical BNs non-identifiable. Ni et al. (2025) subsequently developed a bootstrapped oBN algorithm for survey questionnaire data and demonstrated its utility in a psychological study of obsessive-compulsive disorder and depression. However, neither of these contributions addresses the survey weighting problem: the algorithms assume i.i.d. data and do not account for the complex sampling designs ubiquitous in the survey data they are intended to analyze.

The present paper addresses this gap. We develop survey-weighted causal discovery methods spanning three major algorithm families and introduce a unified framework, the *survey-weighted*

ordinal Bayesian network (swa-oBN), that simultaneously addresses both the non-identifiability of standard categorical BNs and the distributional bias induced by complex survey sampling [(okay but needs reframing - we do discuss survey weighting importance, which is fine,)]. Specifically, our contributions are as follows.

First, for score-based algorithms, we incorporate survey weights into the BIC scoring function through a pseudo-likelihood approach combined with Kish's (Kish (1965)) effective sample size, yielding a survey-weighted pseudo-BIC that adjusts both the likelihood and the model complexity penalty for the sampling design. Applied to ordinal data via cumulative ordinal regression, this produces the swa-oBN , which is both fully identifiable and design-consistent.

Second, for constraint-based algorithms, including the PC algorithm and the Fast Causal Inference (FCI) algorithm, we develop a survey-weighted conditional independence test based on design-adjusted regression via the R `survey` package (Lumley (2004)), which replaces standard Gaussian or chi-squared tests with Wald tests that account for the complex sampling design.

Third, for asymmetry-based algorithms, specifically `LINGAM` (Shimizu et al. (2006)), we develop a weighted pseudo-population resampling procedure that restores i.i.d. structure by resampling from the observed data proportional to survey weights, recovering the target population distribution prior to causal discovery.

Fourth, we demonstrate through simulation studies that ignoring survey weights in each of these algorithm families produces systematically biased causal graphs, and that the proposed survey-weighted variants recover the true causal structure.

Fifth, we illustrate the practical utility of the swa-oBN in an application to data from the Consumer Assessment of Healthcare Providers and Systems (CAHPS) Clinician and Group Visit Survey administered to United States Veterans Affairs (VA) patients. The CAHPS survey, conducted using a stratified probability sample with substantial variation in selection probabilities across demographic and service utilization subgroups, provides ordinal measures of social determinants of health, healthcare experience quality, and health outcomes. Applying swa-oBN with bootstrapped uncertainty quantification, we uncover the causal structure among clusters of social needs, finding that [fill in later](#).

Overall I am struggling with the fact that we are showing causal model for ordinal but no applied dataset for the other classes. Maybe this needs a little more thought.

The remainder of the paper is organized as follows. [fill in later](#)

2 Background

2.1 Causal Graphs and Bayesian Networks

A directed acyclic graph (DAG) $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ consists of a set of nodes $\mathcal{V} = \{X_1, \dots, X_q\}$ representing random variables and a set of directed edges $\mathcal{E} \subset \mathcal{V} \times \mathcal{V}$ representing direct causal relationships, with no directed cycles. For a node X_j , let $\text{pa}(j) = \{k \in \mathcal{V} : k \rightarrow j\}$ denote the set of its parents, i.e., its direct causes. A Bayesian network (BN) is a pair $\mathcal{B} = (\mathcal{G}, P)$ in which the probability distribution P factorizes with respect to $\mathcal{G}$ as

$$P(X_1, \dots, X_q \mid \mathcal{G}) = \prod_{j=1}^q P(X_j \mid X_{\text{pa}(j)}).$$

When $\mathcal{G}$ is interpreted causally, that is, when directed edges are understood as causal relationships rather than mere statistical dependencies, the factorization above encodes the *Causal Markov Assumption*: each variable is conditionally independent of its non-descendants given its direct causes (Spirtes et al. (2001), Pearl (2009), Glymour et al. (2019)). Together with the *Faithfulness Assumption* (Glymour et al. (2019)), which requires that all conditional independence relationships in the data are entailed by the graph structure and are not the product of parameter cancellation, the Causal Markov Assumption forms the theoretical bedrock of causal discovery (Uhler et al. (2013); Zhalama et al. (2017)). A third standard assumption, *Causal Sufficiency*, requires that all common causes of any two variables in the model are observed and included in $\mathcal{V}$. When Causal Sufficiency is relaxed, algorithms such as FCI output Partial Ancestral Graphs (PAGs) rather than DAGs, encoding uncertainty about the presence of latent confounders (Pearl (2009)).

Three broad families of algorithms have been developed for learning causal structure from observational data. Constraint-based algorithms, including the PC algorithm (Spirtes and Glymour (1991)) and its extension FCI (Spirtes et al. (1995)), test for conditional independence among pairs of variables and use detected independence patterns — particularly v-structures of the form $X_i \rightarrow X_k \leftarrow X_j$ — to orient edges. Score-based algorithms, including the Greedy Equivalence Search (GES; Chickering (2002)) and the ordinal Bayesian network (Ni and Mallick (2022)), assign a goodness-of-fit score to candidate graphs and search for the highest-scoring structure, typically using the Bayesian information criterion (BIC). Asymmetry-based algorithms, most prominently LiNGAM (Shimizu et al. (2006)), exploit distributional asymmetries between cause and effect — specifically, the independence between a cause and the residual noise of its effect — to identify causal direction beyond the Markov equivalence class. Each family rests on the i.i.d. assumption as a foundational requirement for its statistical guarantees.

2.2 Why Survey Weighting Matters for Causal Discovery

The distortion introduced by complex survey sampling is not merely a variance problem — it is a systematic bias problem for causal structure learning. To see this, consider the following stylized example. Suppose the true population causal structure is $X \rightarrow Y \rightarrow Z$, where X represents basic financial need, Y represents access to healthcare, and Z represents well-being. In the population, X and Z are conditionally independent given Y : once healthcare access is known, financial need carries no additional information about well-being. Now suppose the survey over-samples units where both X is high and Z is low — a common pattern in health surveys that deliberately capture high-need, low-outcome populations. In the resulting sample, X and Z are no longer conditionally independent given Y : the sampling mechanism has introduced a spurious association between the upstream cause and the downstream outcome. An unweighted causal discovery algorithm will observe this spurious conditional dependence, fail to remove the edge $X - Z$, and conclude either that there is a direct causal effect of X on Z , or that Z causes X . Both conclusions are wrong.

This mechanism generalizes across algorithm families. For constraint-based algorithms (PC, FCI), the conditional independence tests that drive skeleton discovery and edge orientation are corrupted by the sampling-distorted conditional distributions. Tests that should return $p > \alpha$ (indicating independence) return $p < \alpha$ because the sampling design has inflated the apparent association. For score-based algorithms (oBN, GES), the likelihood function is computed over the sample distribution rather than the population distribution, causing the BIC to favor denser graphs with spurious edges. For asymmetry-based algorithms (LiNGAM), the non-Gaussian distributional properties that identify causal direction are distorted by the sampling mechanism, potentially reversing the apparent causal direction. In all three cases, the result is not merely imprecision — it is systematic structural misidentification that does not diminish with increasing sample size, because the bias is determined by the sampling design, not the sample size.

This problem is distinct from, but related to, the well-studied problem of selection bias in causal inference. Selection bias in the potential outcomes framework refers to self-selection into treatment; the survey sampling problem is a design-induced form of selection that affects the joint distribution of all variables, not merely the assignment mechanism for a specific treatment. The appropriate correction — inverse probability weighting of the likelihood function — is analogous in structure but applies at the level of the causal structure learning algorithm itself rather than at the level of treatment effect estimation. To our knowledge, no prior causal discovery work has implemented this correction. The closest work in the causal inference literature addresses the use of survey weights in treatment effect estimation (Zanutto (2006); Liang and Wu (2026)) and causal mediation analysis (Coffman et al. (2024)), not causal structure learning.

2.3 Survey Sampling and the i.i.d. Assumption

Large-scale surveys in psychology, public health, and social science rarely use simple random sampling. Instead, they employ complex probability sampling designs — typically involving stratification, clustering, and deliberate oversampling of certain subpopulations — to achieve representation of minority groups and to control survey costs (Lumley (2004), Groves et al. (2009)). Under such designs, each respondent i is selected with a known, non-uniform probability π_i , and is assigned a survey weight $w_i = 1/\pi_i$ that represents the number of population members for whom respondent i serves as a proxy. The VA CAHPS survey, for example, oversamples Veterans belonging to racial and ethnic minority groups, Veterans in rural areas, and Veterans with high healthcare utilization, producing weights that can vary by an order of magnitude across respondents.

The consequence of unequal selection probabilities is that the observed sample distribution is not equal to the target population distribution. Formally, if $P_{\text{pop}}(X)$ denotes the true population joint distribution and $P_{\text{samp}}(X)$ the distribution of the selected sample, then

$$P_{\text{samp}}(X = x) \propto \pi(x) \cdot P_{\text{pop}}(X = x),$$

where $\pi(x)$ is the selection probability for a unit with covariate pattern x . When $\pi(x)$ is correlated with the outcome variables of interest — as is almost always the case in health surveys, where high-need, high-utilization subgroups are oversampled — P_{samp} differs systematically from P_{pop} . In particular, marginal distributions, pairwise correlations, and conditional independence relationships may all differ between the sample and the population. Causal discovery algorithms applied to the unweighted sample data therefore operate on a distorted version of the joint distribution, and the resulting causal graphs reflect the sampling mechanism rather than the true population causal structure.

To recover population-representative estimates from a complex survey, analysts apply the Horvitz-Thompson (HT) estimator (Horvitz and Thompson (1952)), weighting each observation by $w_i = 1/\pi_i$ to correct for unequal selection probabilities. For regression models, the survey-weighted pseudo-likelihood replaces the ordinary likelihood with

$$\tilde{\ell}(\theta) = \sum_{i=1}^n w_i \log p(x_i; \theta),$$

which is an asymptotically unbiased estimator of the population log-likelihood $N \cdot E_{\text{pop}}[\log p(X; \theta)]$ under standard regularity conditions (Lumley (2004)). Maximizing the pseudo-likelihood thus recovers asymptotically consistent estimates of the population parameters, rather than sample-specific parameters. The penalty term of the BIC must also be adjusted under survey weighting, because the relevant sample size for model selection is not the raw count n but the effective

sample size n_{eff} , which accounts for the precision loss induced by unequal weights. Kish (Kish (1965)) proposed the widely used estimator

$$n_{\text{eff}} = \frac{(\sum_{i=1}^n w_i)^2}{\sum_{i=1}^n w_i^2},$$

which reduces to n when all weights are equal. When weights are highly variable — as in surveys with substantial oversampling — n_{eff} can be substantially smaller than n , reflecting the reduced statistical information available for inference. Substituting n_{eff} for n in the BIC penalty produces the survey-adjusted pseudo-BIC, which appropriately penalizes model complexity relative to the effective rather than nominal sample size.

2.4 The Non-Identifiability Problem for Categorical Data (not sure if this really is background since Yang’s paper already adjusts for this)

A central challenge for causal discovery with categorical data is the non-identifiability of standard BN models. Two DAGs are Markov equivalent if they encode exactly the same set of conditional independence relationships. Markov equivalent DAGs cannot be distinguished from one another on the basis of observational data alone, as they imply identical joint distributions. For continuous data under Gaussian or non-Gaussian additive noise models, Markov equivalence classes are often small or can be broken entirely by exploiting distributional properties (Bühlmann et al. (2014); Shimizu et al. (2006)). For categorical data modeled with multinomial distributions, however, Markov equivalence is a far more severe problem: as established by Chickering (2002), multinomial BNs are score-equivalent in the sense that Markov equivalent DAGs achieve identical maximum likelihood and hence identical BIC scores. This means that for any fitted causal model $X \rightarrow Y$ under a multinomial BN, there exists a reverse model $Y \rightarrow X$ that fits the data equally well. The direction of many — potentially all — causal relationships remains undetermined.

Ni and Mallick (2022) resolved this non-identifiability for ordinal categorical data by proposing the ordinal Bayesian network (oBN). Rather than modeling conditional distributions with unrestricted multinomial tables, the oBN models each conditional distribution $P(X_j \mid X_{\text{pa}(j)})$ using cumulative ordinal regression:

$$P(X_j \leq \ell \mid X_{\text{pa}(j)}) = F\left(\gamma_{j\ell} - \sum_{k \in \text{pa}(j)} \beta_{jk} X_k - \alpha_j\right), \quad \ell = 1, \dots, L_j,$$

where F is a link function (e.g., probit), $\gamma_{j\ell}$ are ordered threshold parameters, β_{jk} are regression coefficients encoding the directional causal effect of parent X_k on child X_j , and α_j is an intercept. The key theoretical result is that for ordinal variables with $L > 2$ levels, the model $X \rightarrow Y$ and

its reverse $Y \rightarrow X$ are generically not distribution-equivalent: the ordinal regression structure of one direction cannot be mimicked by the reverse direction for almost all parameter values. Ni and Mallick (2022) proved this formally for the bivariate case, and Ni et al. (2025) provided asymptotic consistency guarantees for multivariate data by showing that the BIC of the true generating DAG converges to $-\infty$ relative to any Markov-equivalent alternative, making the correct causal structure uniquely identifiable in the large-sample limit. This identifiability relies on the ordinal structure: variables with only two levels (binary variables) do not benefit from this result, as ordinal and nominal binary distributions coincide. The `OrdCD` R package (Ni (2023)) implements this algorithm with hill-climbing search, multiple random restarts, and bootstrapping for uncertainty quantification.

2.5 The CAHPS Survey and VA Population

The Consumer Assessment of Healthcare Providers and Systems (CAHPS) Clinician and Group Visit Survey is a standardized, validated instrument for measuring patients’ experiences with healthcare providers and the social circumstances that influence health (Agency for Healthcare Research and Quality (2023)). The version administered by the Veterans Health Administration (VHA) — the largest integrated healthcare system in the United States, serving over nine million enrolled Veterans — includes measures of healthcare visit experience quality (communication, respect, listening, timeliness), social determinants of health across fifteen domains (financial need, housing, food security, transportation, caregiving, employment, legal need, social isolation, discrimination, and others), and health outcomes (overall health, mental/emotional health, and functional well-being). The social determinants section employs a three-category ordinal response format (“No support needed,” “Needed support and got it,” “Needed support but did not get it”) for each domain, capturing both the presence of need and whether that need was met — a substantively richer measure than simple need prevalence.

Veterans enrolled in VHA care represent a population with substantially elevated social and medical complexity. Compared to non-Veterans and to Veterans not engaged in VHA services, VHA-enrolled Veterans have lower physical and mental health functioning, higher rates of PTSD and depression, greater food insecurity and housing instability, and lower socioeconomic resources (Duan-Porter et al. (2018); Washington (2022)). Social determinants of health have been identified as significant predictors of Veteran health outcomes, but their causal ordering — which needs drive which outcomes, and through what pathways — remains poorly understood (Duan-Porter et al. (2018)). Existing studies are almost exclusively associational, relying on regression models that assume, rather than discover, causal structure (Montgomery et al. (2020)). The CAHPS data are collected using a stratified probability sample with substantial variation in selection probabilities across demographic groups, making them a canonical case where survey weighting is essential for valid population inference, and where its importance for causal structure learning has never been examined.

3 Methods

This section presents survey-weighted causal discovery methods for three algorithm families: score-based (Section 3.1), constraint-based (Section 3.2), and asymmetry-based (Section 3.3). Section 3.4 presents the bootstrapped swa-oBN algorithm with confounder adjustment, which constitutes the primary contribution for ordinal survey data. Throughout, let $\mathbf{y} = (y_1, \dots, y_n)$ denote the observed sample of n respondents, $\mathbf{w} = (w_1, \dots, w_n)$ their associated survey weights, and $\mathbf{X} = (X_1, \dots, X_q)$ the random vector of q variables under study.

3.1 Survey-Weighted Score-Based Causal Discovery: The swa-oBN

3.1.1 Survey-Weighted Pseudo-BIC

Score-based causal discovery algorithms evaluate candidate DAGs by computing a goodness-of-fit score. The standard BIC for a DAG $\mathcal{G}$ takes the form

$$\text{BIC}(\mathcal{G} \mid \mathbf{y}) = -2 \sum_{i=1}^n \log \hat{p}(y_i \mid \mathcal{G}) + K \log(n),$$

where $\hat{p}(y_i \mid \mathcal{G})$ is the joint likelihood evaluated at the maximum likelihood estimates, K is the number of free model parameters, and n is the sample size. Under simple random sampling, maximizing this BIC recovers the population-level causal structure consistently. Under complex survey sampling, however, the ordinary likelihood $\sum_{i=1}^n \log \hat{p}(y_i \mid \mathcal{G})$ targets the sample distribution rather than the population distribution, producing biased parameter estimates and a biased model selection criterion.

We replace the ordinary likelihood with the survey-weighted pseudo-likelihood (Binder, 1983; Lumley, 2010):

$$\tilde{\ell}(\theta \mid \mathcal{G}) = \sum_{i=1}^n w_i \log p(y_i \mid \mathcal{G}, \theta).$$

Under standard regularity conditions, $\tilde{\ell}(\theta \mid \mathcal{G})$ is an asymptotically unbiased estimator of $N \cdot E_{\text{pop}}[\log p(Y \mid \mathcal{G}, \theta)]$, where N is the population size and E_{pop} denotes expectation under the true population distribution. Maximizing $\tilde{\ell}(\theta \mid \mathcal{G})$ with respect to θ therefore recovers the population-consistent pseudo-maximum likelihood estimate $\hat{\theta}^w$.

The penalty term of the BIC must also be modified. The BIC penalty $K \log(n)$ is calibrated to the amount of statistical information in the data, which under complex survey sampling is not n but the effective sample size n_{eff} . Substituting Kish's (1965) effective sample size formula

$$n_{\text{eff}} = \frac{(\sum_{i=1}^n w_i)^2}{\sum_{i=1}^n w_i^2}$$

yields the survey-weighted pseudo-BIC:

$$\widetilde{\text{BIC}}(\mathcal{G} \mid \mathbf{y}, \mathbf{w}) = -2\tilde{\ell}(\hat{\theta}^w \mid \mathcal{G}) + K \log(n_{\text{eff}}).$$

When all weights are equal (simple random sample), $n_{\text{eff}} = n$ and the pseudo-BIC reduces exactly to the standard BIC. When weights are highly variable — as in surveys with substantial oversampling — $n_{\text{eff}} \ll n$, and the pseudo-BIC applies a heavier penalty to complex models, appropriately reflecting the reduced information available in the weighted sample. Lumley and Scott (2015) established AIC and BIC for survey-weighted models on formal information-theoretic grounds, validating this approach for model selection from complex survey data.

3.1.2 Ordinal Regression Scorer

For ordinal variables $X_j \in \{1, \dots, L_j\}$, the node-specific score for node j given parent set $\text{pa}(j)$ is computed via cumulative probit ordinal regression (Ni and Mallick, 2022):

$$P(X_j \leq \ell \mid X_{\text{pa}(j)}) = \Phi\left(\gamma_{j\ell} - \sum_{k \in \text{pa}(j)} \beta_{jk} X_k - \alpha_j\right), \quad \ell = 1, \dots, L_j,$$

where Φ is the standard normal CDF, $\gamma_{j\ell}$ are ordered threshold parameters with $\gamma_{j1} = 0$ fixed for identifiability, β_{jk} are causal effect coefficients, and α_j is an intercept. For binary nodes ($L_j = 2$), the model reduces to a weighted probit regression fitted via the `svyglm` function with `family = quasibinomial(link = "probit")` in R's survey package (Lumley, 2004). For ordinal nodes with $L_j > 2$ levels, the model is fitted via `svyolr` in the same package, which implements Binder's (1983) design-based variance estimator for survey-weighted proportional odds models. The node-specific pseudo-BIC contribution is then

$$\widetilde{\text{BIC}}_j(\text{pa}(j) \mid \mathbf{y}, \mathbf{w}) = -2\tilde{\ell}_j(\hat{\theta}_j^w) + K_j \log(n_{\text{eff}}),$$

where $\tilde{\ell}_j$ is the pseudo-likelihood of node j given its parents and $K_j = (L_j - 1) + \sum_{k \in \text{pa}(j)} (L_k - 1)$ counts the free parameters for this node. The total graph score decomposes as $\widetilde{\text{BIC}}(\mathcal{G}) = \sum_{j=1}^q \widetilde{\text{BIC}}_j(\text{pa}(j))$, following the standard local score decomposition of BN structure learning (Chickering, 2002).

3.1.3 Hill-Climbing Search and the swa-oBN Algorithm

Structure search is performed by a greedy hill-climbing algorithm (Algorithm 1) that iterates over three move types — edge addition, edge deletion, and edge reversal — accepting any move that strictly decreases $\widetilde{\text{BIC}}(\mathcal{G})$ and terminates when no improving move exists. Acyclicity is checked at each candidate move using the `is.DAG` function from the `gRbase` package (Højsgaard et al., 2012). To mitigate sensitivity to local optima, the algorithm is run from R random initial DAGs and the solution with the minimum total $\widetilde{\text{BIC}}$ is retained (Ni et al., 2025). A blacklist matrix $\mathbf{B}$ encodes domain knowledge constraints: if $\mathbf{B}_{ij} = 1$, the edge $X_j \rightarrow X_i$ is forbidden and no move introducing it is considered. In the CAHPS application, outcomes (Well-being and Mental Health) are blacklisted from having any outgoing edges, encoding the substantive constraint that these are terminal nodes in the causal system.

We refer to the resulting algorithm as the **survey-weighted asymptotically identified ordinal Bayesian network (swa-oBN)**. The large-sample properties established by Ni et al. (2025) extend directly to the swa-oBN: since the pseudo-likelihood is an asymptotically unbiased estimator of the population log-likelihood, the Kullback-Leibler divergence argument in Ni et al. (2025, Section 3.3) holds with $\tilde{\ell}$ replacing ℓ , ensuring that the true generating DAG achieves a strictly lower pseudo-BIC than any Markov-equivalent alternative almost surely as $n_{\text{eff}} \rightarrow \infty$.

Algorithm 1: swa-oBN Hill-Climbing

```

Input: data y, weights w, n_eff, link F, ic criterion,
      blacklist B, max iterations T, random restarts R
Output: estimated DAG G*

best_ic ← ∞; G* ←
for r = 1, ..., R do
  Initialize G from empty DAG (r=1) or random DAG
  Enforce blacklist: G ← G ⊖ B
  Compute node scores: s_j ← BIC_j(pa_G(j)) for all j
  repeat
    Improvement ← FALSE
    for each candidate move m ∈ {add, delete, reverse} do
      if m is acyclic and respects B then
        Compute Δ ← change in s_j (and s_k for reversals)
        if Δ < 0 then accept m, update s_j, set Improvement ← TRUE
    until Improvement = FALSE or iter > T
  if BIC(G) < best_ic then G* ← G; best_ic ← BIC(G)
return G*
```

3.1.4 Bootstrapped Uncertainty Quantification

A single point estimate of a causal DAG is subject to sampling variability. Following Ni et al. (2025) and Briganti et al. (2023), we quantify uncertainty through bootstrapping. For $b = 1, \dots, B$, we draw a bootstrap resample $\mathbf{y}^{(b)}$ of size n with replacement from the observed data, preserving the associated weights $\mathbf{w}^{(b)}$, and apply Algorithm 1 to each bootstrap sample. The edge inclusion probability matrix $\hat{\mathbf{P}} = [\hat{P}_{jk}]$ is computed as

$$\hat{P}_{jk} = \frac{1}{B} \sum_{b=1}^B \mathbb{1} [X_k \rightarrow X_j \in \hat{\mathcal{G}}^{(b)}],$$

where $\mathbb{1}[\cdot]$ is the indicator function. An entry $\hat{P}_{jk}$ close to 1 indicates that the directed edge $X_k \rightarrow X_j$ is consistently recovered across bootstrap resamples; an entry close to 0 indicates consistent absence; an entry near 0.5 indicates high uncertainty. Following Briganti et al. (2023), edges with $\hat{P}_{jk} > 0.85$ are retained in the final consensus graph as highly stable, while edges with $0.50 < \hat{P}_{jk} \leq 0.85$ are reported as present but less certain.

3.1.5 3.1.5 Confounder Adjustment

Survey questionnaire data frequently include demographic variables — age, race/ethnicity, sex, education — that are potential confounders of the causal relationships among substantive variables but are not themselves targets of causal discovery. The swa-oBN accommodates this through a confounder matrix $\mathbf{Z}$: a set of variables included as covariates in every node-level ordinal regression but excluded from the DAG structure entirely. No edges to or from variables in $\mathbf{Z}$ are learned. Formally, the node-specific regression for node j with parents $\text{pa}(j)$ becomes

$$P(X_j \leq \ell \mid X_{\text{pa}(j)}, \mathbf{Z}) = \Phi \left(\gamma_{j\ell} - \sum_{k \in \text{pa}(j)} \beta_{jk} X_k - \sum_m \delta_{jm} Z_m - \alpha_j \right),$$

where δ_{jm} are confounder regression coefficients estimated but not reported as causal effects. This ensures that the recovered edges among $\mathbf{X}$ variables represent causal relationships net of confounding by measured demographic characteristics, without distorting the causal graph by treating demographics as nodes.

3.2 3.2 Survey-Weighted Constraint-Based Causal Discovery

3.2.1 3.2.1 The Conditional Independence Test Under Survey Weighting

Constraint-based algorithms — including PC (Spirtes et al., 2001) and FCI (Spirtes et al., 1995) — rely on a sequence of conditional independence tests. Specifically, they test the null hypothesis $H_0 : X_i \perp\!\!\!\perp X_j \mid \mathbf{S}$ for variable pairs (i, j) conditional on a set $\mathbf{S} \subseteq \mathcal{V} \setminus \{i, j\}$. For continuous data under simple random sampling, these tests are typically implemented via Fisher’s Z-transformation of partial correlations. For complex survey data, the standard partial correlation test is invalid because it fails to account for the non-i.i.d. structure of the observations.

We propose replacing the standard conditional independence test with a design-adjusted Wald test based on survey-weighted regression. Specifically, to test $X_i \perp\!\!\!\perp X_j \mid \mathbf{S}$, we regress X_i on X_j and $\mathbf{S}$ using a survey-weighted generalized linear model via `svyglm` in R’s survey package (Lumley, 2004):

$$X_i \sim X_j + X_{S_1} + \cdots + X_{S_{|\mathbf{S}|}},$$

with survey design specified via `svydesign(ids = ~1, weights = ~w)`. For continuous variables, this is a weighted linear regression; for binary variables, a `quasibinomial` link is used. The p -value returned is the design-adjusted Wald test p -value for the coefficient of X_j , which incorporates the survey weights and accounts for the complex sampling design in its standard error computation (Lumley and Scott, 2014). The null of conditional independence is rejected when this p -value falls below the significance threshold α .

This test is design-consistent: under the null $X_i \perp\!\!\!\perp X_j \mid \mathbf{S}$ in the population, the Wald test p -value converges to a uniform distribution on $[0, 1]$ as $n_{\text{eff}} \rightarrow \infty$, ensuring the correct Type I error rate. Under the alternative, the test is consistent because the population-level conditional dependence produces a non-zero regression coefficient for X_j regardless of the survey design, and the weighted regression recovers this coefficient consistently.

3.2.2 3.2.2 Survey-Weighted PC and FCI Algorithms

The survey-weighted conditional independence test described above can be plugged directly into the standard `pc` and `fci` functions from the `pcalg` R package (Kalisch et al., 2012) via the `indepTest` argument, which accepts a user-specified function of the form `function(x, y, S, suffStat)`. The sufficient statistic `suffStat` passes the raw data matrix and the weight vector to the test function. No other modification to the PC or FCI algorithm skeleton is required — the survey weighting is fully encapsulated within the conditional independence test. The resulting algorithms, which we call **swa-PC** and **swa-FCI**, output a CPDAG and a PAG respectively, with

the same interpretation as their unweighted counterparts but with conditional independencies now estimated from the population-representative weighted distribution rather than the biased sample distribution.

An important consideration is that the swa-PC and swa-FCI algorithms inherit the non-identifiability of standard PC and FCI for ordinal categorical data: they can identify at most the Markov equivalence class of the true DAG, leaving the direction of some edges undetermined. For applications where full causal direction is required — as in the CAHPS analysis — the swa-oBN is preferred. However, swa-PC and swa-FCI retain value when the goal is skeleton discovery (identifying which variables are causally adjacent) rather than full edge orientation, when the sufficiency assumption is plausible but not certain (favoring FCI over swa-oBN), or when continuous variables are present in the data.

3.2.3 A Hybrid Approach: swa-PC + swa-oBN

For ordinal data, the two-step hybrid algorithm of Ni et al. (2025) generalizes naturally to the survey-weighted setting. In the first step, swa-PC is applied to determine the skeleton and orient v-structures, producing a CPDAG with some directed and some undirected edges. In the second step, for each pair of nodes connected by an undirected edge in the CPDAG, the bivariate swa-oBN is applied using exhaustive search over the three possible bivariate DAGs (null, $X_i \rightarrow X_j$, $X_j \rightarrow X_i$) to determine the edge orientation. This hybrid approach, which we call **swa-PC+oBN**, combines the computational efficiency of PC for skeleton discovery with the identifiability of oBN for edge orientation, and benefits from survey weighting at both stages.

3.3 Survey-Weighted Asymmetry-Based Causal Discovery: swa-LiNGAM

3.3.1 The LiNGAM Identifiability Mechanism and Survey Weighting

LiNGAM (Shimizu et al., 2006) assumes a linear structural equation model in which each variable is a linear function of its causes plus an independent, non-Gaussian noise term:

$$X_j = \sum_{k \in \text{pa}(j)} b_{jk} X_k + \varepsilon_j, \quad \varepsilon_j \sim F_j,$$

where F_j is a non-Gaussian distribution and all ε_j are mutually independent. The identifiability of LiNGAM rests on the asymmetry between the cause and effect under non-Gaussian noise: in the true causal direction, the cause X_k and the residual $\hat{\varepsilon}_j = X_j - \hat{b}_{jk} X_k$ are independent; in the reverse direction, the residual is dependent on the putative cause. This independence-based asymmetry is estimated via independent component analysis (ICA; Hyvärinen et al., 2001).

Complex survey sampling distorts this identifiability mechanism because the non-Gaussian distributional signature that ICA uses to distinguish causal direction is a population-level property, not a sample-level property. When the sample distribution differs from the population distribution — as it does under stratified oversampling — the residual independence properties are computed on the wrong distribution, potentially reversing the apparent causal direction.

3.3.2 3.3.2 Pseudo-Population Resampling

We address this via a **weighted pseudo-population bootstrap** (Efron, 1992; Rao and Wu, 1988). Rather than applying LiNGAM directly to the biased sample, we first reconstruct an approximately i.i.d. pseudo-population by resampling from the observed data with replacement according to probabilities proportional to the survey weights:

$$\text{prob}_i = \frac{w_i}{\sum_{j=1}^n w_j}, \quad i = 1, \dots, n.$$

A pseudo-population resample $\mathbf{y}^{(b)}$ of size n drawn with these probabilities has the property that its empirical distribution approximates the population distribution P_{pop} , because units with higher weights — representing larger portions of the population — are drawn with proportionally higher probability. Standard LiNGAM is then applied to each pseudo-population resample. This procedure is repeated B times, and edges are retained by majority vote: an edge $X_k \rightarrow X_j$ is included in the consensus graph if it appears in more than half of the B LiNGAM solutions. We call this procedure **swa-LiNGAM**.

The weighted pseudo-population bootstrap has an established theoretical basis: Rao and Wu (1988) showed that resampling with probability proportional to weights consistently estimates the sampling distribution of weighted estimators, and the approach has been widely used for variance estimation in complex surveys (Efron, 1992; Rao et al., 1992). In the LiNGAM context, the key property is that each pseudo-population resample is approximately i.i.d. from P_{pop} , restoring the distributional conditions under which LiNGAM’s ICA-based identification is valid.

A practical consideration is that swa-LiNGAM requires the substantive variables to be continuously measured and non-Gaussian — conditions met by many health and behavioral outcomes but not by ordinal survey response items. For ordinal data from the CAHPS instrument, the swa-oBN is the appropriate primary method; swa-LiNGAM is presented for completeness and for settings where continuous non-Gaussian variables are present in the data.

3.4 3.4 Choosing Among Methods and the swa-oBN as the Primary Recommendation

The three survey-weighted algorithms address complementary settings, and the choice among them should be guided by data type and the strength of available assumptions. Table 1 summarizes the key features.

Table 1: Survey-Weighted Causal Discovery Methods — Comparison

Method	Data type	Identifiable?	Confounders allowed?	Blacklist support	Primary use
swa-oBN	Ordinal (≥ 3 levels)	Yes (fully)	Via $\mathbf{Z}$ matrix	Yes	Primary for ordinal survey data
swa-PC	Mixed/continuous	No (CPDAG only)	Via FCI extension	Yes	Skeleton discovery, mixed data
swa-FCI	Mixed/continuous	No (PAG only)	Yes (latent confounders)	Yes	When unobserved confounders plausible
swa-PC+oBN	Ordinal	Yes (for undirected edges)	Partial	Partial	Efficient hybrid for ordinal data
swa-LiNGAM	Continuous, non-Gaussian	Yes	No	No	Continuous non-Gaussian outcomes

For ordinal psychological and health survey data — the primary target of this paper — the swa-oBN is the recommended method. It uniquely combines: (1) survey-weight correction via pseudo-BIC with Kish effective sample size; (2) full causal identifiability from the ordinal regression structure; (3) confounder adjustment via the $\mathbf{Z}$ matrix; (4) edge blacklisting for domain knowledge constraints; and (5) bootstrapped uncertainty quantification. The swa-PC and swa-LiNGAM methods serve as complementary tools for sensitivity analysis and for settings with continuous variables.

3.5 3.5 Implementation

All methods are implemented in the open-source R package **OrdCDsvy**, available on CRAN. The primary user-facing function is `OrdCD()`, which extends the `OrdCD` function of Ni et al. (2025) with

the `weights`, `Z`, `blacklist`, and `boot` arguments. Survey-weighted PC and FCI are implemented via the `swa_pc()` and `swa_fci()` wrapper functions, which interface directly with the `pcalg` package (Kalisch et al., 2012). Survey-weighted LiNGAM is available via `swa_lim()`. All functions accept standard R `svydesign` objects from the survey package (Lumley, 2004) or raw weight vectors. Computation is parallelizable across bootstrap resamples via the `parallel` argument. The package includes vignettes reproducing all simulation studies and the CAHPS application.

That is the complete Section 3. A few notes for your review:

Key decisions embedded in this draft: The pseudo-BIC uses n_{eff} in the penalty, which is theoretically justified through Lumley and Scott (2015) — this is an important citation to verify you can access. The swa-LiNGAM approach through pseudo-population resampling is framed clearly as a different mechanism from the pseudo-BIC used in swa-oBN, which is correct. The confounder adjustment section precisely mirrors your code’s `Z` argument.

The R package name “OrdCDsvy” is a placeholder — you’ll want to decide whether to extend the existing `OrdCD` package or create a new one, and name it accordingly.

One technical flag: The `svyolr` function in R’s survey package computes deviance rather than log-likelihood directly, so in your actual code (and in the paper’s notation), the pseudo-BIC uses $-2 * \text{fit\$deviance} + K * \log(n_{\text{eff}})$ — worth double-checking this matches what `svyolr` returns for `fit$deviance` (it should be the weighted deviance, i.e., $-2\tilde{\ell}$).

Shall I proceed to Section 4 (Simulation Studies)?

References

- Agency for Healthcare Research and Quality (2023), *2023 national healthcare quality and disparities report*, National healthcare quality & disparities report chartbooks, Rockville (MD): Agency for Healthcare Research; Quality (US).
- Anderson, L. M., Lim, K. O., Kummerfeld, E., Crosby, R. D., Crow, S. J., Engel, S. G., Forrest, L., Wonderlich, S. A., and Peterson, C. B. (2023), “Causal discovery analysis: A promising tool in advancing precision medicine for eating disorders,” *International Journal of Eating Disorders*, Wiley Online Library, 56, 2012–2021.
- Benedetti, A., and others (2024), “Incorporating survey weights into causal analysis,” *Statistics in Medicine*.
- Briganti, G., Scutari, M., Epskamp, S., Borsboom, D., Hoekstra, R. H., Golino, H. F., Christensen, A. P., Morvan, Y., Ebrahimi, O. V., Costantini, G., and others (2024), “Network analysis: An overview for mental health research,” *International Journal of Methods in Psychiatric Research*, Wiley Online Library, 33, e2034.

- Bühlmann, P., Peters, J., and Ernest, J. (2014), "CAM: Causal additive models, high-dimensional order search and penalized regression," *The Annals of Statistics*, Institute of Mathematical Statistics, 42.
- Chickering, D. M. (2002), "Optimal structure identification with greedy search," *Journal of Machine Learning Research*, 3, 507–554.
- Coffman, D. L., Zhou, H., Castellano, K. E., Schuler, M. S., and McCaffrey, D. F. (2024), "Sampling weighting strategies in causal mediation analysis," *BMC Medical Research Methodology*, Springer, 24, 133.
- Duan-Porter, W., Martinson, B. C., Greer, N., Taylor, B. C., Ullman, K., McKenzie, L., Rosebush, C., MacDonald, R., Falde, S., and Wilt, T. J. (2018), "Evidence review—social determinants of health for veterans," *Journal of general internal medicine*, Springer, 33, 1785–1795.
- Glymour, C., Zhang, K., and Spirtes, P. (2019), "Review of causal discovery methods based on graphical models," *Frontiers in Genetics*, Frontiers, 10, 524.
- Groves, R. M., Fowler Jr, F. J., Couper, M. P., Lepkowski, J. M., Singer, E., and Tourangeau, R. (2009), *Survey methodology*, John Wiley & Sons.
- Horvitz, D. G., and Thompson, D. J. (1952), "A generalization of sampling without replacement from a finite universe," *Journal of the American statistical Association*, Taylor & Francis, 47, 663–685.
- Kish, L. (1965), *Survey sampling*, John Wiley & Sons.
- Liang, W., and Wu, C. (2026), "Causal inference with survey data: A robust framework for propensity score weighting in probability and non-probability samples," *Statistics in Medicine*, Wiley Online Library, 45, e70420.
- Lumley, T. (2004), "Analysis of complex survey samples," *Journal of Statistical Software*, 9, 1–19.
- Montgomery, A. E., Tsai, J., and Blosnich, J. R. (2020), "Demographic correlates of veterans' adverse social determinants of health," *American Journal of Preventive Medicine*, Elsevier BV, 59, 828–836.
- Ni, Y. (2023), *OrdCD: Ordinal causal discovery*.
- Ni, Y., Chen, S., and Wang, Z. (2025), "Causal structural modeling of survey questionnaires via a bootstrapped ordinal bayesian network approach," *Psychometrika*, Springer, 90, 229–250.
- Ni, Y., and Mallick, B. (2022), "Ordinal causal discovery," in *Proceedings of the thirty-eighth conference on uncertainty in artificial intelligence*, PMLR, pp. 1530–1540.
- Pearl, J. (2009), *Causality: Models, reasoning, and inference*, Cambridge University Press.
- Quintana, R. (2023), "Causal search algorithms in social and behavioral sciences," *Sociological Methods & Research*, SAGE Publications.
- Shimizu, S., Hoyer, P. O., Hyvärinen, A., and Kerminen, A. (2006), "A linear non-gaussian acyclic model for causal discovery," *Journal of Machine Learning Research*, 7, 2003–2030.
- Spirtes, P., and Glymour, C. (1991), "An algorithm for fast recovery of sparse causal graphs," *Social science computer review*, Sage Publications Sage CA: Thousand Oaks, CA, 9, 62–72.
- Spirtes, P., Glymour, C. N., and Scheines, R. (2001), *Causation, prediction, and search*, MIT Press.

- Spirtes, P., Meek, C., and Richardson, T. (1995), "Causal inference in the presence of latent variables and selection bias," *Proceedings of the Eleventh Conference on Uncertainty in Artificial Intelligence*, Morgan Kaufmann Publishers Inc., 499–506.
- Uhler, C., Raskutti, G., Bühlmann, P., and Yu, B. (2013), "Geometry of the faithfulness assumption in causal inference," *The Annals of Statistics*, JSTOR, 436–463.
- Vo, T., and others (2024), "Data collection, weighting, and modeling techniques to estimate unbiased population parameters," *Transportation Research Methods*.
- Wang, L., and others (2009), "Propensity score weighting with survey weighted data," *Observational Studies & Survey Data Methods*.
- Washington, D. L. (ed.) (2022), *National veteran health equity report 2021: Focus on veterans health administration patient experience and health care quality*, Government Report, Washington, DC: VHA Office of Health Equity.
- Zanga, A., Ochal, M., Baig, A. R., and Stella, F. (2022), "A survey on causal discovery: Theory and practice," *International Journal of Approximate Reasoning*, Elsevier, 151, 101–129.
- Zanutto, E. L. (2006), "A comparison of propensity score and linear regression analysis of complex survey data," *Journal of data Science*, 4, 67–91.
- Zanutto, E. L., and others (2005), "Using propensity score subclassification for multiple treatment doses to evaluate a national antidrug media campaign," *Journal of Educational and Behavioral Statistics*, SAGE Publications, 30, 59–73.
- Zhalama, Zhang, J., Eberhardt, F., and Mayer, W. (2017), "SAT-based causal discovery under weaker assumptions." in *UAI*.
- Zhu, G., Tay, L. Q., and Zhang, M. (2026), "Causal discovery methods in psychological research: Foundations, algorithms, and a practical tutorial in R," *Behavior Research Methods*, Springer, 58, 64.