

(a)
Tb: raw GC estimate
All molecules (no fitting)
Parity
GC estimate, Tb (K)
Experimental Tb (K)
$R^2_{GC} = 0.77$
$MAE_{GC} = 25.80$
$RMSE_{GC} = 32.85$
(b)
Tm: raw GC estimate
All molecules (no fitting)
Parity
GC estimate, Tm (K)
Experimental Tm (K)
$R^2_{GC} = -0.33$
$MAE_{GC} = 77.82$
$RMSE_{GC} = 102.30$
(c)
Pc: raw GC estimate
All molecules (no fitting)
Parity
GC estimate, Pc (bar)
Experimental Pc (bar)
$R^2_{GC} = 0.82$
$MAE_{GC} = 3.37$
$RMSE_{GC} = 5.43$
(d)
Vc: raw GC estimate
All molecules (no fitting)
Parity
GC estimate, Vc (cm³/mol)
Experimental Vc (cm³/mol)
$R^2_{GC} = 0.98$
$MAE_{GC} = 12.98$
$RMSE_{GC} = 20.59$
(e)
Tc: raw GC estimate
All molecules (no fitting)
Parity
GC estimate, Tc (K)
Experimental Tc (K)
$R^2_{GC} = 0.88$
$MAE_{GC} = 23.41$
$RMSE_{GC} = 30.36$
(f)
ΔHvap: raw GC estimate
All molecules (no fitting)
Parity
GC estimate, Hvap (kJ/mol)
Experimental Hvap (kJ/mol)
$R^2_{GC} = 0.88$
$MAE_{GC} = 1.94$
$RMSE_{GC} = 2.40$
(g)
log P: raw GC estimate
All molecules (no fitting)
Parity
GC estimate, logP
Experimental logP
$R^2_{GC} = 0.86$
$MAE_{GC} = 0.47$
$RMSE_{GC} = 0.75$