

# ECMM450 Stochastic Processes: Assignment 1

700054986

2023-02-19

1 Consider the discrete random variable  $X$  that has the probability generating function

$$G_X(\theta) = 2(3 - \theta)^{-1}$$

(a) Suppose that  $Y = X^2$ . Calculate  $P(Y = k)$  for  $0 \leq k \leq 10$  and the expectation  $E[Y]$

$$\begin{aligned} G_X(\theta) &= 2(3 - \theta)^{-1} \\ &= \frac{2}{3} \left(1 - \frac{\theta}{3}\right)^{-1} \\ &= \frac{2}{3} \left(1 + \frac{\theta}{3} + \left(\frac{\theta}{3}\right)^2 + \left(\frac{\theta}{3}\right)^3 + \dots\right) \\ &= \frac{2}{3} \theta^0 + \frac{2}{3^2} \theta^1 + \frac{2}{3^3} \theta^2 + \frac{2}{3^4} \theta^3 + \dots \end{aligned}$$

We observe from the above equation that  $P(X = 0) = \frac{2}{3}, P(X = 1) = \frac{2}{3^2}, P(X = 2) = \frac{2}{3^3}, \dots$

$$G_Y(\theta) = \sum_k \theta^k P(Y = k) = \sum_k \theta^k P(X^2 = k)$$

$$= \theta^0 P(X^2 = 0) + \theta^1 P(X^2 = 1) + \dots + \theta^4 P(X^2 = 4) + \dots + \theta^9 P(X^2 = 9) + \theta^{10} P(X^2 = 10)$$

As  $X$  can only take on positive integer values, the only terms in the above equation which are non-zero are those associated where  $X^2 \in \{0, 1, 4, 9\}$

$$\text{Therefore, } G_Y(\theta) = \theta^0 P(X^2 = 0) + \theta^1 P(X^2 = 1) + \theta^4 P(X^2 = 4) + \theta^9 P(X^2 = 9)$$

$$G_Y(\theta) = \theta^0 P(X = 0) + \theta^1 P(X = 1) + \theta^4 P(X = 2) + \theta^9 P(X = 3)$$

$$G_Y(\theta) = \frac{2}{3} \theta^0 + \frac{2}{3^2} \theta^1 + \frac{2}{3^3} \theta^4 + \frac{2}{3^4} \theta^9 \quad \mathbf{Ans}$$

$$\begin{aligned} \text{Now, to calculate } E[Y], \text{ we evaluate } G'_Y(\theta)|_{\theta=1} &= \left[\frac{2}{3^2} + 4\theta^3 \frac{2}{3^3} + 9\theta^8 \frac{2}{3^4}\right]_{\theta=1} \\ &= \frac{2}{9} + \frac{4 \cdot 2}{27} + \frac{9 \cdot 2}{81} = 0.7407 \quad \mathbf{Ans} \end{aligned}$$

(b) Suppose that  $Z$  is the random variable with probability generating function  $G_Z(\theta) = G_X(G_X(\theta))$ . Calculate  $P(Z = 0), P(Z = 2)$ , and  $E[Z]$ .

$$G_Z(\theta) = G_X(2(3 - \theta)^{-1}) = 2\left(3 - \frac{2}{3 - \theta}\right)^{-1} = \frac{2(3 - \theta)}{3(3 - \theta) - 2} = \frac{2(3 - \theta)}{7 - 3\theta}$$

$$\implies G_Z(\theta) = \frac{2}{3} \frac{9 - 3\theta}{7 - 3\theta} = \frac{2}{3} \left(1 + \frac{2}{7 - 3\theta}\right) = \frac{2}{3} + \frac{2}{3} \cdot 2 \cdot (7 - 3\theta)^{-1}$$

$$\implies G_Z(\theta) = \frac{2}{3} + \frac{4}{3} \cdot \left(7\left(1 - \frac{3}{7}\theta\right)\right)^{-1} = \frac{2}{3} + \frac{4}{3} \cdot \frac{1}{7} \left(1 - \frac{3}{7}\theta\right)^{-1}$$

$$\implies G_Z(\theta) = \frac{2}{3} + \frac{4}{21} \cdot \left(1 + \frac{3}{7}\theta + \left(\frac{3}{7}\theta\right)^2 + \left(\frac{3}{7}\theta\right)^3 + \dots\right)$$

$$\implies G_Z(\theta) = \frac{2}{3} + \frac{4}{21} + \frac{4}{21} \cdot \left(\frac{3}{7}\theta + \left(\frac{3}{7}\theta\right)^2 + \left(\frac{3}{7}\theta\right)^3 + \dots\right)$$

$$\Rightarrow G_Z(\theta) = \frac{18}{21} + \frac{4}{21} \cdot (\frac{3}{7}\theta + (\frac{3}{7}\theta)^2 + (\frac{3}{7}\theta)^3 + \dots)$$

From the above equation, we see that  $P(Z = 0) = \frac{18}{21} = 0.8571$  **Ans**,  $P(Z = 2) = \frac{4}{21} \cdot \frac{3^2}{7^2} = 0.0349$  **Ans**

Since  $G_Z(\theta) = G_X^2(\theta)$ , we know that  $E[Z] = \mu_X^2$ , where  $\mu_X = E[X]$ .

To find this, we evaluate  $G'_X(\theta)|_{\theta=1} = \frac{d}{d\theta}(2(3-\theta)^{-1}) = 2 \frac{d}{d\theta} \frac{1}{3-\theta} = 2 \cdot (-1) \cdot \frac{1}{(3-\theta)^2} \cdot (-1) = \frac{2}{(3-\theta)^2}$

Putting the value of  $\theta = 1$ , we have,  $E[X] = \frac{2}{(3-1)^2} = \frac{2}{2^2} = 0.5$  Therefore,  $E[Z] = 0.5^2 = 0.25$  **Ans**

(c)  $W = X_1 + X_2 + X_3$ , where  $X_1, X_2, X_3$  are independent

Therefore,  $G_W(\theta) = \{2(3-\theta)^{-1}\}^3 = 2^3(3-\theta)^{-3} = 8(3-\theta)^{-3} = \frac{8}{27}(1 - \frac{\theta}{3})^{-3}$

Using binomial expansion for negative coefficients, we have

$$G_W(\theta) = \frac{8}{27}(1 + (-3)(-\frac{\theta}{3}) + \frac{(-3)(-3-1)}{2!} \frac{\theta^2}{3^2} + \dots)$$

$$G_W(\theta) = \frac{8}{27}(1 + \theta + \frac{2}{3}\theta^2 + \dots)$$

From the above equation, it can be observed that

$$P(W = 0) = \frac{8}{27} = 0.2962 \text{ **Ans**}, P(W = 2) = \frac{8 \cdot 2}{27 \cdot 3} = 0.1975 \text{ **Ans**}$$

For  $E[W]$ , we have  $G'_W(\theta)|_{\theta=1} = 2^3(-3)(3-\theta)^{-4}(-1)|_{\theta=1} = \frac{24}{(3-1)^4} = \frac{24}{16} = 1.5$  **Ans**

**2** Suppose that  $X$  is a discrete random variable with probability generating function:  $G_X(\theta) = \frac{1-2\lambda}{1-\lambda\theta-\lambda\theta^2}$ , where  $\lambda > 0$  is a constant. Consider a branching process  $\{S_n\}_{n \geq 0}$  denoting the number of individuals at stage  $n$  (notation as in lectures), with  $S_0 = 1$ . Assume the number of offspring  $X$  arising from any individual in any generation has the probability distribution governed by  $G_X(\theta)$ , and that all individuals evolve independently of all others.

(a) Write down the range of values of  $\lambda$  for which  $G_X$  is a valid probability generating function.

$$G_X(\theta) = (1-2\lambda)(1-\lambda\theta^2-\lambda\theta)^{-1} = (1-2\lambda)(1-\lambda\theta(\theta+1))^{-1}$$

Expanding, we have

$$G_X(\theta) = (1-2\lambda)[1 + \lambda\theta(\theta+1) + (\lambda\theta(\theta+1))^2 + (\lambda\theta(\theta+1))^3 + \dots]$$

It is clear from the above equation that  $P(X = 0)$  or the coefficient of  $\theta^0$  is  $(1-2\lambda)$

For  $G_X(\theta)$  to be a valid PGF,  $0 \leq (1-2\lambda) \leq 1 \Rightarrow 0 < \lambda \leq \frac{1}{2} \Rightarrow \lambda \in [0, \frac{1}{2}]$  **Ans**

$$(b) \text{ For } \lambda = \frac{1}{4}, G_X(\theta) = \frac{1-2\lambda}{1-\lambda\theta-\lambda\theta^2} = \frac{1-2 \cdot \frac{1}{4}}{1-\frac{1}{4}\theta-\frac{1}{4}\theta^2} = \frac{2}{4-\theta-\theta^2}$$

$$\mu_X = G'_X(\theta)|_{\theta=1} = \frac{2}{(4-\theta-\theta^2)^2} \cdot (-1) \cdot (-1-2\theta) = \frac{2}{(4-\theta-\theta^2)^2} \cdot (1+2\theta)$$

Putting the value of  $\theta$ ,  $\mu_X = \frac{2}{(4-1-1^2)^2}(1+2 \cdot 1) = \frac{6}{4} = 1.5$

$$E[S_n] = \mu_X^n = (1.5)^n \text{ **Ans**}$$

$$G_{S1}(\theta) = G_X^1(\theta) = G_X(\theta) = (1-2\lambda)[1 + \lambda\theta(\theta+1) + (\lambda\theta(\theta+1))^2 + (\lambda\theta(\theta+1))^3 + \dots]$$

$$\Rightarrow G_{S1}(\theta) = (1-2\lambda)[1 + \lambda\theta(\theta+1) + \lambda^2\theta^2(\theta+1)^2 + \dots]$$

$$\Rightarrow G_{S1}(\theta) = (1-2\lambda)[1 + \lambda\theta^2 + \lambda\theta + \lambda^2\theta^2(\theta^2 + 2\theta + 1) + \dots]$$

$$\Rightarrow G_{S1}(\theta) = (1-2\lambda)[1 + \lambda\theta^2 + \lambda\theta + \lambda^2\theta^2(\theta^2 + 2\theta + 1) + \dots]$$

$$\Rightarrow G_{S1}(\theta) = (1-2\lambda)[1 + \lambda\theta^2 + \lambda\theta + \lambda^2\theta^4 + 2\lambda^2\theta^3 + \lambda^2\theta^2 + \dots]$$

Here,  $P(S_1 = 2)$  is the coefficient of  $\theta^2$ , which is  $(1 - 2\lambda)(\lambda^2 + \lambda) = (1 - \frac{2}{4})(\frac{1}{16} + \frac{1}{4}) = 0.5 * 0.3125 = 0.15625$   
**Ans**

Probability of extinction is given by the smallest root of the equation  $G_X(e) = e$

$$G_X(e) = e \implies \frac{1-2\lambda}{1-\lambda e-\lambda e^2} = e \implies 1-2\lambda = e \cdot (1-\lambda e-\lambda e^2) \implies 1-2\lambda = e - \lambda e^2 - \lambda e^3 \implies \lambda e^3 + \lambda e^2 - e - 2\lambda + 1 = 0$$

We know that  $(e-1)$  must always be a factor as the probability of extinction is upper-bounded by 1.

Factorizing,

$$(e-1)(\lambda e^2 + 2\lambda e - 1 + 2\lambda) = 0$$

Now, either  $e = 1$  or  $(\lambda e^2 + 2\lambda e - 1 + 2\lambda) = 0$

Using the Sridharacharya formula for solving quadratics, we have

$$e = \frac{-2\lambda \pm \sqrt{4\lambda^2 - 4\lambda(2\lambda-1)}}{2\lambda}$$

$$e = \frac{-2\lambda \pm \sqrt{4\lambda^2 - 4\lambda^2 \cdot \frac{1}{\lambda}(2\lambda-1)}}{2\lambda}$$

$$e = \frac{-2\lambda \pm \sqrt{4\lambda^2(1 - \frac{1}{\lambda}(2\lambda-1))}}{2\lambda}$$

$$e = \frac{-2\lambda \pm 2\lambda \sqrt{(1 - \frac{1}{\lambda}(2\lambda-1))}}{2\lambda}$$

$$e = -1 \pm \sqrt{(1 - \frac{1}{\lambda}(2\lambda-1))}$$

$$e = -1 \pm \sqrt{(1 - 2 + \frac{1}{\lambda})}$$

$$e = -1 \pm \sqrt{(\frac{1}{\lambda} - 1)}$$

Using  $\lambda = \frac{1}{4}$ , we have  $e = -1 \pm \sqrt{3}$ . Since the negative solution is nonsensical,  $e = -1 + \sqrt{3} = 0.732$  **Ans**

(c) Compute the range of values of  $\lambda$  for which the population is guaranteed to become extinct

To achieve this, we need  $\mu_X < 1$

$$\mu_X = G'_x(\theta)|_{\theta=1} = \frac{1-2\lambda}{(1-\lambda\theta-\lambda\theta^2)^2} \cdot (-1) \cdot (-\lambda - 2\lambda\theta) < 1$$

$$\text{Substituting } \theta = 1, \text{ we have } \mu_X = \frac{1-2\lambda}{(1-\lambda-\lambda)^2}(\lambda + 2\lambda) = \frac{1-2\lambda}{(1-2\lambda)^2}3\lambda < 1$$

$$(1-2\lambda)(3\lambda) < (1-2\lambda)^2$$

$$(1-2\lambda)(3\lambda) - (1-2\lambda)^2 < 0$$

$$(1-2\lambda)(3\lambda - (1-2\lambda)) < 0$$

$$(1-2\lambda)(5\lambda - 1) < 0$$

$$(2\lambda - 1)(5\lambda - 1) > 0$$

$$(\lambda - \frac{1}{2})(\lambda - \frac{1}{5}) > 0$$

This inequality yields  $\lambda \in (-\infty, \frac{1}{5}) \cap (\frac{1}{2}, \infty)$  Combining this with the earlier bounded region for  $\lambda$  where  $(\lambda \in [0, \frac{1}{2}))$  for  $G_X(\theta)$  to be a valid PGF, we have  $\lambda \in [0, \frac{1}{5})$  **Ans**

(d) For ultimate extinction as a function of  $\lambda$ , we refer to part (b). It is  $-1 + \sqrt{\frac{1}{\lambda} - 1}$  **Ans**

**3** At the University of Exeter, the frequency of cars passing the Harrison Building are recorded. It is noticed that Mercedes, BMWs and Ferraris pass by at a combined rate of 8 cars per hour. Out of these cars, it is noticed that 40% are Mercedes, 50% are BMWs and the remaining 10% are Ferraris. You may assume that all passing car types are independent Poisson processes.

(a) Find the probability that the next car observed is a Mercedes.

Since all of the passing cars follow independent Poisson processes,  $P(\text{next Mercedes}) = 0.4$  **Ans**

- (b) Given that the last car observed was a BMW, find the probability that the next car to be observed passing the Harrison Building is a Ferrari.

We have independence of events in the future from events in the past. Therefore  $P(\text{next car Ferrari}) = 0.1$  **Ans**

- (c) Calculate the probability that exactly 2 BMWs pass by the Harrison Building in 30 minutes.

The combined rate of passing is known to be  $\lambda_{combined} = 8/hr$ . Since BMWs account for 50% of the volume,  $\lambda_{BMW} = 0.5 * 8 = 4/hr$

Recall that formula for Poisson process is  $P((N = t) = k) = \frac{(\lambda t)^k e^{-\lambda t}}{k!}$  Hence,  $P(N(t = 1/2) = 2) = \frac{(4 \cdot \frac{1}{2})^2 \cdot e^{-2}}{2!} = 2e^{-2} = 0.2706$  **Ans**

- (d) Calculate the probability that at least three cars (of any of those three types) pass by the Harrison building in 45 minutes.

$$P(N(t = \frac{3}{4}) \geq 3) = 1 - P(N(t = \frac{3}{4}) < 3) = 1 - P(N(t = \frac{3}{4}) \leq 2) = 1 - \sum_{k=0}^2 P(N(t = \frac{3}{4}) = k)$$