

- Joint inflammation can occur weeks later (leading to an unusual form of arthritis).
- Infection of vessels and heart valves is a special characteristic of *C. fetus*. Immunocompromised patients may develop repeated episodes of passage of bacteria into the bloodstream from these sites of infection.
- The gallbladder, pancreas, and bone may be affected.

Diagnosis

Campylobacter is only one of many causes of acute diarrhea. Culture (growing the bacteria in the laboratory) of freshly obtained diarrhea fluid is the only way to be certain of the diagnosis.

Treatment

The first aim of treatment is to keep up **nutrition** and avoid dehydration. Medications used to treat diarrhea by decreasing intestinal motility, such as Loperamide or Diphenoxylate are also useful, but should only be used with the advice of a physician. **Antibiotics** are of value, if started within three days of onset of symptoms. They are indicated for those with severe or persistent symptoms. Either an erythromycin type drug or one of the **fluoroquinolones** (such as ciprofloxacin) for five to seven days are the accepted therapies.

Prognosis

Most patients with *Campylobacter* infection rapidly recover without treatment. For certain groups of patients, infection becomes chronic and requires repeated courses of antibiotics.

Prevention

Good hand washing technique as well as proper preparation and cooking of food is the best way to prevent infection.

Resources

BOOKS

- Blaser, Martin J. "Infections due to *Campylobacter* and Related Species." In *Harrison's Principles of Internal Medicine*, ed. Anthony S. Fauci, et al. New York: McGraw-Hill, 1997.
- Hamer, Davidson H., and Sherwood L. Gorbach. "Campylobacter." In *Sleisenger & Fordtran's Gastrointestinal and Liver Disease*, ed. Mark Feldman, et al. Philadelphia: W. B. Saunders Co., 1997.
- Thielman, Nathan M., and Richard L. Guerrant. "Food-Borne Illness." In *Conn's Current Therapy, 1996*, ed. Robert E. Rakel. Philadelphia: W. B. Saunders Co., 1996.

Wolfe, Martin S. "Acute Infectious Diarrhea." In *Conn's Current Therapy, 1996*, ed. Robert E. Rakel. Philadelphia: W. B. Saunders Co., 1996.

PERIODICALS

- Lew, Edward A., Michael A. Poles, and Douglas T. Dieterich. "Diarrheal Disease Associated with HIV Infection." *Gastroenterology Clinics of North America* (June 1997): 259-290.
- "Traveler's Diarrhea: Don't Let It Ruin Your Trip." *Mayo Clinic Health Letter* (Jan. 1997).
- "When Microbes are on the Menu." *Harvard Health Letter* (Dec. 1994) : 4-5.

ORGANIZATIONS

Centers for Disease Control and Prevention. 1600 Clifton Rd., NE, Atlanta, GA 30333. (800) 311-3435, (404) 639-3311. <<http://www.cdc.gov>>.

OTHER

Centers for Disease Control. <<http://www.cdc.gov/nccddphp/ddt/ddthome.htm>>.

David Kaminstein, MD

Cancer

Definition

Cancer is not just one disease, but a large group of almost one hundred diseases. Its two main characteristics are uncontrolled growth of the cells in the human body and the ability of these cells to migrate from the original site and spread to distant sites. If the spread is not controlled, cancer can result in **death**.

Description

One out of every four deaths in the United States is from cancer. It is second only to heart disease as a cause of death in the states. About 1.2 million Americans are diagnosed with cancer annually; more than 500,000 die of cancer annually.

Cancer can attack anyone. Since the occurrence of cancer increases as individuals age, most of the cases are seen in adults, middle-aged or older. Sixty percent of all cancers are diagnosed in people who are older than 65 years of age. The most common cancers are skin cancer, lung cancer, **colon cancer**, **breast cancer** (in women), and **prostate cancer** (in men). In addition, cancer of the kidneys, ovaries, uterus, pancreas, bladder, rectum, and blood and lymph node cancer (leukemias and lymphomas) are also included among the 12 major cancers that affect most Americans.

Cancer, by definition, is a disease of the genes. A gene is a small part of DNA, which is the master molecule of the cell. Genes make “proteins,” which are the ultimate workhorses of the cells. It is these proteins that allow our bodies to carry out all the many processes that permit us to breathe, think, move, etc.

Throughout people’s lives, the cells in their bodies are growing, dividing, and replacing themselves. Many genes produce proteins that are involved in controlling the processes of cell growth and division. An alteration (mutation) to the DNA molecule can disrupt the genes and produce faulty proteins. This causes the cell to become abnormal and lose its restraints on growth. The abnormal cell begins to divide uncontrollably and eventually forms a new growth known as a “tumor” or neoplasm (medical term for cancer meaning “new growth”).

In a healthy individual, the immune system can recognize the neoplastic cells and destroy them before they get a chance to divide. However, some mutant cells may escape immune detection and survive to become tumors or cancers.

Tumors are of two types, benign or malignant. A benign tumor is not considered cancer. It is slow growing, does not spread or invade surrounding tissue, and once it is removed, it doesn’t usually recur. A malignant tumor, on the other hand, is cancer. It invades surrounding tissue and spreads to other parts of the body. If the cancer cells have spread to the surrounding tissues, then, even after the malignant tumor is removed, it generally recurs.

A majority of cancers are caused by changes in the cell’s DNA because of damage due to the environment. Environmental factors that are responsible for causing the initial mutation in the DNA are called carcinogens, and there are many types.

There are some cancers that have a genetic basis. In other words, an individual could inherit faulty DNA from his parents, which could predispose him to getting cancer. While there is scientific evidence that both factors (environmental and genetic) play a role, less than 10% of all cancers are purely hereditary. Cancers that are known to have a hereditary link are breast cancer, colon cancer, **ovarian cancer**, and uterine cancer. Besides genes, certain physiological traits could be inherited and could contribute to cancers. For example, inheriting fair skin makes a person more likely to develop skin cancer, but only if they also have prolonged exposure to intensive sunlight.

There are several different types of cancers:

- Carcinomas are cancers that arise in the epithelium (the layers of cells covering the body’s surface and lining the internal organs and various glands). Ninety percent of human cancers fall into this category. Carcinomas can be

subdivided into two types: adenocarcinomas and squamous cell carcinomas. Adenocarcinomas are cancers that develop in an organ or a gland, while squamous cell carcinomas refer to cancers that originate in the skin.

- Melanomas also originate in the skin, usually in the pigment cells (melanocytes).
- Sarcomas are cancers of the supporting tissues of the body, such as bone, muscle and blood vessels.
- Cancers of the blood and lymph glands are called leukemias and lymphomas respectively.
- Gliomas are cancers of the nerve tissue.

Causes and symptoms

The major risk factors for cancer are: tobacco, alcohol, diet, sexual and reproductive behavior, infectious agents, family history, occupation, environment and pollution.

According to the estimates of the American Cancer Society (ACS), approximately 40% of the cancer deaths in 1998 will be due to tobacco and excessive alcohol use. An additional one-third of the deaths will be related to diet and **nutrition**. Many of the one million skin cancers that are expected to be diagnosed in 1998 will be due to over-exposure to ultraviolet light from the sun’s rays.

Tobacco

Eighty to ninety percent of the lung cancer cases occur in smokers. **Smoking** has also been shown to be a contributory factor in cancers of upper respiratory tract, esophagus, larynx, bladder, pancreas, and probably liver, stomach, and kidney as well. Recently, scientists have also shown that second-hand smoke (or passive smoking) can increase one’s risk of developing cancer.

Alcohol

Excessive consumption of alcohol is a risk factor in certain cancers, such as **liver cancer**. Alcohol, in combination with tobacco, significantly increases the chances that an individual will develop mouth, pharynx, larynx and esophageal cancers.

Diet

Thirty-five percent of all cancers are due to dietary causes. Excessive intake of fat leading to **obesity** has been associated with cancers of the breast, colon, rectum, pancreas, prostate, gall bladder, ovaries and uterus.

Sexual and reproductive behavior

The human papilloma virus, which is sexually transmitted, has been shown to cause cancer of the cervix.

Having too many sex partners and becoming sexually active early has been shown to increase one's chances of contracting this disease. In addition, it has also been shown that women who don't have children or have children late in life have an increased risk for both ovarian and breast cancer.

Infectious agents

In the last 20 years, scientists have obtained evidence to show that approximately 15% of the world's cancer deaths can be traced to viruses, bacteria, or parasites. The most common cancer-causing pathogens and the cancers associated with them are shown in table form.

Family history

Certain cancers like breast, colon, ovarian and uterine cancer recur generation after generation in some families. A few cancers, such as the **eye cancer** "retinoblastoma," a type of colon cancer, and a type of breast cancer known as "early-onset breast cancer," have been shown to be linked to certain genes that can be tracked within a family. It is therefore possible that inheriting particular genes makes a person susceptible to certain cancers.

Occupational hazards

There is evidence to prove that certain occupational hazards account for 4% of all cancer deaths. For example, asbestos workers have an increased incidence of lung cancer. Similarly, a higher likelihood of getting **bladder cancer** is associated with dye, rubber and gas workers; skin and lung cancer with smelters, gold miners and arsenic workers; leukemia with glue and varnish workers; liver cancer with PVC manufacturers; and lung, bone and bone marrow cancer with radiologists and uranium miners.

Environment

Radiation is believed to cause 1–2% of all cancer deaths. Ultra-violet radiation from the sun accounts for a majority of melanoma deaths. Other sources of radiation are x rays, radon gas, and ionizing radiation from nuclear material.

Pollution

Several studies have shown that there is a well-established link between asbestos and cancer. Chlorination of water may account for a small rise in cancer risk. However, the main danger from pollution occurs when dangerous chemicals from the industries escape into the surrounding environment. It has been estimated that 1% of cancer deaths are due to air, land and water pollution.

Frequency Of Cancer-Related Death

Cancer Site	Number of Deaths Per Year
Lung	160,100
Colon and rectum	56,500
Breast	43,900
Prostate	39,200
Pancreas	28,900
Lymphoma	26,300
Leukemia	21,600
Brain	17,400
Stomach	13,700
Liver	13,000
Esophagus	11,900
Bladder	12,500
Kidney	11,600
Multiple myeloma	11,300

Cancer is a progressive disease, and goes through several stages. Each stage may produce a number of symptoms. Some symptoms are produced early and may occur due to a tumor that is growing within an organ or a gland. As the tumor grows, it may press on the nearby nerves, organs and blood vessels. This causes **pain** and some pressure which may be the earliest warning signs of cancer.

Despite the fact that there are several hundred different types of cancers, producing very different symptoms, the ACS has established the following seven symptoms as possible warning signals of cancer:

- changes in the size, color, or shape of a wart or a mole
- a sore that does not heal
- persistent **cough**, hoarseness, or **sore throat**
- a lump or thickening in the breast or elsewhere
- unusual bleeding or discharge
- chronic **indigestion** or difficulty in swallowing
- any change in bowel or bladder habits

Many other diseases, besides cancer, could produce the same symptoms. However, it is important to have these symptoms checked, as soon as possible, especially if they linger. The earlier a cancer is diagnosed and treated, the better the chance of it being cured. Many cancers such as breast cancer may not have any early symptoms. Therefore, it is important to undergo routine screening tests such as breast self-exams and mammograms.

Diagnosis

Diagnosis begins with a thorough **physical examination** and a complete medical history. The doctor will observe, feel and palpate (apply pressure by touch) different parts of the body in order to identify any variations from the normal size, feel and texture of the organ or tissue.