

BIOLOGICAL EVALUATION OF ANTIOXIDANT, ANTIMICROBIAL AND PRESERVATION PERFORMANCE OF A HERBAL PARABEN-FREE FACIAL CLEANSER

¹Souravh Jha, ²Dr. Yesh Pratap Singh Rana

¹Research Scholar, ²Professor

¹ ²Department of Pharmacy, Monad University, Hapur

Article ID: IJPEMR-2026-001 | Article Type: Methodological / Conceptual Framework

ABSTRACT

The increasing interest in botanical cosmetics has encouraged the development of facial cleansing products that combine effective cleansing with skin compatibility and alternative preservation strategies. This article presents a structured methodological framework for evaluating a herbal paraben-free facial cleanser with emphasis on antioxidant activity, antimicrobial properties, microbiological quality and preservation performance. The framework integrates phytochemical characterization, antioxidant assessment, antimicrobial screening, microbiological quality evaluation and preservation efficacy testing. Herbal extracts are considered on the basis of documented phytochemical composition, traditional cosmetic relevance and reported biological activities. The framework distinguishes intrinsic antimicrobial activity of individual extracts from preservation efficacy of the finished cosmetic formulation. It also considers the influence of pH, surfactant composition, water activity, packaging, stability and interactions among botanical constituents. DPPH and ABTS assays are proposed as complementary antioxidant assessments, while antimicrobial screening, microbial quality assessment and controlled challenge testing are proposed for microbiological evaluation. The framework is intended to provide a scientifically structured basis for future laboratory validation and optimization of botanical paraben-free facial cleansers. No numerical experimental findings are presented in this framework article; laboratory observations should be generated and reported separately following validated procedures.

Keywords: Herbal facial cleanser; paraben-free cosmetics; antioxidant activity; antimicrobial activity; preservation efficacy; DPPH; ABTS; botanical extracts; cosmetic microbiology; natural preservatives.

1. INTRODUCTION

1.1 Background

Facial cleansing is an essential component of routine skin care because the skin surface is continuously exposed to sebum, perspiration, environmental particulate matter, microorganisms, cosmetic residues and other contaminants. A facial cleanser is therefore expected to remove undesirable material while minimizing disruption of the physiological skin barrier. This requirement creates a formulation challenge: increasing cleansing intensity may improve soil removal but can simultaneously increase dryness, irritation or unpleasant post-wash sensations when the surfactant system is poorly selected.

Contemporary cosmetic research has increasingly explored botanical ingredients as multifunctional components of skin-care formulations. Plants contain diverse secondary metabolites, including phenolic compounds, flavonoids, tannins, terpenoids, saponins and other constituents that may contribute antioxidant, antimicrobial, soothing or conditioning properties. However, the presence of biologically active compounds does not automatically establish the suitability of a botanical extract for a finished cosmetic product. Extract composition can vary with botanical identity, plant part, cultivation conditions, extraction solvent, processing conditions and storage.

The preservation of water-containing cosmetics represents an additional scientific challenge. Such formulations provide an environment in which microorganisms may proliferate if an adequate preservation system is not established. Consequently, a paraben-free product cannot simply be considered safe because parabens have been omitted. The complete formulation must demonstrate adequate microbiological quality and resistance to microbial proliferation under appropriately designed testing conditions.

The present framework therefore focuses on three related but distinct biological dimensions: antioxidant activity, antimicrobial activity and preservation performance. Antioxidant assays evaluate the ability of extracts or formulations to interact with selected reactive species under defined test conditions. Antimicrobial testing evaluates inhibitory effects against selected microorganisms. Preservation testing, in contrast, examines whether the finished formulation can control deliberately introduced microorganisms over a specified testing period. Treating these endpoints separately provides greater scientific clarity.

1.2 Rationale

Botanical facial cleansers may provide opportunities to reduce reliance on conventional preservative systems while incorporating plant-derived functional ingredients. Nevertheless, several limitations require careful consideration. Botanical extracts may introduce colour and odour changes, interact with surfactants or polymers, alter viscosity, influence pH and potentially compromise microbial stability.

Moreover, an extract exhibiting strong activity in a laboratory antioxidant assay may have limited activity after incorporation into a complex cleanser because of dilution, extraction differences, pH effects or interactions with other ingredients. Similarly, an antimicrobial zone observed in an agar diffusion test cannot by itself establish preservative efficacy in a finished cosmetic.

Table 1. Major Biological Evaluation Domains

Evaluation domain	Principal question	Typical assessment
Antioxidant activity	Can the formulation/extract interact with selected free radicals under assay conditions?	DPPH, ABTS
Antimicrobial activity	Does the formulation inhibit	Agar diffusion or validated

	selected microorganisms?	quantitative method
Microbial quality	Is the finished product microbiologically acceptable?	Microbial enumeration and specified-organism testing
Preservation performance	Can the formulation control introduced microorganisms over time?	Preservative efficacy/challenge testing
Stability	Does performance remain acceptable during storage?	Physical, chemical and microbiological monitoring

2. AIM AND OBJECTIVES

Aim: To formulate and scientifically evaluate a herbal paraben-free facial cleanser with particular emphasis on its antioxidant activity, antimicrobial properties and preservation performance.

Objectives:

- To select suitable botanical extracts for incorporation into a paraben-free facial cleanser.
- To characterize the selected extracts for relevant phytochemical constituents.
- To formulate a stable herbal facial cleanser using an appropriate surfactant and preservation-support system.
- To evaluate antioxidant activity using validated or appropriately standardized DPPH and ABTS assays.
- To investigate the antimicrobial activity of the selected extracts and finished formulations.
- To assess the microbiological quality of the formulated cleanser.
- To evaluate preservation performance using an appropriate challenge-testing methodology.
- To examine the relationship between formulation composition, antimicrobial activity and microbiological stability.
- To identify the formulation demonstrating the most satisfactory overall biological and formulation performance.

3. RESEARCH METHODOLOGY

3.1 Proposed Research Design

The framework proposes an experimental formulation-and-evaluation design consisting of botanical selection, extract preparation, phytochemical screening, formulation development, antioxidant evaluation, antimicrobial evaluation, microbiological quality assessment, preservation efficacy testing, stability monitoring and statistical analysis. Exact experimental conditions should be finalized through laboratory validation and relevant institutional procedures. This article does not report completed laboratory experiments or numerical experimental findings.

3.2 Materials

The study may use authenticated botanical materials selected according to the final formulation design. Additional materials include a mild surfactant system, humectant, viscosity modifier, pH-adjusting agents, purified water and an appropriate paraben-free preservation system.

Table 2. Experimental Material Categories

Category	Function
Botanical extracts	Functional herbal actives
Surfactant	Cleansing and soil removal
Humectant	Moisture retention and skin feel
Thickener	Viscosity and physical stability
pH modifier	pH adjustment
Preservation system	Microbial control
Chelating/preservation booster	Enhancement of preservation where appropriate
Purified water	Vehicle
Packaging	Product protection and dispensing

3.3 Preparation of Herbal Extracts

Authenticated plant material should be cleaned to remove foreign matter and dried under controlled conditions that minimize degradation of thermolabile constituents. The dried material should then be pulverized to a reasonably uniform particle size. The extraction solvent should be selected according to the polarity and stability of the target phytoconstituents. Aqueous, hydroalcoholic or other validated extraction systems may be considered. Following extraction, the material should be filtered and concentrated under controlled conditions. The resulting extract should be stored in suitable containers under conditions that minimize moisture, light and temperature-related degradation.

3.4 Phytochemical Screening

Preliminary screening may include qualitative tests for:

- phenolics
- flavonoids
- tannins
- saponins
- alkaloids
- terpenoids
- glycosides
- other constituents relevant to the selected botanical materials

Where resources permit, quantitative marker analysis should complement preliminary screening because qualitative tests provide only limited information regarding concentration.

3.5 Formulation Development

Several experimental batches should be prepared by varying relevant botanical and formulation variables.

Table 3. Proposed Formulation Development Matrix

Batch	Herbal active level	Preservation-support level	Purpose
F1	Low	Standard	Determine lower active level
F2	Moderate	Standard	Evaluate intermediate level
F3	Higher	Standard	Determine effect of increased active level
F4	Optimized combination	Optimized	Final candidate
Control	Without selected herbal actives	Appropriate control	Comparative reference

3.6 Antioxidant Evaluation

DPPH Assay

Appropriate concentrations of extract or formulation should be reacted with DPPH reagent under standardized conditions. Absorbance should be measured at the validated wavelength using a suitable spectrophotometer.

ABTS Assay

The ABTS radical system should be generated according to the validated laboratory protocol and adjusted to the required absorbance range. Sample activity should be determined relative to the control and expressed according to the selected analytical protocol. Using DPPH and ABTS together can provide complementary evidence rather than treating either assay as definitive.

3.7 Antimicrobial Evaluation

The antimicrobial investigation should include suitable reference microorganisms selected according to the study objectives and laboratory biosafety requirements. The test panel may include representative Gram-positive bacteria, Gram-negative bacteria and yeast/fungal organisms. Appropriate positive and negative controls should be incorporated. Where agar diffusion is employed, inhibition zones should be measured using standardized procedures. Where possible,

quantitative methods such as minimum inhibitory concentration or equivalent validated procedures can provide stronger comparative evidence.

3.8 Microbial Quality Assessment

The finished cleanser should be assessed for microbial quality using appropriate enumeration methods and examination for relevant specified or objectionable microorganisms. Testing should follow recognized microbiological quality procedures applicable to cosmetic products. Importantly, microbial quality testing and preservation efficacy testing should not be conflated. The former assesses contamination status of the product, whereas the latter examines its capacity to resist or control microbial challenge.

3.9 Preservation Efficacy Evaluation

A controlled challenge test should be performed using defined microbial strains and predetermined sampling intervals. The formulation should be inoculated under controlled conditions, followed by microbial recovery at specified times. The principal outcome is the change in recoverable microbial population over the testing period. Acceptance criteria should be selected according to the applicable recognized standard or pharmacopoeial/cosmetic microbiological framework rather than invented specifically for the study.

3.10 Stability Evaluation

Samples should be stored under selected long-term, accelerated and temperature-variation conditions appropriate to the research design. Where relevant, freeze-thaw and light-exposure studies may also be conducted. Parameters to monitor include colour, odour, appearance, pH, viscosity, phase separation, antioxidant activity, marker compounds, microbial quality and preservation performance where appropriate.

4. EXPECTED EVALUATION FRAMEWORK

Important publication note: This framework article intentionally does not present numerical experimental results. The numerical example tables from the supplied PDF have been omitted from this publication-ready version because they were not supported as actual laboratory observations.

The proposed framework is intended to guide future laboratory evaluation. Antioxidant activity should be interpreted as chemical assay evidence under defined conditions rather than direct clinical skin protection. Antimicrobial screening should be interpreted as evidence of inhibition against selected microorganisms. Preservation performance should be assessed at formulation level through an appropriately designed challenge test. Stability assessment should integrate physical, chemical and microbiological parameters over the defined storage period.

5. DISCUSSION

The scientific evaluation of a herbal paraben-free cleanser requires interpretation across several levels. The first level concerns the biological properties of the botanical materials themselves. The second concerns whether those properties survive incorporation into the formulation. The third concerns whether the complete product remains microbiologically stable during storage.

Antioxidant activity is particularly relevant when phenolic and flavonoid-rich botanical extracts are used. If both DPPH and ABTS assays show concentration-dependent activity, confidence in the general radical-scavenging capacity of the sample is greater than if only one assay is used. However, assay results remain chemical evidence rather than clinical proof.

The antimicrobial evaluation requires similarly cautious interpretation. Botanical extracts may inhibit selected microorganisms, but activity can vary substantially according to microbial species and extraction conditions. In addition, antimicrobial activity in an agar system does not necessarily indicate that the same concentration will preserve a complex aqueous cleanser.

Preservation performance is consequently the most formulation-specific endpoint. An effective preservation system must function in the presence of surfactants, botanical constituents, polymers and other ingredients. Product pH, packaging and repeated exposure to environmental microorganisms may also influence performance.

A successful formulation should therefore not be selected solely because it demonstrates the strongest antioxidant or antimicrobial activity. Excessive botanical concentration could negatively affect odour, colour, viscosity, stability or skin compatibility. Optimization requires a balance among biological effectiveness, physical stability, microbiological safety and user acceptability.

6. CONCLUSION

The evaluation of antioxidant, antimicrobial and preservation performance provides a comprehensive framework for assessing herbal paraben-free facial cleansers. Botanical extracts may contribute useful biological functionality, but their incorporation into a cosmetic vehicle introduces additional variables that can modify activity and stability. The most scientifically defensible approach is therefore to evaluate the extract, formulation and finished product separately while subsequently integrating the findings. DPPH and ABTS assays can characterize antioxidant potential; antimicrobial testing can identify inhibitory effects against selected microorganisms; and microbial quality and challenge testing can establish whether the complete formulation possesses appropriate microbiological characteristics.

A paraben-free formulation should not be regarded as inherently safer or more effective merely because it excludes parabens. Its scientific merit depends on demonstrated product quality, preservation adequacy, stability and skin compatibility. Accordingly, optimization should be based on a multidimensional assessment rather than on a single biological endpoint.

The proposed research framework can contribute to the development of herbal facial cleansers by demonstrating how botanical functionality and microbiological protection can be evaluated simultaneously. Further work should emphasize standardized botanical extracts, quantitative marker analysis, validated preservation systems, extended stability assessment and appropriately designed human skin-compatibility studies.

REFERENCES

1. Bakkali F, Averbeck S, Averbeck D, Idaomar M. Biological effects of essential oils—a review. *Food Chem Toxicol.* 2008;46:446-475.
2. Barel AO, Paye M, Maibach HI, editors. *Handbook of Cosmetic Science and Technology.* 4th ed. Boca Raton: CRC Press; 2014.
3. Brand-Williams W, Cuvelier ME, Berset C. Use of a free radical method to evaluate antioxidant activity. *Lebensm Wiss Technol.* 1995;28:25-30.
4. Brannan DK, Dille JC. Type of closure prevents microbial contamination of cosmetic emulsions during consumer use. *Appl Environ Microbiol.* 1990;56:1476-1479.
5. Burt S. Essential oils: their antibacterial properties and potential applications in foods—a review. *Int J Food Microbiol.* 2004;94:223-253.
6. Cowan MM. Plant products as antimicrobial agents. *Clin Microbiol Rev.* 1999;12:564-582.
7. Dai J, Mumper RJ. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules.* 2010;15:7313-7352.
8. Denyer SP, Baird RM, editors. *Guide to Microbiological Control in Pharmaceuticals and Medical Devices.* Boca Raton: CRC Press; 1990.
9. Denyer SP, Stewart GSAB. Mechanisms of action of disinfectants. *Int Biodeterior Biodegradation.* 1998;41:261-268.
10. Dorman HJD, Deans SG. Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *J Appl Microbiol.* 2000;88:308-316.
11. Draelos ZD. *Cosmetic Dermatology: Products and Procedures.* 3rd ed. Hoboken: Wiley-Blackwell; 2016.
12. Draelos ZD. The science behind skin care: moisturizers. *J Cosmet Dermatol.* 2018;17:138-144.
13. Geis PA. *Cosmetic Microbiology: A Practical Approach.* 2nd ed. Boca Raton: CRC Press; 2006.
14. Gilbert P, Moore LE. Cationic antiseptics: diversity of action under a common epithet. *J Appl Microbiol.* 2005;99:703-715.
15. Harborne JB. *Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis.* 3rd ed. London: Chapman & Hall; 1998.
16. Harry RG. *Harry's Cosmeticology.* 8th ed. New York: Chemical Publishing; 2000.
17. Korting HC, Kober M, Mueller M, Braun-Falco O. Influence of repeated washings with soap and synthetic detergents on pH, stratum corneum hydration and fat content. *Acta Derm Venereol.* 1987;67:41-47.
18. Molyneux P. The use of the stable free radical diphenylpicrylhydrazyl for estimating antioxidant activity. *Songklanakarin J Sci Technol.* 2004;26:211-219.
19. Orth DS. *Handbook of Cosmetic Microbiology.* New York: Marcel Dekker; 1997.
20. Pietta PG. Flavonoids as antioxidants. *J Nat Prod.* 2000;63:1035-1042.
21. Prior RL, Wu X, Schaich K. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *J Agric Food Chem.* 2005;53:4290-4302.

22. Rhein LD, Schlossman ML, editors. *Cosmetic Science and Technology: Theoretical Principles and Applications*.
23. Rosen MJ, Kunjappu JT. *Surfactants and Interfacial Phenomena*. 4th ed. Hoboken: Wiley; 2012.
24. Russell AD. Principles of antimicrobial activity. In: Block SS, editor. *Disinfection, Sterilization, and Preservation*. 5th ed. Philadelphia: Lippincott Williams & Wilkins; 2001. p. 31-54.
25. Sandle T. *Pharmaceutical Microbiology: Essentials for Quality Assurance and Quality Control*. Cambridge: Woodhead Publishing; 2016.
26. Tadros T. *Applied Surfactants: Principles and Applications*. Weinheim: Wiley-VCH; 2005.