Please check the examination details held	w before entering your candidate information
Candidate surname	Other names
Pearson Edexcel Intern	
Friday 7 June 2024	
Morning (Time: 2 hours)	Paper reference 4PM1/02
Further Pure Math PAPER 2	nematics
Calculators may be used.	Total Marks

## **Instructions**

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Without sufficient working, correct answers may be awarded no marks.
- Answer the questions in the spaces provided
  - there may be more space than you need.
- You must NOT write anything on the formulae page.
   Anything you write on the formulae page will gain NO credit.

## Information

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

## **Advice**

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.

Turn over ▶





## **International GCSE in Further Pure Mathematics Formulae sheet**

#### Mensuration

**Surface area of sphere** =  $4\pi r^2$ 

Curved surface area of cone =  $\pi r \times \text{slant height}$ 

Volume of sphere =  $\frac{4}{3}\pi r^3$ 

#### Series

#### **Arithmetic series**

Sum to *n* terms,  $S_n = \frac{n}{2} [2a + (n-1)d]$ 

## Geometric series

Sum to *n* terms, 
$$S_n = \frac{a(1-r^n)}{(1-r)}$$

Sum to infinity,  $S_{\infty} = \frac{a}{1-r} |r| < 1$ 

#### **Binomial series**

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{r!}x^r + \dots$$
 for  $|x| < 1, n \in \mathbb{Q}$ 

#### **Calculus**

## **Quotient rule (differentiation)**

$$\frac{\mathrm{d}}{\mathrm{d}x} \left( \frac{\mathrm{f}(x)}{\mathrm{g}(x)} \right) = \frac{\mathrm{f}'(x)\mathrm{g}(x) - \mathrm{f}(x)\mathrm{g}'(x)}{\left[\mathrm{g}(x)\right]^2}$$

## **Trigonometry**

#### Cosine rule

In triangle ABC:  $a^2 = b^2 + c^2 - 2bc \cos A$ 

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

$$\cos(A+B) = \cos A \cos B - \sin A \sin B$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

## Logarithms

$$\log_a x = \frac{\log_b x}{\log_b a}$$



## Answer all ELEVEN questions.

# Write your answers in the spaces provided.

## You must write down all the stages in your working.

1  $f(x) = 6x^3 - 13x^2 + ax - 10$  where a is a constant

Given that (3x - 2) is a factor of f(x)

(a) show that a = 21

**(2)** 

(b) Hence show algebraically that the curve y = f(x) has only one intersection with the x-axis.

(4)

(1)
(Total for Operation 1 is ( more)
(Total for Question 1 is 6 marks)



2	The quadratic equation $3x^2 - 5x + 1 = 0$ has roots $\alpha$ and $\beta$	
	Without solving the equation,	
	form a quadratic equation with integer coefficients, that has roots $\frac{\alpha}{2\beta}$ and $\frac{\beta}{2\alpha}$ (8)	








Figure 1

Figure 1 shows the sector AOB of a circle with centre O and radius 3r cm

A circle with radius r cm touches OA and OB and the arc AB

Angle *AOB* is  $\theta$  radians, where  $0 < \theta < \frac{\pi}{2}$ 

(a) Find the exact value of  $\theta$ 

(2)


The area of the region shown shaded in Figure 1 is  $8\pi$  cm<sup>2</sup>

(b) Find the value of r









# Figure 2

Figure 2 shows part of the curve with equation  $y = \frac{x^2}{3} - \frac{1}{2x}$  for -4 < x < 0By drawing a suitable straight line on the grid, obtain estimates, to one decimal place, of the roots of the equation  $4x^3 + 3x^2 - 36x - 6 = 0$  in the interval -4 < x < 0 (4)




_
5

	$y = e^{2x} \left( x^2 - 5x \right)$	
Show that $2e^{2x} = \frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y$		(7)








Figure 3

Figure 3 shows part of the curve C with equation  $y = \frac{1}{4x}$ , x > 0 and part of the curve S with equation  $y = 2x^2$ ,  $x \ge 0$ 

The curve C and the curve S intersect at the point A

(a) Find the coordinates of point A

(3)

The finite region R, shown shaded in Figure 3, bounded by the curve C, the curve S and the straight line y = 4 is rotated through 360° about the y-axis.

(b) Find, using algebraic integration, the exact volume of the solid formed.

**(7)** 




7 (a) Expand  $(1+2x^2)^{-\frac{3}{4}}$  in ascending powers of x up to and including the term in  $x^6$  Express each coefficient as an exact fraction in its lowest terms.

(3)

$$f(x) = \frac{(2+kx)}{(1+2x^2)^{\frac{3}{4}}}$$
 where  $k \neq 0$ 

(b) Obtain a series expansion for f(x) in ascending powers of x up to and including the term in  $x^5$ 

Give each coefficient in terms of k where appropriate.

**(2)** 

The coefficient of the term in  $x^5$  is fourteen times the coefficient of the term in  $x^2$ 

(c)	Find	the	value	of k	
-----	------	-----	-------	------	--

(2)

14



Question 7 continued	








Figure 4

Figure 4 shows a solid right triangular prism ABCDEF

The cross section of the prism is an isosceles triangle.

• 
$$\angle DEC = \angle AFB = 90^{\circ}$$

• 
$$AB = DC = x \text{ cm}$$

• 
$$AD = BC = FE = v \text{ cm}$$

• 
$$AF = BF = DE = CE$$

The triangular faces of the prism are vertical and the edges AD, BC and FE are horizontal.

The volume of the prism is 3.6 cm<sup>3</sup>

The total external surface area of the prism is  $S \text{ cm}^2$ 

(a) Show that S satisfies the equation

$$S = \frac{x^2}{2} + \frac{72\left(\sqrt{2} + 1\right)}{5x} \tag{4}$$

Given that x can vary,

(b) use calculus, to find to 3 significant figures, the value of x for which S is a minimum.


Justify that this value of x gives a minimum value of S

**(4)** 

(c) Hence find, to 2 significant figures, the minimum value of S

(2)





					Ś		
			C	7	₹		
	7	١	d		à	b	
					à		
			ŕ				
					ú		
					è		
					į		
					ķ	ø	
						ú	
		4	è	6	è		
					ģ		
					ś		
					è		
					2		
					N		
					É		
		d	輷		Ļ	Ħ	
		Š	Ž	9	Ż	9	
	3	à	6	ė	6	ė	
>	2	à	6	ė	6	ė	
	?	à	6	ė	6	ė	
	3	à	6	ė		ė	
	3	Ì					
	3	Ì					
	2	Ì					
	3	4					
	3	4					
	3	4					
	?	4					
	3						
	3						
	?						
	?						
	3						
	?						
>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>							

						3		
					è			
				É				
						i i		
					ś			
						ė		
>	Κ	$\geq$	ς	2	≤	2	٤	
>	S	?	S	į	Ś	2	٤	
>	8	3	ξ		Ź	2	٤	
>	5	?	5			2	٤	
	8	3	8			2	٤	
	5	3	S			2	٤	
>	5	3	{			2	٤	
	3	3	S	d				
>	3	3	S	d				
>	3	3	S	d				
	3	3	S	d				
	3	3	S	d				
		3						
	\ \ \	?						
	\ \ \	3						
	\ \ \	?						
		3						
		?						
>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>								
		?						
		3						
		?						
		3						

D				
	ļ			
	ø			
	έ			
à				
K		6	÷	é
13	٠	٠	٠	
э				
- 2		2	s	
а	ř	1		
٩				
κ,	ø	6	×	ú
		7		
Κ	d		۰	
K				
	ę			
	þ			
N				
. 2	2	2	2	5
М	è	쁴	#	d
À	8		7	d
Ν	₽	У	Р	
٦				
А	Ŕ			
Ν	ν	ч		
z				
	è	۷	2	
3	Ì	þ	è	ś

Question 8 continued	








Figure 5

Figure 5 shows a right triangular prism ABCDEF where ABCD is a rectangle.

$$AF = DE$$
  $BF = CE$   $AD = FE = BC$   $AB = DC = 24$  cm  
 $\angle ABF = \angle DCE = 45^{\circ}$   $\angle BAF = \angle CDE = 60^{\circ}$ 

Using a formula from page 2,

(a) show that 
$$\sin AFB = \frac{\sqrt{2} + \sqrt{6}}{4}$$

(3)

Without using a calculator,

(b) show that 
$$BF = 12(3\sqrt{2} - \sqrt{6})$$
 cm

(5)

The angle between the plane AEB and the plane ABCD is  $65^{\circ}$ 

(c) Find, in cm to 2 significant figures, the length of EF

(3)

(d) Find, in degrees to one decimal place, the size of the angle between the line *CF* and the plane *ABCD* 

**(4)** 



				S			
	à	í	è		á		
⟨	J	ľ	>	ζ	ď		
<		d	9	ę	5		
d	b	ø	9	ø	b		
2	9	Ġ	S	à	ø		
2		Ź	Š	7			
2	9	ė	9	Ì	ġ.		
2							
>	⋖	ž	S	Š	ď		
$\geq$	4	r	₹		ď		
S	ð	٩	ú	ø	۳		
S	2	2	2	2			
⟨	2	7	7	V	ř		
			2	۷	2		
	Ì	g		c	3		
<	ì	ø	ë	P	5		
2				,	۹		
$\rangle$	Þ	9		×			
$\rangle$	4		ņ	×	ĸ		
$\rangle$	4	þ	ę	þ	ij		
>	4	è	6				
S							
	d	P	9	ĺ	ľ		
ς	Į	ĸ,	F		P		
ς	2	2	2	S	è		
ς	Ž	S	2	ς			
◁		ζ	ā	P	?		
	)	ę	7		,		
Ś		ζ	>	3	è		
Ċ	ž	ę	9	d			
Ċ	)	é	è	í	ř		
	S	Ž	1	Ľ	S		
2	Š	Ź	3	Ź	Š		
2	S	Ź	Š	Ź	Š		
2	4	Ė	I	Š	á		
>		2	⋖	ž	ζ		
>	á	ò	۷	2			
>	Ś	j			ľ		
S	ð	S	Ź				
	Ź	Š	į	Š	ĺ		
	2	Š	2				
ζ	J	ŕ	1				
ζ	3	Š	Ź	S	2		
<	3	₫			ġ		
	j	ę	2	Č	Š		
<		Ż	>	<			
				?			
				?			
2				2			
2				2			
>	<	>	<	>			
>	Ċ	$\geq$	<	>			
>			<	>			
>			<	>			
5			2	S			
S				Ś			
S			2	Ś			
S				Ś			
S				Ś			
5			<	>			
$\rangle$			<	>			
$\rangle$			₹	>			
2	2	$\rangle$	<	2			
3			>	S			
3		3	>	5			
3			>	3			
3		?	>	3			
3		?		3			
		?	>				
	\ \ \ \						
					)		
					)		
					)		
					)		

	ς	2	۱
2	2	ä	
7	Ή	r	
	0		i
2		⋖	
	А	я	,
	2	4	i
	d		
2		S	i
/	Ч	r	
	a	в	ś
2	2	₹	۰
	۲	2	
	ð	۹	2
2	À	ú	
7	1	7	۰
	ď		
	2	₹	۱
۲	ς	ü	i
	ľ	3	
5	Э	ø	ø
	G	÷	ś
2	2	₹	
1	ì	2	
	ð	۹	ę
	Ч	۴	
	d	Ŀ	
5	3		
$\rangle$	þ	ą	į
ζ	ζ	4	ì
8		1	ì
3	۶	ď	
3			
3	S		
3	۶		
3	S		
3	S		
3	S		
?	S		
3	S		
3	S		
3	S		
	S		
?	S		
3	S		
3	S		
	S		
	S		
	S		
	S		
	S		
	S		
	S		
	S		
	S		

э				
'n				
d				
	è			
	2			
	r			
4				
	ġ			
	è			
			P	
'n				
'n		ø	۴	•

Question 9 continued	





10 The points A, B, C and D are the vertices of a quadrilateral such that

$$\overrightarrow{AB} = 3\mathbf{a} + 4\mathbf{b}$$
  $\overrightarrow{AC} = 7\mathbf{a} + 9\mathbf{b}$   $\overrightarrow{AD} = 4\mathbf{a} + 5\mathbf{b}$ 

(a) Show that ABCD is a parallelogram.

(3)

BC is extended to the point E such that BCE is a straight line.

Point F lies on CD such that CF : FD = 1 : 2

Given that A, F and E are collinear,

(b) find the vector  $\overrightarrow{AE}$  in the form  $X\mathbf{a} + Y\mathbf{b}$  where X and Y are rational numbers to be found.

(8)




$\times\!\!\times\!\!\!\times$	XXX.
$\times\!$	$\cdots$
$\times\!\!\!\sim$	
$\times \mathbb{Z}$	3KX
T 0 2	
80	
	₹XX.
882	¥XX.
$\sim$	
	₹XX.
C	XXX.
$\propto$	WXX.
$\circ$	m(XX
	mXX.
	₩XX
	<b>5</b> 000
$\diamond \circ \circ$	
	WXX.
$\sim$	
$\propto \times$	<u> </u>
ΧĤ	1XX.
$\infty$	XXX.
	=88
2	ĕXX.
$\times\!\!\times\!\!\times$	900.
X	
××	
$\times 2$	ZXX.
$\times \times$	
THIS ARE	
ΧV	
$\times \!\! \times \!\! \sim$	$\infty$
X	\$₩.
X	∞.
X	
$\times \nearrow$	
$\times$	
82	<b>*</b>
88	$\times\!\!\times\!\!\!\times$
$\times\!\!\times\!\!\times$	$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times$	$\times$
$\times\!\!\times\!\!\times$	$\times$
882	XX.
882	XXX.
	XXX.
$\sim\sim$	XXX.
$\times\!\!\times\!\!\times$	XXX.
$\times\!\!\times\!\!\times$	XXX.
$\sim \times$	$\times\!\!\times\!\!\times$
$\propto \propto$	$\times\!\!\times\!\!\times$
	$\times\!\!\times\!\!\times$
XX.	
$\times\!\!\times$	$\times\!\!\times\!\!\times$
$\otimes$	$\times\!\!\times\!\!\times$
$\otimes$	$\bowtie$
$\overset{\times}{\otimes}$	XX
t	
Ę	
t	
50	
001	
Ç	
Ç	) ) E
Ç	5
Ç	
Ç	
Ç	
Ç	
Ç	
Ç	
Ç	
Ç	
Ç	
CINCI WKI	
CINCI WKI	
Ç	
O NOT WALE	
O NOT WALE	
O WOJ WKIJEH	
ONCHWANTENIA	
O NOT WRITE IN THIS AR	
ONCHWANTENIA	

		1	41	r	7	3	
				ú	ń	6	
		0	G.	Р	₹	9	
	2		я	L		S	
2	۱	2	7	ø	8	ρ	í
				á			
					Ä		
					Ā		
					è		
					ì		
			١,	2	S	2	
	$\rangle$	<	þ	ę	è	è	į
>	⟨	S	ξ	į	È	Š	ĺ
>	2	ζ	}	į		į	ĺ
	3	5	}			į	
	2	3	}				
	3	5	3	1		Ź	
	3	3	3	1		Ź	
	3	3	3	1		Ź	
	3	3	3				
	3	3	3				
	3	3	3				
	?	3	3				
	3						
	?						
	3						
	3						
	\ \ \ \						
	3						
	\ \ \						

а	F	7	ę	
y	٩	ù	d	ø
S	à	S	n	š
ч	r	₹	5	
ď	9	ij	è	e
Ż				
)	۹	2		ļ
À	d		Z	
	Z	2	Z	3
Ý	۴	7	7	
Q	9	é	ġ	ø
4	2	4		
ė	7	7	7	
2		2		ú
ì		è	ę	۹
Z	Z	3		Ď
S	7	Ę	Ŕ	ĕ
4	è	6	è	í
K,	7	ä	٥	
å	S	2	2	Z
2	7	7	5	5
'n		è	ø	
S				2
٩	į	٩	P	
d	9)	₹	9	8
K	ì	2	_	ž
ż	Š	7	7	7
J	۳	7		ı
ì	1		í	û
				٦
'n	è	á	ú	
K				1
d	þ	۹	è	ă
Ġ	2	å	Ŀ	ú
Z	Ζ	Z	Ξ	=
Ž	S	7	Š	
J	Ľ	Z	f	
K.	7	۲		۹
4	ù	≤		
K	7			
d	ø	₹	٦	
μ	ă	ġ	ŕ	9
À	ά	ø	ķ	d
٦	á	à	á	í
Ч	ľ	1	P	
4	è	4	ì	
K	1		ž	ğ
J	۹	7	5	

Question 10 continued	





11 Using formulae from page 2, show that

(a) (i) 
$$\cos 2A = 2\cos^2 A - 1$$

(3)

(ii) 
$$\sin 2A = 2\sin A\cos A$$

(1)

(b) Show that 
$$\cos^3 A = \frac{\cos 3A + 3\cos A}{4}$$

**(4)** 

Hence, or otherwise,


(c) solve, giving exact values in terms of  $\pi$ 

$$8\cos^{3}\left(\frac{\theta}{2}\right) - 6\cos\left(\frac{\theta}{2}\right) - 1 = 0 \quad \text{for } 0 \leqslant \theta \leqslant 2\pi$$
(4)

(d) use algebraic integration to find the exact value of

$$\int_0^{\frac{\pi}{6}} \left( 4\cos^3\theta - \sin 2\theta \right) d\theta \tag{4}$$





Question 11 continued	
	(Total for Question 11 is 16 marks)
	TOTAL FOR PAPER IS 100 MARKS

