

LBL-TAGRISSO-OPENFDA

=====

Regulatory evidence card

Document ID: LBL-TAGRISSO-OPENFDA

Category: regulatory_submissions

Source entity: 04_regulatory_submissions/LBL-TAGRISSO-OPENFDA.json

Regulatory Identity

Document ID: LBL-TAGRISSO-OPENFDA

Document type: product_label

Source system: openfda_label

Application number: NDA208065

Brand name: TAGRISSO

Generic name: OSIMERTINIB

Sponsor/manufacturer: AstraZeneca Pharmaceuticals LP

Label Extract

Effective time: 20240925

Route: ORAL

Indications: 1 INDICATIONS AND USAGE TAGRISSO is a kinase inhibitor indicated for: ? adjuvant therapy after tumor resection in adult patients with non-small cell lung cancer (NSCLC) whose tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test. (1.1 , 2.2) ? the treatment of adult patients with locally advanced, unresectable (stage III) NSCLC whose disease has not progressed during or following concurrent or sequential platinum-based chemoradiation therapy and whose tumors have EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test. (1.2 , 2.2) ? the first-line treatment of adult patients with metastatic NSCLC whose tumors have EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test. (1.3 , 2.2) ? in combination with pemetrexed and platinum-based chemotherapy, the first-line treatment of adult patients with locally advanced or metastatic NSCLC whose tumors have EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test. (1.4 , 2.2) ? the treatment of adult patients with metastatic EGFR T790M mutation-positive NSCLC, a...

Warnings: 5 WARNINGS AND PRECAUTIONS ? Interstitial Lung Disease (ILD)/Pneumonitis : Monitor for new or worsening pulmonary symptoms indicative of ILD/pneumonitis. For patients receiving TAGRISSO who have not received recent definite platinum-based chemoradiation therapy, permanently discontinue TAGRISSO in patients diagnosed with ILD/Pneumonitis. For patients who received recent definitive platinum-based chemoradiation therapy with Grade 1 ILD/pneumonitis continue TAGRISSO or interrupt and restart, as appropriate. Permanently discontinue TAGRISSO in patients diagnosed with Grade ?2 ILD/pneumonitis. (2.5 , 5.1) ? QTc Interval Prolongation : Monitor electrocardiograms and electrolytes in patients who have a history or predisposition for QTc prolongation, or those who are taking medications that are known to prolong the QTc interval. Withhold, then restart at a reduced dose or permanently disco...

Provenance

Provenance: derived_from_raw_layer

Privacy posture:

Raw source trace:

00_raw/_corpus/json/regulatory/OPENFDA_LABEL_TAGRISSO/OPENFDA_LABEL_TAGRISSO.json