

F. Turtle Mission: Robot and the Earthquake

time limit per test: 3 seconds

memory limit per test: 256 megabytes

input: standard input

output: standard output

The world is a grid with n rows and m columns. The rows are numbered $0, 1, \dots, n - 1$, while the columns are numbered $0, 1, \dots, m - 1$. In this world, the columns are **cyclic** (i.e. the top and the bottom cells in each column are adjacent). The cell on the i -th row and the j -th column ($0 \leq i < n, 0 \leq j < m$) is denoted as (i, j) .

At time 0, the cell (i, j) (where $0 \leq i < n, 0 \leq j < m$) contains either a **rock** or **nothing**. The state of cell (i, j) can be described using the integer $a_{i,j}$:

- If $a_{i,j} = 1$, there is a rock at (i, j) .
- If $a_{i,j} = 0$, there is nothing at (i, j) .

As a result of aftershocks from the earthquake, the columns follow tectonic plate movements: each column moves cyclically **upwards** at a velocity of 1 cell per unit of time. Formally, for some $0 \leq i < n, 0 \leq j < m$, if (i, j) contains a rock at the moment, it will move from (i, j) to $(i - 1, j)$ (or to $(n - 1, j)$ if $i = 0$).

The robot called RT is initially positioned at $(0, 0)$. It has to go to $(n - 1, m - 1)$ to carry out an earthquake rescue operation (to the bottom rightmost cell). The earthquake doesn't change the position of the robot, they only change the position of rocks in the world.

Let RT's current position be (x, y) ($0 \leq x < n, 0 \leq y < m$), it can perform the following operations:

- Go one cell cyclically upwards, i.e. from (x, y) to $((x + n - 1) \bmod n, y)$ using 1 unit of time.
- Go one cell cyclically downwards, i.e. (x, y) to $((x + 1) \bmod n, y)$ using 1 unit of time.
- Go one cell to the right, i.e. (x, y) to $(x, y + 1)$ using 1 unit of time. (RT may perform this operation only if $y < m - 1$.)

Note that RT cannot go left using the operations nor can he stay at a position.

Unfortunately, RT will explode upon colliding with a rock. As such, when RT is at (x, y) and there is a rock at $((x + 1) \bmod n, y)$ or $((x + 2) \bmod n, y)$, RT cannot move down or it will be hit by the rock.

Codeforces Round 929 (Div. 3)

Finished

→ Practice?

Want to solve the contest problems after the official contest ends? Just register for practice and you will be able to submit solutions.

[Register for practice](#)

→ Virtual participation

Virtual contest is a way to take part in past contest, as close as possible to participation on time. It is supported only ICPC mode for virtual contests. If you've seen these problems, a virtual contest is not for you - solve these problems in the archive. If you just want to solve some problem from a contest, a virtual contest is not for you - solve this problem in the archive. Never use someone else's code, read the tutorials or communicate with other person during a virtual contest.

[Start virtual contest](#)



→ Problem tags

dfs and similar dp graphs
shortest paths


No tag edit access

→ Contest materials


- Announcement (en)



Similarly, if $y + 1 < m$ and there is a rock at $((x + 1) \bmod n, y + 1)$, RT cannot move right or it will be hit by the rock.



However, it is worth noting that if there is a rock at $(x \bmod n, y + 1)$ and $((x + 1) \bmod n, y)$, RT can still move right safely.



Find the minimum amount of time RT needs to reach $(n - 1, m - 1)$ without colliding with any rocks. If it is impossible to do so, output -1 .

Input

The first line of the input contains one integer t ($1 \leq t \leq 10^4$) — the number of test cases.

In each test case, the first line contains two integers n, m ($3 \leq n, m \leq 10^3$) — the size of the planet's boundaries.

Each of the next n lines contains m integers. The $(j + 1)$ -th integer on the $(i + 1)$ -th line ($0 \leq i < n, 0 \leq j < m$) is $a_{i,j}$ ($0 \leq a_{i,j} \leq 1$), which denotes whether or not there is a rock at (i, j) at time 0.

Additionally, it is guaranteed that $a_{0,0} = 0$, and $a_{i,m-1} = 0$ for $0 \leq i < n$. In other words,

there is no rock at RT's initial position as well as column $m - 1$.

The sum of $n \cdot m$ over all test cases does not exceed 10^6 .

Output

For each test case:

- If the destination can be reached without colliding with any rocks, output a single integer — the minimum amount of time RT needs to reach $(n - 1, m - 1)$.
- Otherwise, output -1 .

Examples

input	Copy
6 4 5 0 1 0 0 0 0 0 1 0 0 1 0 1 1 0 0 0 0 0 0 3 3 0 0 0 1 0 0 0 0 0 5 3 0 0 0 0 0 0 1 0 0 0 0 0 1 0 0 3 7 0 0 1 0 0 1 0 1 0 1 0 1 0 0 0 1 0 0 0 0 0 3 4 0 1 0 0 1 0 0 0 0 1 1 0 5 5 0 0 0 0 0 0 1 0 1 0 0 1 0 1 0 0 1 0 1 0 0 0 0 1 0	
output	Copy
7 3 3 8 -1 12	

input	Copy
6 3 3 0 0 0 0 0 0 0 0 0 4 3 0 1 0 1 0 0 0 1 0 1 0 0 4 3 0 1 0 0 1 0 0 1 0 0 1 0 3 3 0 0 0 1 1 0 0 0 0 3 3 0 1 0 0 0 0 0 1 0 5 5 0 0 0 0 0 0 1 1 0 0 0 1 1 0 0 0 0 0 0 0	

0 0 1 0 0


output

Copy

3
3
-1
-1
3
8

Note

Visual explanation of the first test case in the example:



Supported by

