

PRECISE summer school  
Warsaw

July 05, 2023

# hands-on cosmological simulations

session 2: cosmological integration and initial conditions

Sownak Bose  
&  
Shaun Brown

[sownak.bose@durham.ac.uk](mailto:sownak.bose@durham.ac.uk)

 @Swnk16



# choice of softening

$$\Phi(x) = -G \sum_{j=1}^N \frac{m_j}{\left[\left(\mathbf{x} - \mathbf{x}_j\right)^2 + \epsilon^2\right]^{1/2}}$$

gravitational softening,  $\epsilon$ , is introduced in N-body simulations to avoid large-angle scattering due to close encounters. higher-resolution simulations use smaller softening [and therefore also shorter timesteps]

simple choice:  $\epsilon = \alpha \cdot d = \alpha \frac{L_{\text{box}}}{N_p^{1/3}}$

$$\alpha = 0.01 - 0.05$$

mean inter-particle separation

# choice of softening

$$\Phi(x) = -G \sum_{j=1}^N \frac{m_j}{\left[\left(\mathbf{x} - \mathbf{x}_j\right)^2 + \epsilon^2\right]^{1/2}}$$

gravitational softening,  $\epsilon$ , is introduced in N-body simulations to **avoid large-angle scattering due to close encounters**. higher-resolution simulations use smaller softening [and therefore also shorter timesteps]

simple choice:  $\epsilon = \alpha \cdot d = \alpha \frac{L_{\text{box}}}{N_p^{1/3}}$

$$\alpha = 0.01 - 0.05$$

mean inter-particle separation

stochastic acceleration from close encounters

stricter choice:  $\frac{Gm_{\text{DM}}}{\epsilon^2} \lesssim \frac{GM_{200}}{r_{200}^2} \Rightarrow \epsilon \gtrsim \frac{r_{200}}{\sqrt{N_{200}}}$

minimum mean-field acceleration in DM halo

Power+ (2003)

# higher-order time integration

once forces on particles have even calculated, we need to propagate these forward in time and update their (1) **positions** and (2) **velocities**

kick

$$\mathbf{v}_j^{n+1/2} = \mathbf{v}_j^n + \frac{1}{2} \mathbf{a}_j^n \cdot \Delta t$$

drift

$$\mathbf{x}_j^{n+1} = \mathbf{x}_j^n + \mathbf{v}_j^{n+1/2} \cdot \Delta t$$

kick

$$\mathbf{v}_j^{n+1} = \mathbf{v}_j^{n+1/2} + \frac{1}{2} \mathbf{a}_j^{n+1} \cdot \Delta t$$

2nd order  
“leapfrog” method

much more stable  
“symplectic”  
[~energy is conserved]

# cosmological time integration

once forces on particles have even calculated, we need to propagate these forward in time and update their (1) **positions** and (2) **momenta**,  $\mathbf{p} = a\dot{\mathbf{x}}$

# cosmological time integration

once forces on particles have even calculated, we need to propagate these forward in time and update their (1) **positions** and (2) **momenta**,  $p = a\dot{x}$



# cosmological time integration

once forces on particles have even calculated, we need to propagate these forward in time and update their (1) **positions** and (2) **momenta**,  $\mathbf{p} = a\dot{\mathbf{x}}$

kick

$$\mathbf{p}_j(\tau_n + \Delta\tau) = \mathbf{p}_j(\tau_n) + \mathbf{a}_j(\tau_n) \int_{\tau_n}^{\tau_n + \Delta\tau} \frac{d\tau}{aH(a)}$$

cosmological scale factor

Hubble parameter

# cosmological time integration

once forces on particles have even calculated, we need to propagate these forward in time and update their (1) **positions** and (2) **momenta**,  $\mathbf{p} = a\dot{\mathbf{x}}$

kick

drift

$$\mathbf{p}_j(\tau_n + \Delta\tau) = \mathbf{p}_j(\tau_n) + \mathbf{a}_j(\tau_n) \int_{\tau_n}^{\tau_n + \Delta\tau} \frac{d\tau}{aH(a)}$$

$$\mathbf{x}_j(\tau_n + \Delta\tau) = \mathbf{x}_j(\tau_n) + \mathbf{p}_j(\tau_n) \int_{\tau_n}^{\tau_n + \Delta\tau} \frac{d\tau}{a^2 H(a)}$$

cosmological  
scale factor

Hubble  
parameter

# cosmological time integration

once forces on particles have even calculated, we need to propagate these forward in time and update their (1) **positions** and (2) **momenta**,  $\mathbf{p} = a\dot{\mathbf{x}}$

kick  
drift  
kick—drift—kick

$$\mathbf{p}_j(\tau_n + \Delta\tau) = \mathbf{p}_j(\tau_n) + \mathbf{a}_j(\tau_n) \int_{\tau_n}^{\tau_n + \Delta\tau} \frac{d\tau}{aH(a)}$$

$$\mathbf{x}_j(\tau_n + \Delta\tau) = \mathbf{x}_j(\tau_n) + \mathbf{p}_j(\tau_n) \int_{\tau_n}^{\tau_n + \Delta\tau} \frac{d\tau}{a^2 H(a)}$$

$$E(\Delta\tau) = K\left(\frac{\Delta\tau}{2}\right) \circ D(\Delta\tau) \circ K\left(\frac{\Delta\tau}{2}\right)$$

cosmological  
scale factor

Hubble  
parameter

# a note on the Hubble parameter

we define the Hubble  
parameter as  $H(t) \equiv \frac{\dot{a}(t)}{a(t)}$ .  
denoting the present-day  
value as

$$H_0 \equiv H(t = t_{\text{now}}) = 100h \text{ km s}^{-1} \text{ Mpc}^{-1}$$

# a note on the Hubble parameter

we define the Hubble parameter as  $H(t) \equiv \frac{\dot{a}(t)}{a(t)}$ . denoting the present-day value as

$$H_0 \equiv H(t = t_{\text{now}}) = 100h \text{ km s}^{-1} \text{ Mpc}^{-1}$$

$$H(a) = H_0 E(a)$$

$$E(a) = \sqrt{\Omega_m a^{-3} + \Omega_r a^{-4} + \Omega_k a^{-2} + \Omega_\Lambda \exp(3\tilde{w}(a))}$$

$$\tilde{w}(a) = (a - 1) w_a - (1 + w_0 + w_a) \log(a)$$

$$w(a) \equiv w_0 + w_a (1 - a)$$

# a note on the Hubble parameter

we define the Hubble parameter as  $H(t) \equiv \frac{\dot{a}(t)}{a(t)}$ . denoting the present-day value as

$$H_0 \equiv H(t = t_{\text{now}}) = 100h \text{ km s}^{-1} \text{ Mpc}^{-1}$$

$$H(a) = H_0 E(a)$$

$$E(a) = \sqrt{\Omega_m a^{-3} + \Omega_r a^{-4} + \Omega_k a^{-2} + \Omega_\Lambda \exp(3\tilde{w}(a))}$$



$$\tilde{w}(a) = (a - 1) w_a - (1 + w_0 + w_a) \log(a)$$

$$w(a) \equiv w_0 + w_a (1 - a)$$


the cosmological model is therefore **fully specified** through  $\Omega_m, \Omega_r, \Omega_k, \Omega_\Lambda, h, w_0$  and  $w_a$


**setting up initial conditions**

measurement of the **cosmic microwave background radiation** gives us *precise constraints* on cosmological initial conditions




## homogeneous & isotropic ICs







**these tiny fluctuations  
will grow into non-linear  
structures (e.g. filaments,  
walls, haloes, clusters)  
through gravitational  
instability**

**perturbed ICs**





## glass initial conditions

start with a Poisson [random] particle distribution and evolve this with the sign of gravity “inverted” [repulsive force] until particles “freeze” in comoving coordinates




helps wash out the memory of the initial lattice

**start with a Poisson [random]  
particle distribution and  
evolve this with the sign of  
gravity “inverted” [repulsive  
force] until particles “freeze”  
in comoving coordinates**




**helps wash out the  
memory of the initial  
lattice**


**goal:** create a **Gaussian random distribution** that is statistically consistent with the fluctuations measured in the CMB




**goal:** create a **Gaussian random distribution** that is statistically consistent with the fluctuations measured in the CMB



**goal:** create a **Gaussian random distribution** that is statistically consistent with the fluctuations measured in the CMB




**goal:** create a **Gaussian random distribution** that is statistically consistent with the fluctuations measured in the CMB



$$\langle \delta(\mathbf{x}) \delta(\mathbf{x}') \rangle = \xi(r)$$

correlation function

**goal:** create a **Gaussian random distribution** that is statistically consistent with the fluctuations measured in the CMB



$$\langle \delta(\mathbf{x}) \delta(\mathbf{x}') \rangle = \xi(r)$$

$$\langle \delta(\mathbf{k}) \delta^*(\mathbf{k}') \rangle = P(k)$$


correlation function

---

power spectrum

---

**goal:** create a **Gaussian random distribution** that is statistically consistent with the fluctuations measured in the CMB



$$\langle \delta(\mathbf{x}) \delta(\mathbf{x}') \rangle = \xi(r)$$

**correlation function**

$$\langle \delta(\mathbf{k}) \delta^*(\mathbf{k}') \rangle = P(k)$$

**power spectrum**

the statistical properties of a Gaussian random field are characterised fully by  $P(k)$

**fluctuation amplitudes**  $\propto \sqrt{P(k)}$

**phases**  $\in [0, 2\pi]$

# the primordial power spectrum

inflation makes a specific prediction for the primordial density field that is a power-law:

$$P(k) \propto k^{n_s}$$

where  $n_s = -1$  corresponds to a **scale-invariant** power spectrum

# the primordial power spectrum

inflation makes a specific prediction for the primordial density field that is a power-law:

$$P(k) \propto k^{n_s}$$

where  $n_s = -1$  corresponds to a **scale-invariant** power spectrum

depending on what the Universe is made of and how fluctuations grow in time and space, the inflationary power spectrum is modified by a **transfer function**

$$P(k, t) = A k^{n_s}$$

# the primordial power spectrum

inflation makes a specific prediction for the primordial density field that is a power-law:

$$P(k) \propto k^{n_s}$$


where  $n_s = -1$  corresponds to a **scale-invariant** power spectrum

depending on what the Universe is made of and how fluctuations grow in time and space, the inflationary power spectrum is modified by a **transfer function**

$$P(k, t) = A k^{n_s} |T(k, t)|^2$$


computed by public codes  
like CAMB, CLASS etc.

**Stücker+  
(2018)**



one way to model different DM candidates is to change the  $P(k)$  used to generate ICs

Stücker+  
(2018)



one way to model different DM candidates is to change the  $P(k)$  used to generate ICs

now, we more or less have everything we need: we just need to **perturb the initial particle configuration according to these fluctuations**. the position of a particle initially at coordinate  $\mathbf{q}$  after some time  $t$  is given by:

$$\mathbf{x}(t) = \mathbf{q} +$$

now, we more or less have everything we need: we just need to **perturb the initial particle configuration according to these fluctuations**. the position of a particle initially at coordinate  $\mathbf{q}$  after some time  $t$  is given by:

$$\mathbf{x}(t) = \mathbf{q} + \mathbf{S}(\mathbf{q})$$

now, we more or less have everything we need: we just need to **perturb the initial particle configuration according to these fluctuations**. the position of a particle initially at coordinate  $\mathbf{q}$  after some time  $t$  is given by:

$$\mathbf{x}(t) = \mathbf{q} + D(t) \cdot \mathbf{S}(\mathbf{q})$$


now, we more or less have everything we need: we just need to **perturb the initial particle configuration according to these fluctuations**. the position of a particle initially at coordinate  $\mathbf{q}$  after some time  $t$  is given by:

$$\mathbf{x}(t) = \mathbf{q} + D(t) \cdot \mathbf{S}(\mathbf{q})$$


**Zel'dovich approximation**

---


now, we more or less have everything we need: we just need to **perturb the initial particle configuration according to these fluctuations**. the position of a particle initially at coordinate  $\mathbf{q}$  after some time  $t$  is given by:




now, we more or less have everything we need: we just need to **perturb the initial particle configuration according to these fluctuations**. the position of a particle initially at coordinate  $\mathbf{q}$  after some time  $t$  is given by:



$\mathbf{S}(\mathbf{q}) \propto -\nabla\Phi(\mathbf{q})$ , and we know that  $\Phi$  is related to  $\delta(\mathbf{k})$  through **Poisson's equation**



credit: [Johan Hidding](#)



credit: [Johan Hidding](#)

# the non-linear density field

while the initial conditions of our cosmos may be approximated as a Gaussian random field, the late-time, non-linear density is anything but Gaussian! gravitational instability naturally transitions the matter field away from Gaussianity

$$\langle \delta(\mathbf{k}, z) \delta^*(\mathbf{k}', z) \rangle = P(k, z)$$

# the non-linear density field

while the initial conditions of our cosmos may be approximated as a Gaussian random field, the late-time, non-linear density is anything but Gaussian! gravitational instability naturally transitions the matter field away from Gaussianity

$$\langle \delta(\mathbf{k}, z) \delta^*(\mathbf{k}', z) \rangle = P(k, z)$$

given a density field,  $\delta(\mathbf{x})$ , you can also quantify the degree of non-Gaussianity by computing moments:

size of the grid  
used to smooth  
the density field

$$\langle \delta^n \rangle = \frac{1}{N_g} \sum_i^{N_g} (\delta^i - \langle \delta \rangle)^n$$

- $n = 0$  (mean)
- $n = 2$  (variance)
- $n = 3$  (skewness)
- $n = 4$  (kurtosis)

**lecture notes:**

[https://github.com/sownakbose/PRECISE Summer School Sims](https://github.com/sownakbose/PRECISE_Summer_School_Sims)

**jupyter notebooks:**

[https://github.com/Shaun-T-Brown/Summer school](https://github.com/Shaun-T-Brown/Summer_school)