

Health Insurance and Fertility: Evidence from the Advent of Blue Cross

Soyoung Han *

October 7, 2026

[Please click here for the latest version](#)

Abstract

After decades of decline, U.S. fertility began to rise in the mid-1930s, well before the postwar Baby Boom. At the same time, employer-sponsored health insurance was expanding nationwide for the first time. Did the emergence of employer-sponsored hospital insurance contribute to this reversal? I construct a new dataset tracing the geographic rollout and enrollment of Blue Cross hospital plans and link it to county-level hospital births and vital statistics. Using both stacked difference-in-differences and an instrumental variables designs, I find that Blue Cross entry increased insurance enrollment and hospital childbirth. Hospital births rose by 0.37 per 100 women aged 15–44, with little evidence of a corresponding decline in non-hospital births. Total fertility also increased, although the estimate is imprecise. The effects are larger in areas with higher baseline medical costs, suggesting that the financial cost of childbirth played an important role. Meanwhile, infant and maternal mortality show little change. Blue Cross contributed meaningfully to the mid-1930s fertility decline reversal in the United States, highlighting how health-care policy can shape demographic outcomes beyond its direct effects on medical care.

*University of Michigan. Email: soyoungh@umich.edu

1 Introduction

U.S. fertility rates declined for much of the nineteenth and early twentieth centuries. Yet this long-run decline came to an end in mid-1930s before the postwar Baby Boom [Bailey and Hershbein, 2018]. At the same time, another major institutional development was occurring in the United States—the emergence of employer-sponsored prepaid hospital insurance (Figure 1). During the Great Depression, Baylor University Hospital pioneered a prepaid hospital plan for groups of employees, an arrangement that became an early model for Blue Cross hospital plans. These plans became one of the earliest widespread forms of employer-sponsored prepaid hospital coverage in the United States. The coincidence of these developments raises a natural question: did the expansion of Blue Cross hospital plan contribute to the reversal of fertility decline that began in the mid-1930s?

By the 1930s, childbirth was already shifting rapidly from homes to hospitals, with the share of births occurring in hospitals rising from less than five percent in 1900 to roughly 50 percent by 1939 [Leavitt, 1986; Wertz and Wertz, 1979]. This transition had begun well before the emergence of Blue Cross. However, hospital delivery represented a substantial medical expense for many families. Hospital insurance changed not only how households financed medical care, but potentially the economic cost of childbirth and having a child. Its effect on fertility is nevertheless theoretically ambiguous. Insurance could increase fertility by reducing the financial burden of childbirth. Greater access to medical care could also improve maternal health, potentially encouraging childbearing by reducing its health risks or discouraging it by increasing women’s opportunity costs. Improvements in infant survival could further encourage fertility by increasing the likelihood that a birth results in a surviving child. Despite the central role that employer-sponsored health insurance eventually came to play in the U.S. healthcare system, relatively little is known about how its initial large-scale expansion affected childbearing. Studying the emergence of Blue Cross provides an opportunity to examine the broader consequences of health insurance while also evaluating an important institutional change that occurred alongside the reversal of the long-run decline in U.S. fertility.

To study this question, I construct a new dataset tracing the geographic expansion of Blue Cross hospital plans across the United States. I digitize historical records on each plan’s first year of

enrollment, geographic coverage, and subsequent plan-level enrollment. I link these data to county-level measures of hospital births from the American Medical Association’s *Hospital Service in the United States* and to birth registration records, which allow me to quantify hospital, non-hospital, and total births at the county-year level. These data allow me to examine the impact of Blue Cross on hospital delivery and total fertility.

My empirical strategy exploits variation in the timing and geographic rollout of Blue Cross plans. I first estimate the reduced-form effects of Blue Cross entry using weighted stacked difference-in-differences approach [Wing et al., 2024]. I then use the introduction of Blue Cross as an instrument for insurance enrollment to estimate the effect of enrollment on hospital childbirth and fertility.

I find that the introduction of Blue Cross substantially increased insurance enrollment share and the use of hospital childbirth. Plan-level enrollment rose by approximately 8.8 percentage points on average after introduction, while the hospital birth rate increased by 0.37 births per 100 women aged 15–44, relative to a pre-treatment mean of 0.91. However, I do not find evidence of an impact on the non-hospital birth rate, suggesting that the increase in hospital births was not driven solely by a shift in the location of deliveries that would otherwise have occurred outside hospitals. The estimated effect on the total birth rate is also positive: the birth rate increases by approximately 0.31 births per 100 women, although I cannot reject the null hypothesis that the effect on total births is zero. Instrumental variables estimates, which use the introduction of Blue Cross plans as an instrument for insurance enrollment, yield qualitatively similar results: greater Blue Cross enrollment significantly increases hospital births, while the estimated effects on total fertility are positive but is imprecisely estimated.

I then study the potential mechanisms through which the availability of health insurance affected birth rates, considering financial, maternal health, and infant survival channels. The evidence points to the alleviation of the financial costs of childbirth as an important factor linking Blue Cross availability to higher fertility. First, I examine hospitals with high and low baseline medical costs. The effects on hospital births are significantly larger in counties with high baseline medical costs, and the estimated fertility response is also larger in these areas, although the difference in total

fertility is imprecisely estimated. In contrast, I find little evidence that Blue Cross improved infant or maternal mortality. The results show that Blue Cross substantially increased the use of hospital delivery and provide suggestive evidence that reducing the financial burden of hospital childbirth also increased fertility.

A back-of-the-envelope calculation suggests that the expansion of Blue Cross could account for roughly 13 percent of the observed increase in the U.S. birth rate over this period. Although this estimate is imprecise, it suggests that the emergence of employer-sponsored hospital insurance may have contributed meaningfully to the rise in fertility that began in the mid-1930s.

By focusing on the turning point when fertility began to rise after decades of decline, I first contribute to the literature on the reversal of U.S. fertility in the mid-1930s and the origins of the subsequent Baby Boom, a period of elevated fertility in the United States from 1946 to the mid-1960s. A wide range of explanations has been proposed for this rise in fertility, including the relative income hypothesis, which argues that individuals who grew up during the Great Depression had relatively low material aspirations and were therefore more inclined to have children as their incomes rose during the prosperous postwar period [Easterlin, 1966; Hill, 2015], improvements in maternal health [Albanesi and Olivetti, 2014], the diffusion of household technologies [Greenwood et al., 2005; Bailey and Collins, 2011; Lewis, 2018], the economic disruption of the Great Depression [Bellou et al., 2025], the effects of World War II [Doepke et al., 2015; Brodeur and Kattan, 2022], the expansion of modern U.S. mortgage markets [Dettling and Kearney, 2025], and the expansion of labor unions following the 1935 National Labor Relations Act, which improved male workers' earnings and job security while reducing women's labor market opportunities [Downes, 2024]. While much of this literature focuses on the Baby Boom more broadly, this paper focuses on the earlier turning point when fertility began to rise in the mid-1930s. I identify the role of a major institutional development occurring during this period—the emergence of employer-sponsored hospital insurance—and show that the expansion of Blue Cross substantially increased hospital childbirth and also contributed to the rise in fertility.

Second, I contribute to the literature on the effects of health insurance on fertility by providing new

historical evidence from the first nationwide employer-sponsored health insurance system. Existing evidence is mixed, with most studies focusing on modern policy changes and relatively limited evidence from historical settings. Several papers exploit quasi-experimental variation from Medicaid eligibility expansions for pregnant women and children and document mixed effects on fertility outcomes [DeLeire et al., 2011; Zavodny and Bitler, 2010]. Using experimental evidence, Leibowitz [1990] studies a setting in which medical costs were directly reduced and finds a positive effect on fertility. Historical evidence is more limited. Troland and Figinski [2019] examine an expansion of health insurance coverage for unionized workers in the mining industry and find negative effects on fertility. A related literature studies insurance mandates for assisted reproductive technologies. These studies find that reducing the financial burden associated with infertility treatments increases fertility, particularly among women of advanced maternal age [Bitler and Schmidt, 2012; Schmidt, 2007]. My paper contributes to this literature by providing new historical evidence on the relationship between health insurance and fertility, during the early expansion of employer-sponsored hospital insurance in the United States. By studying Blue Cross hospital plans, I examine a broad form of insurance that covered hospital childbirth and therefore directly altered the financial cost of delivery.

The remainder of the paper is organized as follows. I begin in Section 2 with the historical background on the development and expansion of Blue Cross hospital plans and the broader transition toward hospital childbirth. Section 3 then develops the conceptual framework, highlighting the channels through which hospital insurance may affect fertility. The data sources and construction of the main variables are described in Section 4, followed by the empirical strategy in Section 5. Section 6 presents the main empirical results and examines the mechanisms through which Blue Cross may have affected hospital childbirth and fertility. Section 7 discusses hospital capacity and the absence of detectable mortality improvements despite greater hospital use. Section 8 concludes.

2 Historical Background

2.1 Childbirth Practices

Childbirth practices in the United States underwent a profound transformation over the nineteenth and early twentieth centuries. While childbirth was historically a female-centered event attended by women in the community and often a midwife, the presence of male physicians at home births began increasing in the late eighteenth century. [Leavitt, 1986].¹

Despite the growing medicalization of childbirth, hospitals initially played only a limited role. Before the turn of the twentieth century, hospitals primarily served as charitable institutions for poor or unmarried women who lacked the resources to have a physician or midwife attend a home birth. This pattern changed rapidly in the early twentieth century as hospital delivery became increasingly common, rising from less than five percent of births in 1900 to roughly 50 percent (and nearly 75 percent in urban areas) by 1939 [Leavitt, 1986; Wertz and Wertz, 1979].

From the perspective of healthcare providers, hospitals offered important advantages. Physicians could care for patients in a controlled hospital environment, eliminating the need to travel to patients' homes, transport medical equipment, and remain there through lengthy labors. Concentrating childbirth within hospitals made the provision of obstetric care more efficient. In addition, maternity hospitals became important training grounds for medical students and young physicians, providing a steady flow of patients for clinical observation and instruction [Leavitt, 1986; Wertz and Wertz, 1979].

Hospital delivery also became increasingly attractive from the perspective of expectant mothers. Prenatal and postnatal care eventually came to be viewed as increasingly important for managing pregnancy-related risks. Hospitals adopted practices such as routine urine testing to detect conditions like eclampsia at earlier stages. Influential figures such as John Whitridge Williams—often

¹The decline in midwife-attended births continued into the twentieth century. Table 1 shows that the share of births attended by midwives in New Jersey declined from over 40 percent in 1919 to 20 percent in 1926. In New York, the share declined from 16 percent in 1916 to 8 percent in 1924. In Michigan, the share declined from 6.9 percent in 1921 to 4.4 percent in 1924 [U.S. Department of Labor, Children's Bureau, 1926]. In Richmond, Virginia, the share declined from 41 percent in 1907 to 18 percent in 1922 [vir, 1923].

regarded as the founder of academic obstetrics—promoted hospital delivery as a safer and more modern form of childbirth, reinforcing public confidence in institutional care. The introduction of innovations such as Twilight Sleep, which promised to alleviate the pain of childbirth, as well as access to emergency interventions such as C-sections, further increased the appeal of hospital delivery [Wertz and Wertz, 1979].

Despite their growing popularity, hospitals did not yet have an established safety advantage. Medical practice was still evolving, and physicians frequently experimented with new techniques, sometimes drawing criticism for excessive or unnecessary intervention [Leavitt, 1986]. Contemporaneous reports from the U.S. Children’s Bureau illustrate the uncertain relationship between hospital delivery and maternal safety. In a report by Children’s Bureau, 57 percent of maternal deaths were classified as hospital cases, while the remaining deaths were classified as non-hospital cases [U.S. Children’s Bureau, 1934]. Although these figures do not provide a direct comparison of mortality risk between hospital and non-hospital births, they illustrate that a substantial share of maternal deaths continued to occur in hospitals during this period. The Children’s Bureau noted that “In normal cases delivery can be safely carried out in the home” [U.S. Children’s Bureau, 1938].

At the same time, hospital childbirth was expensive. In the early twentieth century, postpartum hospital stays were substantially longer than they are today, often lasting up to two weeks, allowing charges to accumulate over an extended period [Temkin, 1999]. Contemporary estimates illustrate the magnitude of these expenses. In six New York counties in 1935–36, estimates suggest that hospital costs associated with delivery ranged from approximately \$40 to \$59, with the estimated total cost of a hospital delivery ranging from approximately \$65–\$70. In Berkeley, California, total obstetric care in 1928 was estimated at approximately \$200–\$215, including an average physician fee of approximately \$92 [U.S. Children’s Bureau, 1941; U.S. Bureau of Labor Statistics, 1930]. Figure 2 provides examples of hospital receipts issued following childbirth in the late 1920s and 1930s, highlighting the financial burden associated with hospital delivery.

These expenses were substantial relative to household resources at the time: median annual household income during the mid-1930s was only approximately \$1,000–\$1,250 [National Resources Com-

mittee, 1938]. On a monthly basis, this amounted to about \$83–\$104, highlighting the potentially substantial burden of childbirth-related expenses when they came due around the time of delivery. A hospital delivery could therefore absorb a meaningful share of a household’s annual resources, even though childbirth itself was generally anticipated in advance. Contemporary accounts suggest that these costs were salient for fertility decisions. In a 1923 letter published in *Ladies’ Home Journal*, one reader wrote: “The reason for the decrease in American babies is the high cost of obstetrics... Our first baby cost: for layette – with strictest economy – \$25; ten days in a maternity hospital, \$35; dressings and laundry at the hospital, \$5; doctor’s charge, \$25; anesthetic, \$5; total, \$95. My mother had seven babies and I doubt if they all cost her \$100 put together” [Wertz and Wertz, 1979]. The account illustrates both the growing use of medical and hospital-based childbirth and the financial burden that such care could impose on families.

By the time Blue Cross hospital plans emerged, the transition toward hospital-based childbirth was already well underway. Hospital delivery had become increasingly desirable to both patients and providers even though its health advantages were not yet firmly established. At the same time, the substantial cost of institutional childbirth made hospital delivery a significant financial burden for households. This combination of growing appeal and high cost created a setting in which health insurance could play an important role in shaping childbirth decisions.

2.2 Early Health Insurance Initiatives

Despite the growing importance and cost of hospital care, health insurance remained extremely limited in the United States. Prior to the emergence of Blue Cross, efforts to provide health insurance were fragmented and highly localized, often limited to particular cities, employers, or industries, and failed to achieve broad coverage, leaving most households exposed to substantial medical cost risks. One barrier to expansion was the reluctance of private insurance companies to offer health insurance, which they viewed as particularly risky due to adverse selection [Morrissey, 2013]. A second barrier to the expansion came from organized medicine. The American Medical Association (AMA) actively opposed alternative insurance models, including cooperatives and prepaid group practice arrangements, which it viewed as threats to physician autonomy and the prevailing fee-for-

service model, and leveraged its influence over physician licensing and lobbying efforts to restrict their development [Starr, 2017].

In parallel with these private initiatives, repeated efforts were made to establish publicly organized health insurance. In 1912, Teddy Roosevelt endorsed social insurance against sickness as part of the Progressive Party platform. In 1915, the American Association for Labor Legislation developed and promoted a model bill for compulsory health insurance. Although the proposal initially received support from the AMA, political support subsequently weakened amid opposition from organized labor and other groups, the AMA's eventual reversal of its position, and growing anti-German and anti-socialist sentiment tied to World War I [Institute of Medicine, 1993]. Efforts to establish compulsory health insurance continued at the state level, including several proposals during the 1930s, but these initiatives also failed to pass state legislatures [Reed, 1939]. Thus, by the beginning of the 1930s, neither private insurance markets nor compulsory public insurance had produced a system capable of providing health coverage on a broad scale.

2.3 Blue Cross Hospital Plans

Against this backdrop, a new model of hospital insurance emerged that would ultimately achieve broader coverage. The origins of this model can be traced to a 1929 initiative at Baylor University Hospital in Dallas, Texas, which was experiencing financial difficulties as declining patient incomes reduced demand for hospital care. Dr. Justin Ford Kimball developed an experimental hospital plan that allowed schoolteachers to make small monthly payments in exchange for hospital services when needed, attracting more than 1,500 enrollees. Its early success drew interest from other employee groups and hospitals in Dallas and quickly gained attention nationwide, contributing to the emergence of the Blue Cross Plan Movement in 1932. In 1934, the American Hospital Association endorsed the development of group hospitalization plans but favored community-wide plans over single-hospital plans, allowing patients to choose among participating hospitals rather than limiting them to a single institution [Cunningham III and Cunningham Jr., 1997]. The subsequent expansion of Blue Cross was rapid. Figure 1 shows that Blue Cross enrollment increased from fewer than one million enrollees in 1935 to more than 24 million by 1946 [Reed, 1947].

Blue Cross was able to achieve this scale in part because its institutional structure addressed several of the barriers that had limited earlier insurance arrangements. First, enrollment was organized primarily through employment groups rather than through the individual market, which facilitated risk pooling and reduced concerns about adverse selection. Second, participation by employers and employees remained voluntary, avoiding many of the political objections directed toward compulsory insurance. Finally, Blue Cross restricted coverage primarily to hospital services rather than physician fees, reducing direct conflict with organized medicine. Monthly premiums typically ranged from \$0.60 to \$3.60, depending on the plan and type of contract [Reed, 1947]. These institutional features allowed Blue Cross to become the first viable system of large-scale prepaid hospital insurance in the United States.

While the core benefits offered across Blue Cross plans were broadly similar—typically including hospital room and board, nursing services, and use of the operating room—coverage of additional services varied across plans, in part because of disagreements over whether particular services constituted hospital expenses or physician services. For example, some plans covered anesthesia and X-rays, while others did not [Reed, 1947]. This distinction between hospital and physician services was institutionally important because Blue Cross initially insured hospital expenses rather than the full cost of medical care. Resistance from organized medicine delayed the development of Blue Shield plans, which provided coverage for physician services and emerged later than Blue Cross plans (Figure 3).

One particularly important application of this hospital coverage—and the one most relevant to this study—was maternity care. Some early plans were hesitant to provide maternity benefits because the need for maternity care could often be anticipated, raising concerns that households might enroll when they expected to use maternity services. However, the experience of plans that successfully offered maternity coverage contributed to its broader adoption, and maternity benefits eventually became standard across Blue Cross plans [Cunningham III and Cunningham Jr., 1997]. Maternity coverage was generally available through couple and family plans [Reed, 1947]. Depending on the plan, maternity benefits covered hospital stays of approximately 10 to 30 days at full benefits, reflecting the substantially longer postpartum hospital stays common during this period [Temkin,

1999]. To limit adverse selection, plans commonly imposed waiting periods of approximately 9 to 12 months before maternity benefits became available although these waiting periods could sometimes be waived for employer groups with sufficiently high participation [Reed, 1947].

Maternity hospitalization became one of the most important uses of hospital insurance. Table 2 shows that maternity accounted for the largest share of cases covered by the New York plan during the 1940s. A similar pattern appears in the New Jersey plan: as shown in Table 3, maternity accounted for the largest share of cases, total days of hospital care, and total hospital expenses in 1943 [Reed, 1947]. The prominence of maternity care in Blue Cross utilization highlights the importance of childbirth-related hospital expenses to the early development of hospital insurance and raises the question of whether insurance coverage for these expenses affected households' fertility decisions.

3 Conceptual Framework

The historical evidence above shows that maternity care was an important component of early Blue Cross coverage and that hospital childbirth represented a substantial expense for households during this period [United States Department of Labor, Children's Bureau, 1938; U.S. Children's Bureau, 1941; U.S. Bureau of Labor Statistics, 1930]. The expansion of Blue Cross therefore changed the costs and conditions associated with childbearing. Whether these changes affected fertility depends on how insurance altered households' incentives to attempt conception and, through this behavioral response, the probability of a realized live birth.

I use a simple conceptual framework to illustrate three channels through which health insurance may affect fertility: the financial cost of childbirth, maternal health, and infant survival. Households choose whether to attempt conception, while a live birth is realized probabilistically conditional on attempting. In making this decision, households take into account the financial and time costs of childbearing, the health cost to the mother, and the expected benefit of an additional surviving child. Insurance can affect the incentive to attempt conception by changing the costs and benefits associated with a realized birth. I describe the intuition behind each channel here and present the

formal framework in Appendix B.

First, insurance can affect fertility by reducing the financial burden of childbirth. As noted above, hospital delivery represented a substantial household expense during this period.² Blue Cross reduced the amount that insured households paid out of pocket for covered hospital care at the time of delivery. More generally, existing work has emphasized that insurance coverage for pregnancy can affect fertility by reducing the financial cost of having a child [Leibowitz, 1990]. A lower out-of-pocket expense increases the resources available to the household in the event of a live birth and therefore raises the value of the live-birth state relative to the no-live-birth state. This increases the incentive to attempt conception when the financial benefit of coverage is sufficiently large relative to the insurance premium.

The financial channel also generates a useful prediction for heterogeneity. For a given coverage rate, insurance produces a larger dollar reduction in out-of-pocket childbirth expenses where baseline childbirth costs are higher. The model therefore predicts a larger effect of insurance on the incentive to attempt conception in higher-cost markets. If differences across markets in the share of households near the conception-attempt threshold do not offset this larger incentive effect, realized births should also respond more strongly in higher-cost markets.

Second, insurance may affect fertility through maternal health. Improved access to hospital care can improve women’s health, with potentially opposing effects on the conception decision. Better maternal health can reduce the health cost associated with pregnancy and childbirth, increasing the value of a live birth and therefore the incentive to attempt conception [Jayachandran and Lleras-Muney, 2009; Albanesi and Olivetti, 2014]. At the same time, better health can increase women’s ability to engage in market work [Albanesi and Olivetti, 2016; Stephens and Toohey, 2022; Gilleskie et al., 2017], thereby raising the opportunity cost of childbearing. The net effect of the maternal-health channel on conception attempts and realized fertility is therefore theoretically ambiguous. To assess whether maternal health could have been an operative channel, I examine whether Blue Cross entry affected maternal health.

²Contemporary accounts characterized maternity hospitalization as a relatively large and unpredictable expense for the family budget [United States Department of Labor, Children’s Bureau, 1938].

Finally, insurance may affect fertility through infant survival. If access to insured hospital care increases the probability that a live-born infant survives, it raises the expected value of a live birth [Becker and Barro, 1988; Cigno, 1998; Guinnane, 2011]. Within the framework developed here, in which households value an additional surviving child, improved infant survival raises the value of the live-birth state relative to the no-live-birth state and increases the incentive to attempt conception. I consider whether improvements in infant survival provide an alternative explanation for the fertility response to Blue Cross by examining whether infant survival improved following Blue Cross entry.

These channels provide guidance for the empirical analysis. In particular, heterogeneity in the effects of Blue Cross by baseline medical costs is consistent with the financial channel if fertility responds more strongly in higher-cost areas, while effects on maternal and infant mortality provide evidence on whether improvements in maternal health or infant survival may account for the observed fertility response. Evaluating these channels requires data that jointly capture the expansion of Blue Cross, fertility outcomes, health outcomes, and local medical conditions.

4 Data

This paper combines several newly digitized and existing historical data sources to construct a county-level panel spanning 1927 to 1947. The data link the rollout of Blue Cross hospital plans to measures of fertility, mortality, hospital use and supply, and local medical costs.

4.1 Data Sources and Variable Construction

I compile data on Blue Cross hospital plans from Louis S. Reed’s *Blue Cross and Medical Service Plans*, a 1947 report published by the U.S. Public Health Service. As voluntary health insurance expanded during this period, the U.S. Public Health Service undertook a comprehensive survey of Blue Cross and Blue Shield programs. I use the report to identify the first year of Blue Cross enrollment in each county and to construct annual measures of plan enrollment.³ Because the report

³Information on the geographic areas served and the timing of first enrollment comes from Table 3, Appendix B, and the “Note on Recent Changes” on p. 31. Annual plan-level enrollment comes from Appendix C and is directly observed for 1935–1946. Because the analysis sample spans 1927–1947, I impute enrollment for years not covered by these annual enrollment data. Eight of the 85 plans in the sample began enrollment before 1935—one in 1932, four in 1933, and three in 1934. For these plans, I assume enrollment is zero immediately prior to the first enrollment

records plan activity through 1947, I observe first-year enrollments through that year. To construct annual enrollment shares, I interpolate county-level population using the 1920, 1930, 1940, and 1950 Censuses and aggregate these estimates to the plan service-area level. I also collect monthly group subscription rates and Blue Shield enrollment from the report.⁴

I use vital statistics data from multiple sources to measure fertility and mortality outcomes. County-level data on births, infant deaths, and total deaths from 1927 to 1947 come from the ICPSR dataset *U.S. County-Level Natality and Mortality Data, 1915–2007* [Bailey et al., 2018]. I define the birth rate as the number of births per 100 women aged 15 to 44, with annual counts of women in this age group linearly interpolated between the 1920, 1930, 1940, and 1950 Censuses.⁵ The infant mortality rate is defined as the number of infant deaths per 100 births.⁶ I define the total mortality rate as total deaths relative to the county population, with annual population counts likewise interpolated between decennial Censuses. For maternal mortality, I use city-level data from 1929 to 1940 compiled by Thomasson and Treber [2008]. These data are drawn from U.S. vital statistics for cities with populations exceeding 10,000, and maternal mortality is defined as deaths related to childbirth and its complications per 100,000 live births.

All hospital-related data are drawn from the American Medical Association’s annual *Hospital Service in the United States* reports, using the 1927–1942 data digitized by Thomasson and Treber [2008]

year and linearly interpolate between zero and the observed 1935 enrollment level. Appendix Figure A1 supports this interpolation approach: using only observed enrollment data, it plots average enrollment shares by years since first enrollment and shows approximately linear growth during the early years following enrollment. I impute 1947 enrollment using each plan’s enrollment growth rate over 1944–1946. These imputed values are used when enrollment shares enter the regression analysis, either as an explanatory variable or as an outcome.

⁴Group subscription rates come from Appendix D, Table 2, which reports monthly rates as of December 1, 1946. Blue Shield enrollment comes from Appendix I, which reports enrollment in each medical service plan as of January 1 for 1940–1947.

⁵Whenever available, I use births by county of residence rather than county of occurrence. Residence-based births more closely capture fertility among county residents, whereas occurrence-based births assign births to the county in which the delivery takes place. This distinction is particularly relevant in this setting because women may travel to neighboring counties to give birth in hospitals, so changes in hospital access may affect the geographic distribution of births by occurrence. Because county-level births by residence are unavailable before 1937, I use births by occurrence from 1927–1936 and births by residence from 1937 onward. Following Holt et al. [2026] and Downes [2024], I assess the comparability of the two measures during years in which both are available by calculating the cross-county correlation between births by residence and births by occurrence for each year. Appendix Figure A2 reports these correlations, which exceed 0.99 in every overlapping year.

⁶As with birth counts, infant deaths by county of residence are preferred because they more closely capture mortality among county residents. However, county-level infant deaths by residence are available only beginning in 1942. I use infant deaths by occurrence from 1927–1941 and infant deaths by residence from 1942 onward. To maintain a consistent geographic basis for the infant mortality rate, I use births by occurrence as the denominator from 1927–1941 and births by residence from 1942 onward.

and newly digitized reports for 1943–1947. The reports provide measures of hospital use, supply and capacity, including hospital counts, bed capacity, bassinet availability, hospital births, average census, and admissions.⁷ I aggregate the hospital-level measures to the county level.

To capture cross-sectional variation in medical care costs prior to widespread insurance coverage, I digitize data from *Intercity Differences in Costs of Living in March 1935* [Stecker, 1937], a 1937 Works Progress Administration research monograph comparing living costs across 59 U.S. cities.⁸ The report constructs a measure of minimum medical service costs based on the expected minimum medical care required over a lifetime, prorated to an annual amount for a four-person family. The measure reflects the cost of a standardized bundle of medical services, including physicians', hospital, nursing, and dental services, rather than the realized medical expenditures of a particular family in a given year. Because cost data are available for only 59 cities, I assign each county the medical cost measure of the geographically nearest city to proxy for local medical costs. I use this measure to capture baseline differences in the cost of medical care faced by households prior to the expansion of hospital insurance. Figure 4 illustrates the geographic distribution of medical costs, with crosses indicating the 59 cities for which cost data are directly observed. I classify counties with minimum medical service costs above the sample mean as "High Baseline Medical Cost Areas" and those below the mean as "Low Baseline Medical Cost Areas."

Finally, I construct additional county-level control variables from a range of historical sources to account for demographic and economic conditions. Appendix C provides detailed definitions, sources, and construction procedures for these variables.

4.2 Blue Cross Roll-out and Treatment Variation

The county-level Blue Cross data capture variation in both the timing and geographic reach of plan entry. Blue Cross coverage expanded gradually across the country during the 1930s and 1940s.⁹

⁷Hospital counts and bed capacity are observed in all years. Average census, average number of patients occupying hospital beds on a given day, is available in all years except 1928, and bassinet availability is reported beginning in 1928. Hospital birth counts are available in 1929, 1931, and annually from 1933 onward, while admissions are observed beginning in 1931.

⁸Minimum medical service costs are drawn from Appendix Table 14 of the report.

⁹When a county was served by more than one Blue Cross plan, I assign the first enrollment year of the earliest plan.

Plans began enrollment at different points in time and differed in their geographic scope. Some plans served entire states, while others covered only particular cities or groups of counties. Figure 5 maps the first year of Blue Cross enrollment for each county. There is no obvious geographic gradient in the timing of entry, as earlier- and later-adopting areas appear across different regions of the country.

By 1947, 31 states and the District of Columbia were each covered by a Blue Cross plan spanning the entire jurisdiction. In another 15 states, coverage was provided through local plans: seven were fully covered by networks of local plans, while eight contained counties that remained outside the service area of any Blue Cross plan. Two states had no Blue Cross plan. Across these jurisdictions, 85 Blue Cross plans were in operation by 1947.

By the end of the sample period, Blue Cross had reached most U.S. counties. To examine whether earlier- and later-adopting counties differed systematically, I estimate a cross-sectional regression of the first year of Blue Cross enrollment on baseline county demographic and economic characteristics. Table 4 shows that earlier enrollment was more common in larger counties, indicating that the timing of Blue Cross entry was correlated with some baseline county characteristics. These differences in levels motivate an empirical design that compares changes in outcomes over time rather than levels across counties. In the next section, I describe the stacked difference-in-differences design used to exploit the staggered timing of Blue Cross entry.

5 Empirical Strategy

5.1 The Effect of Blue Cross Hospital Plans

This section describes the empirical strategy used to estimate the effects of Blue Cross hospital plans on insurance enrollment, fertility, hospital use, and mortality. Identification exploits variation in the timing and geographic spread of Blue Cross plans across counties.

An important feature of this setting is that the expansion of Blue Cross was not primarily motivated by fertility or maternity-specific objectives. Blue Cross plans were broad hospital insurance

programs intended to finance hospital care across a range of medical conditions, with maternity care representing only one component of covered hospital services. The timing of Blue Cross entry is therefore less likely to have been driven by county-specific changes in fertility or fertility intentions. This institutional feature supports the use of variation in Blue Cross introduction to identify its effects on fertility, subject to the identifying assumptions discussed below.

To exploit this variation, I account for the staggered timing of Blue Cross introduction across counties. In settings with staggered treatment adoption, conventional two-way fixed-effects difference-in-differences estimators may combine comparisons between newly treated counties and counties that were treated earlier. As shown by Goodman-Bacon [2021] and de Chaisemartin and D’Haultfoeulle [2020], such comparisons can complicate the interpretation of the estimated treatment effect when treatment effects vary across cohorts or over time since treatment. This concern is particularly relevant in my setting because relatively few counties remain untreated by the end of the sample period. Therefore, I use the weighted stacked difference-in-differences estimator developed by Wing et al. [2024]. The stacked approach organizes the staggered rollout into a series of cohort-specific sub-experiments. For each treatment cohort, counties receiving Blue Cross in a given year are compared with counties that remain untreated throughout the corresponding event window, excluding comparisons between newly treated and previously treated counties. The resulting sub-experiments are then stacked and reweighted to account for differences in the composition of treated and control observations across sub-experiments. The weighted estimator identifies a trimmed aggregate average treatment effect on the treated while maintaining a consistent composition of treatment cohorts across event time.

For each first-enrollment year, I construct a sub-experiment containing counties first exposed to Blue Cross in that year and control counties that remain untreated through the end of the sub-experiment’s post-treatment window. I retain five pre-treatment years and ten post-treatment event years, from $e = -5$ through $e = 9$, and normalize the year immediately preceding treatment, $e = -1$, to zero. Given the 1927–1947 sample period, a balanced event window is available for treatment cohorts from 1932 through 1938.

I estimate the following weighted event-study specification:

$$\text{Outcome}_{ise} = \beta_0 + \sum_{k \neq -1} \delta_k [\text{Treated}_{is} \times \mathbf{1}(e = k)] + \beta_1 \text{Treated}_{is} + \lambda_e + X'_{ise} \Gamma + \varepsilon_{ise}$$

where i indexes counties, s indexes stacked sub-experiments, and e denotes event time relative to the first enrollment year. Treated_{is} equals one for counties belonging to the treatment cohort in sub-experiment s , and λ_e denotes event-time fixed effects. The vector X_{ise} contains county-level demographic and economic controls, including log population, the shares of the population that are Black, urban, and foreign-born, and retail sales per capita. Standard errors are clustered by Blue Cross plan for counties served by a plan and by state for counties without a Blue Cross plan.

The coefficients δ_k trace the evolution of the effect of Blue Cross relative to the year immediately preceding treatment. Identification requires that, absent Blue Cross introduction, treated and control counties would have followed parallel trends in the outcomes. Estimates for $k < 0$ provide evidence on the plausibility of this assumption by showing whether treated and control counties exhibited differential trends prior to Blue Cross entry, while estimates for $k \geq 0$ capture the dynamic effects following introduction. I also report the average of the post-treatment event-study coefficients as a summary measure of the treatment effect.

5.2 The Effect of Enrollment

The event-study specification above estimates the reduced-form effect of the introduction of Blue Cross hospital plans. I estimate the effect of Blue Cross enrollment as well. Enrollment may be endogenous if it expanded more rapidly in counties experiencing other changes that also affected fertility, such as income and employment growth. This motivates an instrumental-variables approach in which the introduction of Blue Cross is used as an instrument for the county-level enrollment share.¹⁰

¹⁰The weighted stacked difference-in-differences estimator developed by Wing et al. [2024] is not extended to instrumental-variables estimation. I assess the finite-sample performance of the IV estimator using a Monte Carlo simulation that preserves the stacked sample structure, corrective weights, fixed effects, and plan-level clustering used in the empirical analysis. In each replication, I simulate county-year enrollment and outcome errors with separate plan-level components and correlated county-year components, with the latter generating endogeneity between enrollment and the outcome. The first-stage parameters are calibrated to match the empirical strength of the relationship between Blue Cross availability and enrollment: the simulated median first-stage F -statistic is 62.2, compared with

$$\text{Outcome}_{ise} = \gamma_0 + \theta \text{Enrollment}_{ise} + \gamma_1 \text{Treated}_{is} + \mu_e + X'_{ise} \Theta + \nu_{ise}$$

I consider three IV specifications for the county-level enrollment share.¹¹ The baseline Binary IV specification uses an indicator that equals one for treated counties beginning in the year Blue Cross becomes available. The Event-Time IV specification allows the effect of Blue Cross introduction on enrollment to vary flexibly with the number of years since entry by using separate post-treatment event-time indicators as instruments. Finally, the Linear Trend IV specification uses a single instrument that increases linearly with the number of years since Blue Cross entry. Across all specifications, I retain the same stacked sample, corrective weights, controls, and fixed effects as in the reduced-form analysis, and standard errors are clustered at the Blue Cross plan level. The IV interpretation additionally requires that, conditional on the stacked design and included controls, the introduction of Blue Cross affects the outcomes through its effect on insurance enrollment.

The coefficient θ captures the effect of Blue Cross enrollment induced by the introduction of a Blue Cross plan. Because enrollment is measured as a share ranging from zero to one, θ represents the effect on the outcome of a one-unit, or 100-percentage-point, increase in enrollment.

6 Empirical Results

This section presents the main empirical results. I begin by investigating the effect of Blue Cross introduction on insurance enrollment, which establishes the first-stage relationship underlying the instrumental-variables analysis. I then examine effects on hospital childbirth and fertility by Blue Cross entry and enrollment, followed by evidence on potential mechanisms.

61.6 in the data using a binary instrument. Across 10,000 replications under a true treatment effect of zero, the mean IV estimate is -0.008 , the Monte Carlo standard deviation of the estimates is 0.136 , and the mean cluster-robust standard error reported by the estimator is 0.132 . Thus, the reported standard errors closely approximate the actual sampling variation of the IV estimator across repeated samples. In addition, nominal 95 percent confidence intervals contain the true treatment effect in 95.1 percent of the simulated samples. In other words, if the sampling process were repeated many times under the simulated data-generating process, confidence intervals constructed using the estimated cluster-robust standard errors would include the true parameter approximately 95 percent of the time. Overall, the close correspondence between the simulated sampling variation and reported standard errors suggests that the weighted stacked IV estimator performs well for inference in this empirical setting.

¹¹As noted in Section 4, enrollment data are available at the Blue Cross plan level rather than the county level. I construct county-level enrollment shares by assigning each county the enrollment share of the Blue Cross plan serving that county.

6.1 The Effect of Blue Cross Hospital Plans on Enrollment Shares

I first estimate the first-stage effect of Blue Cross hospital plans on enrollment shares. Because enrollment is observed at the plan level, I conduct this analysis at the plan level. Table 5 reports the average pre- and post-treatment estimates for enrollment and fertility outcomes, while Figure 6 plots the dynamic event-study estimates for enrollment share and corresponding 95 percent confidence intervals. The estimates show a gradual and sustained increase in enrollment following the introduction of Blue Cross. This pattern is consistent with the way Blue Cross coverage expanded: the introduction of a plan made coverage available, but enrollment did not occur immediately, as employers first had to offer the plan and households then had to enroll. Enrollment therefore accumulated over time rather than increasing discretely upon plan entry. By nine years after first enrollment, Blue Cross had increased the enrollment share by approximately 19 percentage points, while the average increase across the ten post-treatment event years was approximately 8.8 percentage points.¹²

6.2 The Effect of Blue Cross Hospital Plans on Fertility

I turn to the main outcome of interest: fertility. I begin with hospital births, the form of care most directly affected by hospital insurance. I then consider non-hospital births to assess whether increases in hospital births primarily reflect a shift in the location of delivery or an increase in births overall. Finally, I examine the total birth rate. Table 5 summarizes the average pre- and post-treatment estimates for the three fertility outcomes, while Figures 7–9 present the corresponding dynamic event-study estimates.

Figure 7 presents event-study estimates for hospital births per 100 women aged 15–44. Averaging across the ten post-treatment event years, Blue Cross increases the hospital birth rate by 0.37 births per 100 women, and the estimate is statistically significant at the 1 percent level. The pre-treatment coefficients provide little evidence of differential trends prior to Blue Cross entry. Relative to the pre-treatment mean of 0.91 hospital births per 100 women, the estimated effect corresponds to an increase of approximately 41 percent. The effect grows gradually over the post-treatment period,

¹²The county-level specification produces very similar dynamics: the estimated effect rises to roughly 17 percentage points by event year 9, and the average post-entry effect is 7.5 percentage points. Appendix Figure A3 and Appendix Table A1 report the corresponding county-level estimates.

broadly tracking the expansion of Blue Cross enrollment documented above.

I next turn to non-hospital births to assess whether the increase in hospital births primarily reflects substitution from non-hospital to hospital delivery. Figure 8 presents the corresponding estimates for non-hospital births per 100 women. I find no statistically significant decline in non-hospital birth rate following Blue Cross entry. Thus, the results provide little evidence that the increase in hospital birth rate was driven primarily by a shift of births from non-hospital to hospital settings.

Finally, Figure 9 presents the effect of Blue Cross hospital plans on the total birth rate, measured as births per 100 women aged 15–44 regardless of delivery location. Averaging across the ten post-treatment event years, Blue Cross increases the birth rate by approximately 0.31 births per 100 women. The estimate is similar in magnitude to the increase in hospital birth rate, although it is estimated imprecisely. As with hospital birth rate, the estimated effect grows gradually over the post-treatment period, consistent with the gradual expansion of Blue Cross enrollment. I also examine heterogeneity by hospital availability. Appendix Figure A7 and Appendix Table A2 show that the post-treatment point estimate is larger in counties with a hospital than in counties without one. However, neither subgroup estimate is statistically significant, and I cannot reject equality of the two post-treatment effects.

Taken together, the reduced-form estimates suggest that the introduction of Blue Cross increased hospital birth rates without a corresponding decline in non-hospital birth rates, while the estimated effect on the total birth rate was similar in magnitude to the effect on the hospital birth rate.¹³ This pattern is consistent with Blue Cross entry affecting not only the location of delivery but also fertility, although the total birth rate estimate is imprecisely estimated.

6.3 The Effect of Enrollment on Fertility

The reduced-form estimates above capture the effect of the introduction of Blue Cross hospital plans. I use the introduction of Blue Cross as an instrument for Blue Cross enrollment share to estimate the effect of insurance enrollment on fertility. Table 6 reports estimates from OLS and the three

¹³Corresponding estimates using birth counts rather than rates per 100 women aged 15–44 are reported in Appendix Figures A4–A6 and Appendix Table A3.

IV specifications described in Section 5.2: the Binary IV, Event-Time IV, and Linear Trend IV specifications.

The OLS estimates are somewhat smaller than the corresponding IV estimates, particularly for the total birth rate. This difference may reflect measurement error in the enrollment share or endogenous enrollment growth in areas experiencing other economic or demographic changes that affected fertility.

The IV estimates are broadly consistent across the three specifications. The similarity of the estimates suggests that the results are not particularly sensitive to whether the first-stage relationship is modeled as a discrete shift, allowed to vary flexibly with event time, or approximated by a linear trend. For hospital births, an increase in enrollment from 0 to 100 percent is associated with an increase of approximately 4.8 hospital births per 100 women aged 15–44 across all three specifications. The estimates are statistically significant at the 1 percent level. The corresponding estimates for the total birth rate are also positive, although imprecisely estimated. The Event-Time and Linear Trend IV specifications imply an increase of approximately 2.5–2.6 births per 100 women as enrollment increases from 0 to 100 percent, while the Binary IV estimate is at 3.6. None of the total birth rate IV estimates is statistically significant. The first-stage F -statistics indicate a strong relationship between Blue Cross introduction and enrollment across the three specifications.

To gauge the economic magnitude of the fertility response, I use the Event-Time IV estimate of 2.53 births per 100 women. Blue Cross enrollment increased from less than 1 percent in the mid-1930s to approximately 20 percent by the end of the sample period. Applying the IV estimate to an approximately 20-percentage-point increase in enrollment implies an increase in the birth rate of about 0.5 births per 100 women. Over the same period, the U.S. birth rate rose by 3.75 births per 100 women. The point estimate corresponds to roughly 13 percent of the observed increase in the birth rate over this period. The upper bound of the 95 percent confidence interval corresponds to an increase of approximately 1.5 births per 100 women, or about 41 percent of the overall increase in the birth rate. Because the confidence interval includes zero, the data are also consistent with no effect of Blue Cross enrollment on fertility. I interpret the 13 percent figure as an illustrative calculation

based on the point estimate rather than as a precise estimate of Blue Cross’s contribution to the reversal of fertility.

6.4 Mechanisms

Having established that Blue Cross enrollment increased hospital births and may also have increased fertility, I next investigate the mechanisms through which these effects may have operated. The conceptual framework in Section 3 provides three empirical predictions that help distinguish among potential mechanisms. First, if Blue Cross affects childbirth and fertility through the financial cost of hospital delivery, its effects should be larger where the underlying cost of medical care is higher, since insurance provides greater financial protection in these areas. Second, if maternal health is an important channel, Blue Cross introduction should be accompanied by improvements in maternal health outcomes. Third, if the fertility response operates through improved infant survival, Blue Cross should reduce infant mortality if increased access to hospital delivery and medical care improved infant survival. The following subsections examine these predictions in turn.

6.4.1 Financial Channel

I examine the financial channel by testing whether the effects of Blue Cross are larger in areas with higher baseline medical costs.¹⁴ Because Blue Cross reduced the out-of-pocket cost of hospital care, the financial benefit of coverage should be greater in areas where hospital and medical services were more expensive. To examine this prediction, I divide counties into high- and low- baseline medical cost areas based on 1935 medical costs and estimate the effect of Blue Cross separately for the two groups.

I first focus on the hospital birth rate, the outcome most directly affected by hospital insurance. Figure 10 presents the event-study estimates, and Table 7 reports the corresponding average pre- and post-treatment effects. In counties with high baseline medical costs, Blue Cross increases hospital

¹⁴One potential concern is that Blue Cross premiums may have reflected local medical costs, which could attenuate or remove heterogeneity in the effects of coverage by baseline medical costs. To examine this possibility, Appendix Figure A8 plots 1935 city-level medical costs against the corresponding monthly Blue Cross premiums reported in Reed [1947] for two-person and family plans, separately for semiprivate rooms and wards. I focus on two-person and family plans because these plan types typically included maternity benefits. The figure shows little correlation between Blue Cross premiums and baseline medical costs.

births by 0.51 births per 100 women aged 15–44, compared with 0.16 births per 100 women in counties with low baseline medical costs. The difference in the average post-treatment effects is 0.35 births per 100 women and is statistically significant at the 10 percent level. The dynamic estimates also show a substantially larger increase in hospital births in high-cost counties following Blue Cross entry. This pattern is consistent with the financial benefit of hospital insurance being greater where medical care was more expensive.

I next look at whether the same pattern is present for total birth rate. Figure 11 shows that the estimated effect is also larger in high-cost counties. The average post-treatment estimate is 0.55 births per 100 women in high-cost counties, compared with -0.06 births per 100 women in low-cost counties (Table 7). The difference of 0.61 births per 100 women is economically sizable, although it is not statistically significant. The stronger response in high-cost areas is consistent with a financial channel although the heterogeneity in total fertility is estimated imprecisely.¹⁵

Taken together, the results are consistent with the prediction that Blue Cross had larger effects in areas where medical care was more costly. The stronger response in high-cost areas provides evidence consistent with a financial channel in which insurance reduced the financial burden of hospital childbirth.

6.4.2 Maternal Health Channel

I next turn to the health-related mechanisms, beginning with maternal health. If Blue Cross affected fertility by improving maternal health, its introduction should be associated with better maternal health outcomes. Because broader measures of maternal health are unavailable for this period, I use maternal mortality as a proxy. This is a particularly severe measure of maternal health and may therefore fail to capture improvements that do not translate into changes in mortality.

The evidence does not indicate a decline in maternal mortality following Blue Cross entry. Table 8 shows an average post-treatment estimate of 38, relative to a pre-treatment mean of 813; the estimate is not statistically significant. The event-study estimates in Figure 12 likewise fluctuate

¹⁵Corresponding estimates for hospital birth counts, non-hospital birth rates and counts, and total birth counts are reported in Appendix Figures A9–A12 and Appendix Table A4.

around zero rather than exhibiting a sustained decline.¹⁶

These results should be interpreted with the limitation that maternal mortality captures only the most severe maternal health outcomes. Thus, the absence of a mortality decline does not rule out improvements in other dimensions of maternal health. Nevertheless, within the outcome available in the historical data, I therefore find little evidence from maternal mortality in support of the maternal health channel.¹⁷

6.4.3 Infant Survival Channel

The final mechanism I consider is infant survival. For improved infant survival to explain the fertility response, Blue Cross should have reduced infant mortality following its introduction.

Table 8 reports the average pre- and post-treatment estimates. The results provide little evidence that Blue Cross reduced infant mortality. The average post-treatment effect is 0.044 infant deaths per 100 births and is not statistically significant, relative to a pre-treatment mean of 5.845. Figure 13 presents the corresponding event-study estimates. The coefficients similarly show no systematic decline in infant mortality following the introduction of Blue Cross.¹⁸

These results provide little evidence that the expansion of Blue Cross improved infant survival over this period, suggesting that improvements in infant survival are less likely to account for the estimated fertility response to Blue Cross.

7 Discussion

To put these results in context, I show that the introduction of Blue Cross was not accompanied by an expansion in hospital capacity. This evidence suggests that the null effects on infant and maternal health are unlikely to reflect capacity constraints, strengthening the interpretation that

¹⁶The event-study window is shorter than for the other outcomes because the maternal mortality data span only 1928–1940.

¹⁷The main specification uses a balanced panel of 262 cities. Estimates using an unbalanced panel of more than 800 cities are qualitatively similar. Thomasson and Treber [2008] also reports maternal mortality rates separately by race. Race-specific estimates yield a similar conclusion and are reported in Appendix Table A5 and Appendix Figure A13.

¹⁸Results using the number of infant deaths rather than the infant mortality rate are reported in Appendix Table A5 and Appendix Figure A14.

Blue Cross did not meaningfully affect these outcomes.

7.1 Hospital Capacity

The substantial increase in hospital childbirth following Blue Cross entry raises a natural question about the supply side of the hospital market. If Blue Cross increased demand for hospital delivery, did hospitals have sufficient capacity to accommodate the additional births? By a simple measure of maternity capacity, this appears to have been the case. In 97 percent of county-year observations in the sample, the number of bassinets was at least as large as the number required to accommodate observed hospital births.¹⁹ This suggests that, in most counties and years, existing maternity capacity was sufficient to accommodate the observed level of hospital childbirth.

A related question is whether Blue Cross itself induced an expansion in hospital capacity. Appendix Table A6 reports the average pre- and post-treatment estimates, and Appendix Figure A15 presents the corresponding event-study coefficients. I first examine bassinet capacity, which is most directly relevant for hospital childbirth. I find little evidence that Blue Cross increased bassinet capacity relative to the population of women of childbearing age. The average post-treatment effect on bassinets per 100 women aged 15–44 is 0.001 and is not statistically significant, relative to a pre-treatment mean of 0.097.²⁰

I also examine hospital beds as a broader measure of capacity. Although the post-treatment estimates for hospital beds are positive, the event-study estimates raise concerns about the identifying pre-trends. In particular, the average pre-treatment estimate for beds per capita is negative and statistically significant. I therefore interpret the post-treatment increases in bed capacity with caution rather than as evidence of a causal supply response to Blue Cross.

¹⁹ Assuming a 10-day hospital stay for each birth, I calculate the number of bassinets required to accommodate observed hospital births as the annual number of hospital births multiplied by 10 and divided by 365. I then compare this implied number of required bassinets with the observed number of bassinets in each county and year. I classify a county-year as having sufficient capacity when the observed number of bassinets is at least as large as the implied number required.

²⁰ When I instead use the number of bassinets as the outcome, the average post-treatment estimate is positive and statistically significant: Blue Cross entry is associated with an increase of 6.85 bassinets, relative to a pre-treatment mean of 9.98. However, this increase in the absolute number of bassinets does not translate into an increase in bassinets per 100 women aged 15–44. I therefore interpret the results as providing little evidence that Blue Cross increased bassinet capacity relative to the population potentially requiring maternity care.

Finally, I examine whether Blue Cross was associated with changes in the number of general hospitals. On the extensive margin, the probability that a county had a general hospital is essentially unchanged following Blue Cross entry, with an average post-treatment estimate of -0.008. On the intensive margin, the estimates for both general hospitals per capita and the total number of general hospitals are small and statistically insignificant. The results provide little convincing evidence that the increase in hospital childbirth following Blue Cross entry was accompanied by a contemporaneous expansion in measured hospital capacity.

7.2 Other Health Outcomes

I consider why the substantial increase in hospital childbirth did not translate into detectable improvements in maternal or infant mortality. Contemporary accounts show that greater use of hospital childbirth did not necessarily translate into mortality improvements during this period. Hospital delivery provided access to medical intervention for complicated births, but these interventions could also introduce additional risks. Polak [1929], for example, attributed persistently high maternal mortality in part to unnecessary intervention in uncomplicated deliveries, avoidable cesarean sections, and exposure to cross-infection in hospitals. Consistent with this historical evidence, Thomasson and Treber [2008] find that medical intervention had little effect on maternal mortality before the widespread introduction of sulfa drugs in the late 1930s and became mortality-reducing only after these drugs lowered the risk of infection. Contemporary accounts document similar risks for newborns. In particular, outbreaks of highly fatal diarrhea were reported in hospital newborn nurseries, raising concerns about cross-infection associated with congregate newborn care [Spivek, 1938; Abramson and Frant, 1938; Oppenheimer and United States Children’s Bureau, 1936]. Thus, the expansion of hospital childbirth could have improved care along some dimensions while simultaneously exposing mothers and newborns to risks that attenuated its net effect on mortality.

As a broader check, I examine additional measures of hospital utilization and health outcomes. Appendix Table A7 reports the average pre- and post-treatment estimates, and Appendix Figure A16 presents the corresponding event-study coefficients. Average daily census, defined as the average number of inpatients present in a hospital each day, provides a measure of hospital utilization. The

average post-treatment estimate for average daily census per capita is 0.068, relative to a pre-treatment mean of 0.354, while the level of average daily census increases by approximately 121 patients, relative to a pre-treatment mean of 169. However, both measures exhibit evidence of differential pre-treatment dynamics, so I interpret these increases with caution rather than as clean causal estimates.

I also examine broader mortality outcomes that are not restricted to mothers or infants. I find little evidence of a decline in mortality following Blue Cross entry. The average post-treatment effect on the mortality rate is 0.003 and is not statistically significant, relative to a pre-treatment mean of 1.009. The number of deaths also shows no statistically significant change. The corresponding event-study estimates show no systematic decline in either measure following Blue Cross entry. Thus, although hospital utilization appears to have increased, I find little evidence that greater use of hospital care was accompanied by improvements in mortality. This pattern is consistent with the earlier findings for maternal and infant mortality.

8 Conclusion

This paper examines whether the emergence of employer-sponsored hospital insurance affected childbirth and fertility during the period in which the long-run decline in U.S. fertility began to reverse. I construct a new dataset tracing the geographic expansion and enrollment of Blue Cross hospital plans and link these data to county-level measures of hospital childbirth, fertility, and mortality. Exploiting variation in the timing and geographic rollout of Blue Cross, I estimate weighted stacked difference-in-differences models and use Blue Cross introduction as an instrument for insurance enrollment. Together, these approaches provide evidence on how the early expansion of hospital insurance affected both the use of hospital care and childbearing behavior.

The results show that Blue Cross substantially increased insurance enrollment and hospital childbirth. Following Blue Cross introduction, hospital births increased by 0.37 births per 100 women aged 15–44, relative to a pre-treatment mean of 0.91. I find little evidence of a corresponding decline in non-hospital births, suggesting that the increase in hospital childbirth was not driven solely by

a shift in the location of deliveries that would otherwise have occurred outside hospitals. The estimated effect on the total birth rate is also positive and similar in magnitude, although it is estimated imprecisely. Instrumental-variables estimates yield the same qualitative pattern: greater Blue Cross enrollment significantly increases hospital childbirth, while the estimated effects on total fertility are positive but imprecise.

The mechanism evidence points most strongly toward the financial cost of childbirth. The effects of Blue Cross on hospital births are significantly larger in areas with higher baseline medical costs, where insurance provided greater financial protection, and the estimated fertility response is also larger in these areas, although the difference is less precisely estimated. In contrast, I find little evidence that Blue Cross reduced maternal or infant mortality.

More broadly, the results highlight the demographic consequences of changes in the financing of medical care. Blue Cross emerged not as a fertility policy but as a broad system for financing hospital services, yet maternity care quickly became one of its most important uses. Its expansion therefore altered the economic environment surrounding childbirth at the same time that U.S. fertility was beginning to rise after decades of decline. Although the fertility estimates are imprecise and do not imply that Blue Cross alone explains the reversal, the evidence suggests that the emergence of employer-sponsored hospital insurance may have been one institutional development that contributed to this turning point. More generally, the results show that the consequences of health insurance can extend beyond health-care utilization itself, influencing household decisions about childbearing and family formation.

References

Virginia medical monthly, 1923.

Harold Abramson and Samuel Frant. Epidemic diarrhea of the newborn: Iv. clinical considerations. *American Journal of Diseases of Children*, 55(6):1288–1307, 1938. doi: 10.1001/archpedi.1938.01980120150015. URL <https://doi.org/10.1001/archpedi.1938.01980120150015>.

Stefania Albanesi and Claudia Olivetti. Maternal health and the baby boom. *Quantitative Economics*, 5(2):225–269, 2014. doi: <https://doi.org/10.3982/QE315>. URL <https://onlinelibrary.wiley.com/doi/abs/10.3982/QE315>.

Stefania Albanesi and Claudia Olivetti. Gender Roles and Medical Progress. *Journal of Political Economy*, 124(3):650–695, June 2016. ISSN 0022-3808. doi: 10.1086/686035. URL <https://www.journals.uchicago.edu/doi/10.1086/686035>. Publisher: The University of Chicago Press.

Martha Bailey, Karen Clay, Price Fishback, Michael R. Haines, Shawn Kantor, Edson Severnini, and Anna Wentz. U.s. county-level natality and mortality data, 1915-2007. *Inter-university Consortium for Political and Social Research [distributor]*, 5 2018. URL <https://doi.org/10.3886/ICPSR36603.v2>.

Martha J. Bailey and William J. Collins. Did improvements in household technology cause the baby boom? evidence from electrification, appliance diffusion, and the amish. *American Economic Journal: Macroeconomics*, 3(2):189–217, April 2011. doi: 10.1257/mac.3.2.189. URL <https://www.aeaweb.org/articles?id=10.1257/mac.3.2.189>.

Martha J. Bailey and Brad J. Hershbein. Us fertility rates and childbearing in american history, 1800–2010. In Louis P. Cain, Price V. Fishback, and Paul W. Rhode, editors, *The Oxford Handbook of American Economic History, Volume 1*, pages 75–100. Oxford University Press, Oxford, 2018.

Gary S. Becker and Robert J. Barro. A reformulation of the economic theory of fertility. *The Quarterly Journal of Economics*, 103(1):1–25, 1988. doi: 10.2307/1882640. URL <https://doi.org/10.2307/1882640>.

- Andriana Bellou, Emanuela Cardia, and Joshua Lewis. From bust to boom: The great depression and women’s fertility. IZA Discussion Paper 18120, Institute of Labor Economics (IZA), 2025. URL <https://www.iza.org/publications/dp/18120/from-bust-to-boom-the-great-depression-and-womens-fertility>. IZA Discussion Paper No. 18120, September 2025.
- Marianne P. Bitler and Lucie Schmidt. Utilization of infertility treatments: The effects of insurance mandates. *Demography*, 49(1):125–149, 2012. doi: 10.1007/s13524-011-0078-4. URL <https://read.dukeupress.edu/demography/article/49/1/125/169686/Utilization-of-Infertility-Treatments-The-Effects>.
- Abel Brodeur and Lamis Kattan. World war ii, the baby boom, and employment: County-level evidence. *Journal of Labor Economics*, 40(2):437–471, 2022. doi: 10.1086/715485. URL <https://www.journals.uchicago.edu/doi/10.1086/715485>.
- Alessandro Cigno. Fertility decisions when infant survival is endogenous. *Journal of Population Economics*, 11:21–28, 1998. doi: 10.1007/s001480050056. URL <https://doi.org/10.1007/s001480050056>.
- Robert Cunningham III and Robert M. Cunningham Jr. *The Blues: A History of the Blue Cross and Blue Shield System*. Northern Illinois University Press, DeKalb, IL, 1997. ISBN 0875802249.
- Clément de Chaisemartin and Xavier D’Haultfoeulle. Two-way fixed effects estimators with heterogeneous treatment effects. *American Economic Review*, 110(9):2964–2996, 2020. doi: 10.1257/aer.20181169. URL <https://doi.org/10.1257/aer.20181169>.
- Thomas DeLeire, Leonard M. Lopoo, and Kosali I. Simon. Medicaid expansions and fertility in the united states. *Demography*, 48(2):725–747, May 2011. doi: 10.1007/s13524-011-0031-6.
- Lisa J. Dettling and Melissa Schettini Kearney. Did the modern mortgage set the stage for the us baby boom? NBER Working Paper 33446, National Bureau of Economic Research, 2025. URL <https://www.nber.org/papers/w33446>.
- Matthias Doepke, Oved Hazan, and Yishay Maoz. The baby boom and world war ii: A macroeco-

- conomic analysis. *The Review of Economic Studies*, 82(3):1031–1073, 2015. doi: 10.1093/restud/rdv001. URL <https://academic.oup.com/restud/article-abstract/82/3/1031/1576288>.
- Henry Downes. Did organized labor induce labor? unionization and the american baby boom. SSRN Working Paper 4870124, SSRN, 2024. URL https://papers.ssrn.com/sol3/papers.cfm?abstract_id=4870124.
- Richard A. Easterlin. On the relation of economic factors to recent and projected fertility changes. *Demography*, 3(1):131–153, March 1966. ISSN 0070-3370. doi: 10.2307/2060068. URL <https://doi.org/10.2307/2060068>. eprint: <https://read.dukeupress.edu/demography/article-pdf/3/1/131/908220/131easterlin.pdf>.
- Price Fishback and Shawn Kantor. New deal studies, 2018. URL <https://doi.org/10.3886/E101199V1>. Distributed November 18, 2018.
- Donna B. Gilleskie, Euna Han, and Edward C. Norton. Disentangling the contemporaneous and dynamic effects of human and health capital on wages over the life cycle. *Review of Economic Dynamics*, 25:350–383, 2017. doi: 10.1016/j.red.2017.02.012. URL <https://doi.org/10.1016/j.red.2017.02.012>.
- Andrew Goodman-Bacon. Difference-in-differences with variation in treatment timing. *Journal of Econometrics*, 225(2):254–277, 2021. doi: 10.1016/j.jeconom.2021.03.014. URL <https://doi.org/10.1016/j.jeconom.2021.03.014>.
- Jeremy Greenwood, Ananth Seshadri, and Guillaume Vandenbroucke. The baby boom and baby bust. *American Economic Review*, 95(1):183–207, March 2005. doi: 10.1257/0002828053828680. URL <https://www.aeaweb.org/articles?id=10.1257/0002828053828680>.
- Timothy W. Guinnane. The historical fertility transition: A guide for economists. *Journal of Economic Literature*, 49(3):589–614, 2011. doi: 10.1257/jel.49.3.589. URL <https://doi.org/10.1257/jel.49.3.589>.
- Michael R. Haines and Inter-university Consortium for Political and Social Research. Historical, demographic, economic, and social data: The united states, 1790–2002, 2010. URL <https://doi.org/10.3886/ICPSR02896.v3>. Distributed May 21, 2010.

Matthew J. Hill. Easterlin revisited: Relative income and the baby boom. *Explorations in Economic History*, 56:71–85, 2015. ISSN 0014-4983. doi: <https://doi.org/10.1016/j.eeh.2014.10.001>. URL <https://www.sciencedirect.com/science/article/pii/S0014498314000412>.

Andrew Chase Holt, Arjina Akter, and Jason Edward Taylor. Fertility in the time of depression: the impact of economic conditions on us fertility during the 1930s. *Journal of Demographic Economics*, page 1–19, 2026. doi: 10.1017/dem.2026.10017.

Institute of Medicine. *Employment and Health Benefits: A Connection at Risk*. National Academy Press, Washington, DC, 1993. doi: 10.17226/2044.

Seema Jayachandran and Adriana Lleras-Muney. Life expectancy and human capital investments: Evidence from maternal mortality declines. *The Quarterly Journal of Economics*, 124(1):349–397, 2009. doi: 10.1162/qjec.2009.124.1.349. URL <https://doi.org/10.1162/qjec.2009.124.1.349>.

Judith Walzer Leavitt. *Brought to Bed: Childbearing in America, 1750–1950*. Oxford University Press, New York, 1986. ISBN 9780195056907.

Arleen Leibowitz. The response of births to changes in health care costs. *The Journal of Human Resources*, 25(4):697–711, 1990. ISSN 0022166X. URL <http://www.jstor.org/stable/145672>.

Joshua Lewis. Infant health, women’s fertility, and rural electrification in the united states, 1930–1960. *The Journal of Economic History*, 78(1):118–154, 2018. doi: 10.1017/S0022050718000050. URL <https://www.cambridge.org/core/journals/journal-of-economic-history/article/infant-health-womens-fertility-and-rural-electrification-in-the-united-states-19301960/6C627B2FE2748A48A0781BEC7701306A>.

Michael Morrissey. *Health Insurance*. AUPHA/HAP Book (Health Administration Press), Chicago, IL, 2 edition, 2013. ISBN 9781567936094.

National Resources Committee. *Consumer Incomes in the United States: Their Distribution in 1935–36*. United States Government Printing Office, Washington, D.C., 1938. For sale by the Superintendent of Documents.

- Ella Oppenheimer and United States Children's Bureau. *Infant Mortality in Memphis*. United States Government Printing Office, Washington, DC, 1936. URL <https://fraser.stlouisfed.org/title/infant-mortality-memphis-8508>. Bureau Publication No. 233.
- John Osborn Polak. Puerperal morbidity and mortality: With especial reference to the effects of previous infection and operative delivery. *JAMA*, 93(19):1436–1440, 1929. doi: 10.1001/jama.1929.02710190008002. URL <https://doi.org/10.1001/jama.1929.02710190008002>.
- Louis Reed. *Blue Cross and Medical Service Plans*. U.S. Public Health Service, Washington, D.C., 1947.
- Louis S. Reed. Legislative proposals for compulsory health insurance. *Law and Contemporary Problems*, 6(4):628–644, 1939. URL <https://scholarship.law.duke.edu/lcp/vol6/iss4/13>. Fall.
- Lucie Schmidt. Effects of infertility insurance mandates on fertility. *Journal of Health Economics*, 26(3):431–446, May 2007. doi: 10.1016/j.jhealeco.2006.10.012.
- Mandel L. Spivek. A consideration of present day newborn nursery practice: Suggestions for improvement. *JAMA*, 111(12):1065–1068, 1938. doi: 10.1001/jama.1938.02790380007003. URL <https://doi.org/10.1001/jama.1938.02790380007003>.
- Paul Starr. *The Social Transformation of American Medicine: The Rise of a Sovereign Profession and the Making of a Vast Industry*. Basic Books, New York, 2 edition, 2017.
- Margaret Loomis Stecker. *Intercity Differences in Costs of Living in March, 1935, 59 Cities*. Research Monograph No. XII, Works Progress Administration, Division of Social Research. U.S. Government Printing Office, Washington, D.C., 1937. Includes analysis of basic living costs across 59 U.S. cities.
- Melvin Jr. Stephens and Desmond Toohey. The impact of health on labor market outcomes: Evidence from a large-scale health experiment. *American Economic Journal: Applied Economics*, 14(3): 367–399, 2022. doi: 10.1257/app.20180686. URL <https://doi.org/10.1257/app.20180686>.

- Elizabeth Temkin. Driving through: Postpartum care during world war ii. *American Journal of Public Health*, 89(4):587–595, April 1999. doi: 10.2105/ajph.89.4.587.
- Melissa A. Thomasson and Jaret Treber. From home to hospital: The evolution of childbirth in the united states, 1928–1940. *Explorations in Economic History*, 45(1):76–99, 2008. ISSN 0014-4983. doi: <https://doi.org/10.1016/j.eeh.2007.07.001>. URL <https://www.sciencedirect.com/science/article/pii/S0014498307000241>.
- Erin Troland and Theodore F. Figinski. Does getting health insurance affect women’s fertility? evidence from the united mine workers’ health insurance. *AEA Papers and Proceedings*, 109: 511–15, May 2019. doi: 10.1257/pandp.20191094. URL <https://www.aeaweb.org/articles?id=10.1257/pandp.20191094>.
- United States Bureau of the Census. County and city data book [united states] consolidated file: City data, 1944–1977, 2008. URL <https://doi.org/10.3886/ICPSR07735.v1>. Distributed April 14, 2008.
- United States Department of Labor, Children’s Bureau. *Proceedings of Conference on Better Care for Mothers and Babies*. United States Government Printing Office, Washington, DC, 1938. Conference held January 17–18, 1938. Bureau Publication No. 246.
- U.S. Bureau of Labor Statistics. Monthly labor review, December 1930.
- U.S. Children’s Bureau. *Maternal Mortality in Fifteen States*. Number 223 in Bureau Publication. U.S. Government Printing Office, Washington, DC, 1934.
- U.S. Children’s Bureau. *Proceedings of Conference on Better Care for Mothers and Babies*. Number 246 in Bureau Publication. U.S. Government Printing Office, Washington, DC, 1938.
- U.S. Children’s Bureau. *Maternity Care at Public Expense in Six Counties in New York State, July 1, 1935–June 30, 1936*. Number 267 in Bureau Publication. U.S. Government Printing Office, Washington, DC, 1941.
- U.S. Department of Labor, Children’s Bureau. The promotion of the welfare and hygiene of maternity

and infancy: The administration of the act of congress of november 23, 1921, fiscal year ended june 30, 1925, 1926.

Richard W. Wertz and Dorothy C. Wertz. *Lying-in A History of Childbirth in America*. Schocken Books, 1979.

Coady Wing, Seth M. Freedman, and Alex Hollingsworth. Stacked difference-in-differences. NBER Working Paper 32054, National Bureau of Economic Research, 2024. URL <https://www.nber.org/papers/w32054>.

Madeline Zavodny and Marianne P. Bitler. The effect of medicaid eligibility expansions on fertility. *Social Science Medicine*, 71(5):918–924, September 2010. doi: 10.1016/j.socscimed.2010.05.046.

Tables and Figures

Table 1: Midwife-Attended Births in New Jersey, 1919–1926

Year	Total Births	Midwife-Attended Births	Percent Midwife-Attended
1919	70,935	30,000	42.2
1920	76,431	21,571	28
1921	78,172	21,106	27
1922	74,479	19,205	26
1923	74,611	18,365	24.5
1924	76,530	17,645	23
1925	74,193	16,077	21.6
1926	72,386	14,739	20.3

Notes: The table reports total births, births attended by midwives, and the share of births attended by midwives in New Jersey from 1919 to 1926. Percent midwife-attended is calculated as midwife-attended births divided by total births multiplied by 100.

Source: 50th Annual Report of the Department of Health of the State of New Jersey (1927).

Table 2: Hospital Cases for Selected Diagnoses, Associated Hospital Service of New York, 1943

Disease Classification	Percent of Total Cases
Maternity	
Delivery	20.89
Others	2.87
Respiratory System	
Tonsillectomy	11.75
Influenza, Bronchitis, and Pneumonia	5.72
Others	4.92

Notes: The table reports the most common diagnostic categories among hospital cases under the Associated Hospital Service of New York in 1943, selected based on their share of total cases. Percent of total cases is calculated relative to all hospital cases reported by the plan.

Source: Reed [1947].

Table 3: Hospital Utilization and Expenses for Selected Diagnoses, New Jersey Blue Cross Plan, 1943

Diagnosis	Percent of Total Cases	Percent of Total Days' Care	Percent of Total Expenses
Maternity	22.97	25.29	28.19
Tonsils & Adenoids	16.69	2.73	4.48
Appendicitis	9.97	11.95	12.03
Female Diseases	9.74	10.62	10.84

Notes: The table reports selected diagnoses from among the most common hospital cases under the New Jersey Blue Cross plan in 1943, based on their share of total cases. Percent of total cases is the share of all hospital cases accounted for by each diagnosis; percent of total days' care is the share of all inpatient days attributable to each diagnosis; and percent of total expenses is the share of total hospital expenses attributable to each diagnosis.

Source: Reed [1947].

Table 4: Baseline County Characteristics and Timing of Blue Cross Entry

	First Blue Cross Enrollment Year
Log total population	−1.178*** (0.326)
Retail sales per capita	0.001 (0.001)
Share of Black population	0.011 (0.042)
Share of foreign-born population	−0.136 (0.084)
Share of urban population	0.031** (0.014)
Log medical costs	2.944 (4.11)
Has a hospital	−0.406 (0.428)

Notes: The table reports coefficients from a cross-sectional regression of the first year of Blue Cross enrollment in a county on baseline county characteristics. Explanatory variables are measured in the earliest year in which they are available. Standard errors are reported in parentheses. *** $p < 0.01$, ** $p < 0.05$, and * $p < 0.10$.

Table 5: The Effect of Blue Cross Hospital Plans

	Average Pre	Average Post
Enrollment Share	0.045 (0.355)	8.8*** (0.849)
Hospital Birth Rate	0.010 (0.057)	0.370*** (0.129)
Pre-treatment Outcome Mean	0.91	
Non-Hospital Birth Rate	0.159 (0.172)	0.131 (0.248)
Pre-treatment Outcome Mean	7.96	
Birth Rate	0.046 (0.152)	0.306 (0.298)
Pre-treatment Outcome Mean	8.94	

Notes: The table summarizes event-study estimates of the effect of Blue Cross plan entry on insurance enrollment and birth outcomes. The Average Pre column reports the average of the event-study coefficients for event years -5 through -2, with event year -1 omitted as the reference period. The Average Post column reports the average of the event-study coefficients for event years 0 through 9. Enrollment share is measured on a 0–100 percentage scale. Hospital, non-hospital, and total birth rates are measured as births per 100 women aged 15–44. The enrollment-share analysis is conducted at the Blue Cross plan level, while the birth-outcome analyses are conducted at the county level. Pre-treatment outcome means are calculated using observations from the pre-treatment period. Standard errors are reported in parentheses. *** $p < 0.01$, ** $p < 0.05$, and * $p < 0.10$.

Table 6: The Effect of Enrollment on Fertility

	OLS	IV		
		Binary	Event-Time	Linear Trend
<i>Hospital Birth Rate</i>				
Enrollment Share	4.384*** (1.502)	4.774*** (1.577)	4.798*** (1.540)	4.809*** (1.545)
First-stage F-statistic		61.482	12.886	56.899
<i>Birth Rate</i>				
Enrollment Share	1.747 (1.523)	3.577 (3.751)	2.529 (2.600)	2.637 (2.676)
First-stage F-statistic		61.277	12.761	56.998

Notes: The table reports estimates of the effect of Blue Cross enrollment on hospital birth rates and total birth rates. The OLS specification directly relates Blue Cross enrollment share to each outcome. In the IV specifications, Blue Cross enrollment share is treated as endogenous. The Binary IV specification uses an indicator for the introduction of Blue Cross as the instrument. The Event-Time IV specification allows the first-stage effect of Blue Cross introduction on enrollment to vary by years since entry, while the Linear Trend IV specification assumes that this effect changes linearly with years since entry. Enrollment share ranges from 0 to 1, and birth rates are measured as births per 100 women aged 15–44. First-stage F-statistics are reported. Standard errors are reported in parentheses. *** $p < 0.01$, ** $p < 0.05$, and * $p < 0.10$.

Table 7: The Effect of Blue Cross Hospital Plans by Baseline Medical Costs

	High Baseline Medical Costs	Low Baseline Medical Costs	Difference
<i>Panel A. Hospital Birth Rate</i>			
Average Pre	-0.018 (0.067)	0.048 (0.094)	—
Average Post	0.512*** (0.162)	0.160 (0.147)	0.352* (0.179)
Pre-treatment Outcome Mean	0.821	1.049	—
<i>Panel B. Birth Rate</i>			
Average Pre	0.010 (0.224)	0.146 (0.155)	—
Average Post	0.553 (0.374)	-0.058 (0.324)	0.611 (0.369)
Pre-treatment Outcome Mean	9.016	8.830	—


Notes: The table reports event-study estimates of the effect of Blue Cross plan entry separately for counties with high and low baseline medical costs. Counties are classified as high or low cost based on whether baseline medical costs are above or below the sample mean. The Average Pre estimates average the event-study coefficients for event years -5 through -2, with event year -1 omitted as the reference period. The Average Post estimates average the event-study coefficients for event years 0 through 9. In Panel A, the outcome is hospital births per 100 women aged 15–44. In Panel B, the outcome is total births per 100 women aged 15–44. The Difference column reports the difference in average post-treatment effects between high- and low-cost counties (High minus Low). Standard errors are reported in parentheses. *** $p < 0.01$, ** $p < 0.05$, and * $p < 0.10$.

Table 8: The Effect of Blue Cross Hospital Plans on Maternal and Infant Mortality Rates

	Average Pre	Average Post
Maternal Mortality Rate	-33.583 (50.216)	37.971 (39.021)
Pre-treatment Outcome Mean	812.899	
Infant Mortality Rate	0.107 (0.117)	0.044 (0.162)
Pre-treatment Outcome Mean	5.845	

Notes: The table reports event-study estimates of the effect of Blue Cross plan entry on maternal and infant mortality. The maternal mortality rate is measured as maternal deaths per 100,000 live births, and the infant mortality rate is measured as infant deaths per 100 births. The Average Pre column reports the average of the event-study coefficients for event years -5 through -2, with event year -1 omitted as the reference period. The Average Post column reports the average of the event-study coefficients for event years 0 through 9. Pre-treatment outcome means are calculated using observations from the pre-treatment period. Standard errors are reported in parentheses. *** $p < 0.01$, ** $p < 0.05$, and * $p < 0.10$.

Figure 1: U.S. Birth Rate and Blue Cross Enrollment



Notes: The solid line plots the U.S. birth rate, measured as births per 100 women ages 15–44. The dashed line plots total Blue Cross enrollment, measured in millions of enrollees. The light shaded region denotes World War II, and the darker shaded region denotes the Baby Boom period beginning in 1946.

Sources: National Center for Health Statistics; Reed (1947); Health Insurance Institute (1963).

Figure 2: Historical Receipts for Hospital Deliveries

Form 13 3-26-20 M. Philadelphia, Pa., 4/12 1930

Mr. Ida [redacted] Dr.
Room 7 Maternity 1 Hall
To Women's Homoeopathic Hospital of Philadelphia
 20th Street - Susquehanna Avenue

For Board.....	<u>4/8-4/15</u>	<u>35 00</u>
" " Special Nurse.....	<u>Booby Board</u>	<u>1 50</u>
" Guest Trays.....		<u>10 00</u>
" Operating Room.....		
" Delivery Service.....		
" Anesthetic.....		
" Ambulance Service.....		<u>5 00</u>
" Dressings.....		
" X-Ray.....		
" Pathological Examination.....		<u>51 50</u>
" Special Medicines.....		
" Telephone Calls.....	<u>Booby Board</u>	<u>10 00</u>
" Dispensary Treatments.....		
" Electrocardiographic Examinations.....		
TOTAL		<u>46 50</u>

Received Payment with Thanks 4/13/30
L. H. Cannon Superintendent

in advance
 Bills must be settled before leaving hospital

13M-4-36-WPS
 Bills Payable in Advance Hospital Day Ends at 4 o'clock P. M.

Brunswick General Hospital
 AMITYVILLE, N. Y.
 TELEPHONE AMITYVILLE 1700 C. L. MARKHAM, M. D., Superintendent
 Room No. 26 MAY 26 1936
 M. [redacted]

Nc 6357

Bill from	to	inclusive	per Day
<u>5/16</u>	<u>7/1</u>		<u>44</u>
<u>11</u>	<u>4</u>		

Operating Room.....	
Anaesthetic.....	<u>10 -</u>
Delivery Room.....	<u>5 50</u>
Baby Care.....	<u>1 -</u>
Laboratory.....	<u>1 10</u>
Special Medicines.....	
Dressings.....	
Special Day Nurses' Board.....	
Special Night Nurses' Board.....	
X-Ray.....	
Physio-Therapy.....	
Guest Meals.....	
Out Patient Dept.....	
Telephone Tolls.....	
Ambulance.....	
TOTAL	<u>61 60</u>

TELEPHONE PIEDMONT 1920
 OAKLAND, CALIF. Feb. 25 1928
 M. [redacted]
 IN ACCOUNT WITH
OAKLAND MATERNITY HOSPITAL
 358 THIRTY-FOURTH STREET
 TERMS WEEKLY IN ADVANCE

To Room Feb 21 st - 28 th	<u>\$35.00</u>
To Delivery Room, Dressing etc	<u>10.00</u>
To Medicines	<u>2.50</u>
	<u>\$47.50</u>

Paid Maude G. Barnes
per C.C.


STATEMENT
 Findlay, Ohio, Feb. 27 1932
 M. [redacted]
 IN ACCOUNT WITH
The Home and Hospital

<u>Feb. 19 - Feb. 27 1932</u>	
<u>9 days @ 4th day</u>	<u>32 00</u>
<u>Delivery Room</u>	<u>5 00</u>
<u>Nursery</u>	<u>8 00</u>
<u>Mineralgia</u>	<u>2 00</u>
<u>Dressings</u>	<u>2 25</u>
<u>Drug</u>	<u>1 00</u>
	<u>50 25</u>


THE HOME & HOSPITAL
PAID
 FEB 27 1932
 By C. L. Hearn Supt. M. D.

Figure 3: The Expansion of Blue Cross and Blue Shield

Panel A. Number of Counties with Blue Cross and Blue Shield Plans




Panel B. Blue Cross and Blue Shield Enrollment



Notes: Panel A plots the cumulative number of U.S. counties in which Blue Cross and Blue Shield plans were available from 1929 to 1947. Panel B plots total enrollment in Blue Cross and Blue Shield plans over the same period, measured in millions of enrollees. Blue Cross plans provided hospital-service coverage, while Blue Shield plans provided coverage for physician services.

Source: Reed [1947].


Figure 4: Geographic Distribution of Minimum Medical Care Costs, 1935



Notes: The map shows the geographic distribution of minimum medical service costs across U.S. counties in 1935. The measure represents the annual cost of a standardized bundle of minimum medical services for a four-person family, based on expected medical care needs over a lifetime. Because costs are observed for only 59 cities, each county is assigned the value of the geographically nearest city. Darker shading indicates higher medical service costs, and crosses mark the 59 cities for which costs are directly observed.

Source: Stecker [1937].


Figure 5: Timing of Blue Cross Hospital Plan Entry Across Counties



Notes: The map shows the first year in which Blue Cross hospital plan coverage became available in each U.S. county. Counties are shaded according to the year of initial Blue Cross entry, with colors corresponding to entry years from 1932 to 1947. White counties had no Blue Cross hospital plan coverage by 1947.


Source: Reed [1947].

Figure 6: The Effect of Blue Cross Hospital Plans on Enrollment Share




Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on Blue Cross enrollment share. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. Enrollment share is measured on a 0–100 percentage scale. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure 7: The Effect of Blue Cross Hospital Plans on Hospital Births per 100 Women aged 15–44




Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on the hospital birth rate. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. The hospital birth rate is measured as hospital births per 100 women aged 15–44. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure 8: The Effect of Blue Cross Hospital Plans on Non-Hospital Births per 100 Women aged 15–44



Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on the non-hospital birth rate. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. The non-hospital birth rate is defined as total births minus hospital births, divided by the number of women aged 15–44 and multiplied by 100. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.


Figure 9: The Effect of Blue Cross Hospital Plans on Births per 100 Women aged 15–44




Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on the total birth rate. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. The birth rate is measured as total births per 100 women ages 15–44. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure 10: Heterogeneous Effects of Blue Cross Hospital Plans on Hospital Births per 100 Women aged 15–44 by Baseline Medical Costs

Panel A: High Baseline Medical Costs




Panel B: Low Baseline Medical Costs




Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on the hospital birth rate separately for counties with high and low baseline medical costs. Counties are classified as high or low cost based on whether minimum medical service costs are above or below the sample mean. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. The hospital birth rate is measured as hospital births per 100 women aged 15–44. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure 11: Heterogeneous Effects of Blue Cross Hospital Plans on Births per 100 Women aged 15–44 by Baseline Medical Costs

Panel A: High Baseline Medical Costs




Panel B: Low Baseline Medical Costs




Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on the total birth rate separately for counties with high and low baseline medical costs. Counties are classified as high or low cost based on whether minimum medical service costs are above or below the sample mean. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. The birth rate is measured as total births per 100 women aged 15–44. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure 12: The Effect of Blue Cross Hospital Plans on Maternal Mortality per 100,000 Births



Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on the maternal mortality rate. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. The maternal mortality rate is measured as maternal deaths per 100,000 live births. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure 13: The Effect of Blue Cross Hospital Plans on Infant Mortality per 100 Births



Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on the infant mortality rate. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. The infant mortality rate is measured as infant deaths per 100 live births. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

A Additional Tables and Figures

Table A1: The Effect of Blue Cross Hospital Plans on Enrollment: County-Level Estimates

	Average Pre	Average Post
Enrollment Share	0.005 (0.02)	7.518*** (0.968)

Notes: The table summarizes county-level event-study estimates of the effect of Blue Cross hospital plan entry on Blue Cross enrollment share. The Average Pre column reports the average of the event-study coefficients for event years -5 through -2, with event year -1, the year immediately before entry, omitted as the reference period. The Average Post column reports the average of the event-study coefficients for event years 0 through 9. Enrollment share is measured on a 0–100 percentage scale. Standard errors are reported in parentheses. *** $p < 0.01$, ** $p < 0.05$, and * $p < 0.10$.

Table A2: The Effect of Blue Cross Hospital Plans on Birth Rates and Birth Counts by Hospital Availability

	Has a Hospital	No Hospital	Difference
<i>Panel A. Birth Rate</i>			
Average Pre	0.098 (0.123)	0.033 (0.271)	
Average Post	0.385 (0.339)	0.074 (0.268)	0.311 (0.288)
Pre-treatment Outcome Mean	8.585	9.378	
<i>Panel B. Birth Counts</i>			
Average Pre	33.464 (30.372)	1.441 (9.086)	
Average Post	404.348*** (111.887)	1.990 (8.805)	402.357*** (110.936)
Pre-treatment Outcome Mean	764.736	281.152	

Notes: The table reports event-study estimates of the effect of Blue Cross hospital plan entry separately for counties with and without a hospital at baseline. The Average Pre estimates average the event-study coefficients for event years -5 through -2, with event year -1, the year immediately before entry, omitted as the reference period. The Average Post estimates average the event-study coefficients for event years 0 through 9. In Panel A, the outcome is the birth rate, measured as births per 100 women aged 15–44. In Panel B, the outcome is the total number of births in a county-year. The Difference column reports the difference in average post-treatment effects between counties with and without a hospital (Has a Hospital minus No Hospital). Pre-treatment outcome means are calculated using observations from the pre-treatment period. Standard errors are reported in parentheses. *** $p < 0.01$, ** $p < 0.05$, and * $p < 0.10$.

Table A3: The Effect of Blue Cross Hospital Plans on Birth Counts

	Average Pre	Average Post
Hospital Births	-89.704* (49.11)	321.658** (127.825)
Pre-treatment Outcome Mean	138.936	
Non-Hospital Births	72.534*** (21.937)	-94.227** (38.017)
Pre-treatment Outcome Mean	406.915	
Total Births	29.63 (30.868)	296.878*** (101.348)
Pre-treatment Outcome Mean	545.426	

- Notes: The table summarizes event-study estimates of the effect of Blue Cross hospital plan entry on county-level birth counts. The Average Pre column reports the average of the event-study coefficients for event years -5 through -2, with event year -1, the year immediately before entry, omitted as the reference period. The Average Post column reports the average of the event-study coefficients for event years 0 through 9. Hospital births are the number of births occurring in hospitals, non-hospital births are calculated as total births minus hospital births, and total births are the total number of births in the county-year. Pre-treatment outcome means are calculated using observations from the pre-treatment period. Standard errors are reported in parentheses. *** $p < 0.01$, ** $p < 0.05$, and * $p < 0.10$.

Table A4: The Effect of Blue Cross Hospital Plans by Baseline Medical Costs: Additional Outcomes

	High Baseline Medical Costs	Low Baseline Medical Costs	Difference
<i>Panel A. Hospital Birth Counts</i>			
Average Pre	-133.122* (71.691)	-26.108 (26.683)	—
Average Post	526.946** (203.379)	71.742 (67.141)	455.203** (203.255)
Pre-treatment Outcome Mean	140.995	137.060	—
<i>Panel B. Non-Hospital Birth Rate</i>			
Average Pre	0.112 (0.268)	0.255 (0.161)	—
Average Post	0.181 (0.348)	0.033 (0.233)	0.147 (0.323)
Pre-treatment Outcome Mean	8.154	7.671	—
<i>Panel C. Non-Hospital Birth Counts</i>			
Average Pre	98.711*** (36.198)	54.916 (17.824)	—
Average Post	-137.964** (53.649)	-46.666 (31.364)	-91.297 (58.494)
Pre-treatment Outcome Mean	400.580	416.704	—
<i>Panel D. Total Birth Counts</i>			
Average Pre	20.517 (41.012)	47.035 (23.532)	—
Average Post	499.693*** (157.178)	55.283 (58.339)	444.409*** (165.210)
Pre-treatment Outcome Mean	536.874	559.373	—

Notes: The table reports event-study estimates of the effect of Blue Cross hospital plan entry separately for counties with high and low baseline medical costs. Counties are classified as high or low cost based on whether minimum medical service costs are above or below the sample mean. The Average Pre estimates average the event-study coefficients for event years -5 through -2, with event year -1, the year immediately before entry, omitted as the reference period. The Average Post estimates average the event-study coefficients for event years 0 through 9. In Panel A, the outcome is the number of hospital births in a county-year. In Panel B, the non-hospital birth rate measures births occurring outside hospitals per 100 women ages 15–44 and is constructed as the difference between the total birth rate and the hospital birth rate. In Panels C and D, the outcomes are the number of non-hospital births and total births, respectively, in a county-year. The Difference column reports the difference in average post-treatment effects between high- and low-cost counties (High minus Low). Pre-treatment outcome means are calculated using observations from the pre-treatment period. Standard errors are reported in parentheses. *** $p < 0.01$, ** $p < 0.05$, and * $p < 0.10$.

Table A5: The Effect of Blue Cross Hospital Plans on Additional Mortality Outcomes

	Average Pre	Average Post
<i>Panel A: Infant Deaths</i>		
Infant Deaths	6.362*** (2.265)	0.246 (1.767)
Pre-treatment Outcome Mean	32.315	
<i>Panel B: Maternal Mortality Rate</i>		
White	0.320 (52.316)	48.033 (63.181)
Pre-treatment Outcome Mean	857.107	
Black	247.695 (149.542)	-121.634 (200.583)
Pre-treatment Outcome Mean	1320.682	

Notes: The table summarizes event-study estimates of the effect of Blue Cross hospital plan entry on additional mortality outcomes. Panel A reports the number of infant deaths at the county-year level. The Average Pre estimate averages event-study coefficients for event years -5 through -2, with event year -1, the year immediately before entry, omitted as the reference period; the Average Post estimate averages coefficients for event years 0 through 9. Panel B reports maternal mortality rates separately for White and Black mothers, measured as maternal deaths per 100,000 live births. For maternal mortality, the Average Pre estimate averages coefficients for event years -3 and -2, with event year -1 omitted as the reference period, and the Average Post estimate averages coefficients for event years 0 through 4. Pre-treatment outcome means are calculated using observations from the corresponding pre-treatment period. Standard errors are reported in parentheses. *** $p < 0.01$, ** $p < 0.05$, and * $p < 0.10$.

Table A6: The Effect of Blue Cross Hospital Plans on Hospital Capacity

	Average Pre	Average Post	Pre-treatment Mean
<i>Panel A: Bassinets</i>			
Bassinets per 100 Women	0.002 (0.003)	0.001 (0.005)	0.097
Bassinets	0.255 (1.240)	6.854** (2.602)	9.977
<i>Panel B: Hospital Beds</i>			
Beds per Capita	-0.033** (0.016)	0.073*** (0.026)	0.462
Beds	-31.933 (27.392)	145.038** (55.740)	211.334
<i>Panel C: General Hospitals</i>			
Has a General Hospital	-0.001 (0.012)	-0.008 (0.022)	0.546
General Hospitals per Capita	-0.0002 (0.0001)	0.0002 (0.0003)	0.004
General Hospitals	0.043 (0.048)	0.051 (0.047)	1.060


Notes: The table summarizes event-study estimates of the effect of Blue Cross hospital plan entry on hospital capacity. The Average Pre column reports the average of the pre-treatment event-study coefficients, excluding event year -1, the year immediately before entry, which is omitted as the reference period. The Average Post column reports the average of the post-treatment event-study coefficients. In Panel A, bassinets per 100 women measures the number of hospital bassinets per 100 women aged 15–44, while Bassinets reports the total number of bassinets in the county-year. In Panel B, Beds per Capita measures hospital beds relative to population, while Beds reports the total number of hospital beds in the county-year. In Panel C, Has a General Hospital is an indicator for the presence of at least one general hospital, General Hospitals per Capita measures the number of general hospitals relative to population, and General Hospitals reports the total number of general hospitals in the county-year. Pre-treatment means are calculated using observations from the pre-treatment period. Standard errors are reported in parentheses. *** $p < 0.01$, ** $p < 0.05$, and * $p < 0.10$.

Table A7: The Effect of Blue Cross Hospital Plans on Hospital Utilization and Mortality

	Average Pre	Average Post	Pre-treatment Mean
<i>Panel A: Hospital Utilization</i>			
Average Daily Census per Capita	-0.023* (0.014)	0.068*** (0.024)	0.354
Average Daily Census	-45.767* (21.128)	121.474** (47.568)	169.014
<i>Panel B: Hospital Mortality</i>			
Mortality Rate	0.013 (0.016)	0.003 (0.022)	1.009
Deaths	-17.075 (30.451)	33.218 (33.066)	324.267


Notes: The table summarizes event-study estimates of the effect of Blue Cross hospital plan entry on hospital utilization and hospital mortality. The Average Pre column reports the average of the pre-treatment event-study coefficients, excluding event year -1 , the year immediately before entry, which is omitted as the reference period. The Average Post column reports the average of the post-treatment event-study coefficients. In Panel A, Average Daily Census measures the average number of patients hospitalized on a given day, while Average Daily Census per Capita scales this measure by county population. In Panel B, Mortality Rate is calculated as the number of deaths divided by county population, while Deaths reports the total number of deaths in the county-year. Pre-treatment means are calculated using observations from the pre-treatment period. Standard errors are reported in parentheses. *** $p < 0.01$, ** $p < 0.05$, and * $p < 0.10$.

Figure A1: Blue Cross Enrollment Growth Following the First Year of Enrollment




Notes: The figure plots average Blue Cross enrollment share by year relative to the first year of Blue Cross enrollment. Event year 0 denotes the first year of enrollment, while negative event years denote years prior to enrollment and positive event years denote years following enrollment. Enrollment share is measured on a 0–100 percentage scale.

Figure A2: Correlation Between Births by Residence and Births by Occurrence




Notes: The figure plots the annual correlation between county-level births measured by place of residence and births measured by place of occurrence from 1937 to 1947. Each point represents the correlation coefficient across counties in a given year. Correlation coefficients closer to one indicate a stronger correspondence between the two measures.

Figure A3: The Effect of Blue Cross Hospital Plans on Enrollment Share: County-Level Estimates




Notes: The figure plots county-level event-study estimates of the effect of Blue Cross hospital plan entry on Blue Cross enrollment share. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. Enrollment share is measured on a 0–100 percentage scale. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure A4: The Effect of Blue Cross Hospital Plans on Hospital Birth Counts




Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on the number of hospital births at the county-year level. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure A5: The Effect of Blue Cross Hospital Plans on Non-Hospital Birth Counts



Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on the number of non-hospital births at the county-year level. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. Non-hospital births are calculated as total births minus hospital births. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.


Figure A6: The Effect of Blue Cross Hospital Plans on Total Birth Counts




Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on the total number of births at the county-year level. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure A7: The Effect of Blue Cross Hospital Plans on Births per 100 Women aged 15–44, by Hospital Availability

Panel A. Has a Hospital




Panel B. No Hospital




Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on the birth rate separately for counties with and without a hospital at baseline. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. The birth rate is measured as total births per 100 women ages 15–44. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure A8: Baseline Medical Costs and Monthly Blue Cross Premiums




Notes: Each panel plots city-level monthly Blue Cross premiums against baseline medical service costs. Baseline medical costs are measured using the 1935 cost of a standardized bundle of minimum medical services for a four-person family. Panels A and B report premiums for semiprivate-room coverage under two-person and family plans, respectively; Panels C and D report premiums for ward coverage under two-person and family plans. The solid line shows the fitted linear relationship between baseline medical costs and monthly premiums. The estimated regression coefficient and corresponding p-value are reported in the upper-left corner of each panel.

Figure A9: The Effect of Blue Cross Hospital Plans on Hospital Birth Counts by Baseline Medical Costs




Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on hospital birth counts separately for counties with high and low baseline medical costs. Counties are classified as high or low cost based on whether minimum medical service costs are above or below the sample mean. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. Hospital birth counts are measured as the total number of hospital births in a county-year. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure A10: The Effect of Blue Cross Hospital Plans on Non-Hospital Births per 100 Women aged 15–44 by Baseline Medical Costs




Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on the non-hospital birth rate separately for counties with high and low baseline medical costs. Counties are classified as high or low cost based on whether minimum medical service costs are above or below the sample mean. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. The non-hospital birth rate measures births occurring outside hospitals per 100 women ages 15–44 and is constructed as the difference between the total birth rate and the hospital birth rate. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure A11: The Effect of Blue Cross Hospital Plans on Non-Hospital Birth Counts by Baseline Medical Costs



Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on non-hospital birth counts separately for counties with high and low baseline medical costs. Counties are classified as high or low cost based on whether minimum medical service costs are above or below the sample mean. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. Non-hospital birth counts are calculated as total births minus hospital births at the county-year level. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.


Figure A12: The Effect of Blue Cross Hospital Plans on Total Birth Counts by Baseline Medical Costs




Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on total birth counts separately for counties with high and low baseline medical costs. Counties are classified as high or low cost based on whether minimum medical service costs are above or below the sample mean. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. Total birth counts are measured as the total number of births in a county-year. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure A13: The Effect of Blue Cross Hospital Plans on Maternal Mortality Rates by Race

Panel A. White




Panel B. Black



Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on maternal mortality rates separately for White and Black mothers. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. Maternal mortality rates are measured as maternal deaths per 100,000 live births. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.


Figure A14: The Effect of Blue Cross Hospital Plans on Infant Deaths




Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on the number of infant deaths at the county-year level. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure A15: The Effect of Blue Cross Hospital Plans on Hospital Capacity


Panel A: Bassinets per 100 Women aged 15–44




Panel B: Bassinets




Panel C: Beds per Capita




Panel D: Beds




Panel E: Has a General Hospital



Panel F: General Hospitals per Capita




Panel G: General Hospitals



Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on measures of hospital capacity. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. Panels A and B report the number of hospital bassinets, scaled per 100 women ages 15–44 and in levels, respectively. Panels C and D report hospital beds per capita and the total number of hospital beds. Panels E–G report an indicator for the presence of at least one general hospital, the number of general hospitals per capita, and the total number of general hospitals, respectively. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

Figure A16: The Effect of Blue Cross Hospital Plans on Hospital Utilization and Mortality



Notes: The figure plots event-study estimates of the effect of Blue Cross hospital plan entry on hospital utilization and mortality. Event time is measured relative to the first year of Blue Cross entry, with event year -1 , the year immediately before entry, omitted as the reference period. Panels A and B report average daily census, scaled by county population and in levels, respectively. Average daily census measures the average number of patients hospitalized on a given day. Panel C reports the mortality rate, calculated as the number of deaths divided by county population. Panel D reports the total number of deaths in the county-year. Points indicate estimated coefficients, and vertical bars show 95 percent confidence intervals.

B Detailed Conceptual Framework

B.1 Household Problem

I develop a simple static framework in which households choose whether to attempt conception, while a live birth is realized probabilistically. This distinction separates the behavioral fertility decision from the live-birth outcome observed in the data.

Consider household i in period t . Let

$$a_{it} \in \{0, 1\} \tag{1}$$

denote whether the household attempts conception, where $a_{it} = 1$ indicates that the household chooses to try to conceive. Conditional on attempting conception, a live birth occurs with probability $p_{it} \in (0, 1)$. Let $\xi_{it} \in \{0, 1\}$ indicate whether an attempt results in a live birth:

$$\xi_{it} \sim \text{Bernoulli}(p_{it}) \tag{2}$$

The realized live-birth outcome is therefore

$$b_{it} = a_{it}\xi_{it} \tag{3}$$

Thus, $\Pr(b_{it} = 1 \mid a_{it} = 1) = p_{it}$ and $\Pr(b_{it} = 1 \mid a_{it} = 0) = 0$.

Households differ in their underlying propensity to attempt conception. Let ε_{it} denote an idiosyncratic utility shifter associated with attempting conception, observed by the household before the attempt decision, with

$$\varepsilon_{it} \sim F \tag{4}$$

where F is a continuously differentiable cumulative distribution function with density f . I assume that ε_{it} is independent of insurance status and of the biological realization ξ_{it} , conditional on the household's economic environment.

Let N_{it} denote the number of surviving children at the beginning of the period. Conditional on a live birth, let $z_{it} \in \{0, 1\}$ denote infant survival, where q_{it} is the probability that a live-born infant survives:

$$z_{it} \sim \text{Bernoulli}(q_{it}) \tag{5}$$

If the infant survives, the household has $N_{it} + 1$ surviving children; otherwise it remains at N_{it}

surviving children at the end of the period.

Household utility depends on consumption, the number of surviving children, and female non-market labor time. A live birth also imposes a maternal health cost. Conditional on the attempt decision and realized birth outcome, household utility is

$$u(c_{it}, N_{it} + b_{it}z_{it}, h_{it}) - b_{it}R(H_{it}) + a_{it}\varepsilon_{it} \quad (6)$$

where $u_c > 0$ and $u_N > 0$. The term $R(H_{it})$ captures the health cost associated with pregnancy and childbirth, where H_{it} denotes maternal health.

Insurance status is determined prior to the conception-attempt decision. Let $I_{it} \in \{0, 1\}$ denote insurance status and let Π_{it} denote premium expenditure over the period conditional on being insured. Actual premium expenditure is therefore $I_{it}\Pi_{it}$ and is incurred regardless of whether a birth subsequently occurs.

Conditional on conception-attempt choice $a \in \{0, 1\}$ and realized live-birth outcome $j \in \{0, 1\}$, define household value as

$$V_{it}^{a,j} = \max_{c_{it}, L_{it}, h_{it}} \{E_t[u(c_{it}, N_{it} + jz_{it}, h_{it})] - jR(H_{it}) + a\varepsilon_{it}\} \quad (7)$$

subject to

$$c_{it} = y_{it} + w(H_{it})L_{it} - I_{it}\Pi_{it} - jP_{it}(I_{it}, C_i) \quad (8)$$

and

$$L_{it} + h_{it} + \ell j = 1 \quad (9)$$

Here, y_{it} denotes non-wife labor household income, L_{it} female labor supply, h_{it} wife's non-market time, $w(H_{it})$ the wage as a function of maternal health, C_i the baseline uninsured cost of childbirth in the local market, $P_{it}(I_{it}, C_i)$ the out-of-pocket childbirth expense, and ℓ the time cost associated with childbirth.

Let $\lambda_{it}^{a,j} > 0$ denote the multiplier on the budget constraint conditional on (a, j) . By the envelope theorem, $\lambda_{it}^{a,j}$ is the marginal value of household resources in that state. Because the attempt-specific utility shifter is additive, it affects the decision to attempt conception but does not alter the optimal allocation of consumption, labor supply, or non-market time conditional on a realized birth outcome.

The household chooses whether to attempt conception before the live-birth outcome is realized. Let W_{it}^a denote the household's expected value associated with conception-attempt choice a , evaluated before the live-birth outcome is realized. If the household does not attempt conception, then $b_{it} = 0$ with certainty and

$$W_{it}^0 = V_{it}^{0,0} \quad (10)$$

If the household attempts conception, a live birth occurs with probability p_{it} , so

$$W_{it}^1 = p_{it}V_{it}^{1,1} + (1 - p_{it})V_{it}^{1,0} \quad (11)$$

Define the relative value of attempting conception as

$$\Delta_{it} \equiv W_{it}^1 - W_{it}^0 \quad (12)$$

Because the states $(a, j) = (1, 0)$ and $(a, j) = (0, 0)$ have identical budget and time constraints and differ only by the additive attempt-specific utility shifter,

$$V_{it}^{1,0} - V_{it}^{0,0} = \varepsilon_{it} \quad (13)$$

It follows that

$$\Delta_{it} = p_{it} \left(V_{it}^{1,1} - V_{it}^{1,0} \right) + \varepsilon_{it} \quad (14)$$

Define the systematic component of the relative value of attempting conception as

$$\bar{\Delta}_{it} \equiv p_{it} \left(V_{it}^{1,1} - V_{it}^{1,0} \right) \quad (15)$$

The household attempts conception when

$$a_{it}^* = \mathbf{1} \{ \bar{\Delta}_{it} + \varepsilon_{it} \geq 0 \} \quad (16)$$

Thus,

$$\Pr(a_{it}^* = 1) = 1 - F(-\bar{\Delta}_{it}) \quad (17)$$

Since a live birth occurs with probability p_{it} conditional on attempting conception,

$$\Pr(b_{it} = 1) = p_{it} [1 - F(-\bar{\Delta}_{it})] \quad (18)$$

For a variable x that affects household value but does not directly affect p_{it} ,

$$\frac{\partial \Pr(b_{it} = 1)}{\partial x} = p_{it}^2 f(-\bar{\Delta}_{it}) \frac{\partial \left(V_{it}^{1,1} - V_{it}^{1,0} \right)}{\partial x} \quad (19)$$

Hence, for households with positive density around the conception threshold, the direction of the

effect on realized live births is determined by the direction of the effect on the value of the live-birth state relative to the no-live-birth state, conditional on attempting conception. The following sections consider how insurance changes this relative value through three channels.

B.2 Financial Channel

The financial channel operates through the way insurance changes household resources across the live-birth and no-live-birth states. Let $C_i > 0$ denote the baseline uninsured cost of childbirth. Suppose insurance covers a fraction $\kappa \in (0, 1)$ of this cost:

$$P_{it}(I_{it}, C_i) = (1 - \kappa I_{it})C_i \quad (20)$$

Insurance therefore reduces the out-of-pocket expense associated with a live birth by

$$P_{it}(0, C_i) - P_{it}(1, C_i) = \kappa C_i \quad (21)$$

This reduction is realized only in the live-birth state, whereas the premium $I_{it}\Pi_{it}$ is paid regardless of whether a live birth occurs.

Holding maternal health and infant survival fixed, the envelope theorem gives, conditional on attempting conception,

$$\left. \frac{\partial V_{it}^{1,j}}{\partial I_{it}} \right|_{\text{Financial}} = \lambda_{it}^{1,j} \left[-\Pi_{it} - j \frac{\partial P_{it}}{\partial I_{it}} \right], \quad j \in \{0, 1\} \quad (22)$$

Since $\partial P_{it}/\partial I_{it} = -\kappa C_i$,

$$\left. \frac{\partial V_{it}^{1,1}}{\partial I_{it}} \right|_{\text{Financial}} = \lambda_{it}^{1,1} (-\Pi_{it} + \kappa C_i) \quad (23)$$

$$\left. \frac{\partial V_{it}^{1,0}}{\partial I_{it}} \right|_{\text{Financial}} = -\lambda_{it}^{1,0} \Pi_{it} \quad (24)$$

The financial effect of insurance on the value of the live-birth state relative to the no-live-birth state is therefore

$$\left. \frac{\partial (V_{it}^{1,1} - V_{it}^{1,0})}{\partial I_{it}} \right|_{\text{Financial}} = \lambda_{it}^{1,1} \kappa C_i - \Pi_{it} (\lambda_{it}^{1,1} - \lambda_{it}^{1,0}) \quad (25)$$

$$= \lambda_{it}^{1,1} (\kappa C_i - \Pi_{it}) + \Pi_{it} \lambda_{it}^{1,0} \quad (26)$$

Because $\lambda_{it}^{1,1} > 0$, $\lambda_{it}^{1,0} > 0$, and $\Pi_{it} \geq 0$, a sufficient condition for this effect to be positive is

$$\kappa C_i \geq \Pi_{it} \quad (27)$$

Under this condition, insurance raises the relative value of the live-birth state. Since $p_{it} > 0$ and is assumed not to depend on insurance,

$$\left. \frac{\partial \bar{\Delta}_{it}}{\partial I_{it}} \right|_{\text{Financial}} = p_{it} \left[\lambda_{it}^{1,1} \kappa C_i - \Pi_{it} \left(\lambda_{it}^{1,1} - \lambda_{it}^{1,0} \right) \right] > 0 \quad (28)$$

Using equation (19), the financial component of the effect on the probability of a realized live birth is

$$\left. \frac{\partial \Pr(b_{it} = 1)}{\partial I_{it}} \right|_{\text{Financial}} = p_{it}^2 f(-\bar{\Delta}_{it}) \left[\lambda_{it}^{1,1} \kappa C_i - \Pi_{it} \left(\lambda_{it}^{1,1} - \lambda_{it}^{1,0} \right) \right] > 0 \quad (29)$$

for households with positive density around the conception threshold. Thus, when condition (27) holds, insurance increases the incentive to attempt conception and the probability of a realized live birth.

B.2.1 Heterogeneity by Baseline Childbirth Costs

The financial channel also generates a prediction for heterogeneity across markets with different baseline childbirth costs. For a given coverage rate κ , equation (21) shows that insurance removes a larger dollar expense where childbirth is more costly.

Define the financial effect on the systematic value of attempting conception as

$$G(C_i) \equiv \left. \frac{\partial \bar{\Delta}_{it}}{\partial I_{it}} \right|_{\text{Financial}} = p_{it} \left[\lambda_{it}^{1,1}(C_i) \kappa C_i - \Pi_{it} \left(\lambda_{it}^{1,1}(C_i) - \lambda_{it}^{1,0} \right) \right] \quad (30)$$

Baseline childbirth costs enter the household budget only in the live-birth state, so, conditional on the premium, $\lambda_{it}^{1,0}$ does not depend on C_i . In the live-birth state, a higher C_i reduces household resources. Under concavity of the conditional value function in resources,

$$\frac{\partial \lambda_{it}^{1,1}(C_i)}{\partial C_i} \geq 0 \quad (31)$$

To see this, let $m_{it}^{1,1} = y_{it} - I_{it}\Pi_{it} - (1 - \kappa I_{it})C_i$ denote non-labor resources available in the live-birth state. Since $\lambda_{it}^{1,1} = \partial V_{it}^{1,1} / \partial m_{it}^{1,1}$, concavity implies $\partial^2 V_{it}^{1,1} / \partial (m_{it}^{1,1})^2 \leq 0$, while $\partial m_{it}^{1,1} / \partial C_i < 0$. Hence equation (31) follows.

Holding p_{it} , κ , and Π_{it} fixed with respect to C_i , differentiating equation (30) gives

$$\frac{\partial G(C_i)}{\partial C_i} = p_{it} \left[\kappa \lambda_{it}^{1,1}(C_i) + (\kappa C_i - \Pi_{it}) \frac{\partial \lambda_{it}^{1,1}(C_i)}{\partial C_i} \right] \quad (32)$$

Under condition (27), the second term is nonnegative and the first is strictly positive. Therefore,

$$\frac{\partial G(C_i)}{\partial C_i} > 0 \quad (33)$$

Thus, under these conditions, insurance has a larger financial effect on the incentive to attempt conception where baseline childbirth costs are higher.

To translate this prediction into the realized live-birth outcome observed in the data, recall that

$$\left. \frac{\partial \Pr(b_{it} = 1)}{\partial I_{it}} \right|_{\text{Financial}} = p_{it} f(-\bar{\Delta}_{it}) G(C_i) \quad (34)$$

The response of realized births therefore depends both on the effect of insurance on the incentive to attempt conception, $G(C_i)$, and on the share of households near the conception-attempt threshold. If this share is comparable across markets, the larger financial incentive in higher-cost markets translates into a larger increase in realized births.

B.3 Maternal Health Channel

The maternal-health channel operates through the effect of insurance on maternal health. Holding the financial and infant-survival channels fixed, suppose

$$H_{it} = H(I_{it}), \quad H_I \equiv \frac{\partial H_{it}}{\partial I_{it}} > 0 \quad (35)$$

Maternal health affects household value through two components. Better maternal health reduces the health cost of pregnancy and childbirth,

$$R_H(H_{it}) < 0 \quad (36)$$

and raises female labor-market productivity,

$$w_H(H_{it}) > 0 \quad (37)$$

Let $L_{it}^{1,j}$ denote optimal female labor supply conditional on attempting conception and realized birth outcome j . By the envelope theorem,

$$\frac{\partial V_{it}^{1,j}}{\partial H_{it}} = -jR_H(H_{it}) + \lambda_{it}^{1,j}w_H(H_{it})L_{it}^{1,j}, \quad j \in \{0, 1\} \quad (38)$$

Therefore,

$$\frac{\partial (V_{it}^{1,1} - V_{it}^{1,0})}{\partial H_{it}} = -R_H(H_{it}) + w_H(H_{it}) [\lambda_{it}^{1,1}L_{it}^{1,1} - \lambda_{it}^{1,0}L_{it}^{1,0}] \quad (39)$$

The first component is positive because better maternal health reduces the health cost incurred in the live-birth state. The second component captures the labor-market effect. Although $w_H(H_{it}) > 0$, its sign is ambiguous because the value of the wage increase depends on labor supply and the marginal value of resources in the two birth states.

The maternal-health effect of insurance on the systematic value of attempting conception is therefore

$$\left. \frac{\partial \bar{\Delta}_{it}}{\partial I_{it}} \right|_{\text{Maternal}} = p_{it} \left\{ -R_H(H_{it}) + w_H(H_{it}) [\lambda_{it}^{1,1}L_{it}^{1,1} - \lambda_{it}^{1,0}L_{it}^{1,0}] \right\} H_I \geq 0 \quad (40)$$

Using equation (19), the same sign carries through to the probability of a realized live birth:

$$\left. \frac{\partial \Pr(b_{it} = 1)}{\partial I_{it}} \right|_{\text{Maternal}} = p_{it}^2 f(-\bar{\Delta}_{it}) \left\{ -R_H(H_{it}) + w_H(H_{it}) [\lambda_{it}^{1,1}L_{it}^{1,1} - \lambda_{it}^{1,0}L_{it}^{1,0}] \right\} H_I \geq 0 \quad (41)$$

Thus, better maternal health lowers the health cost of childbirth, which can increase the incentive to attempt conception, but can also raise the opportunity cost of childbearing through improved labor-market productivity. The overall maternal-health effect on fertility is therefore theoretically ambiguous.

B.4 Infant Survival Channel

The infant-survival channel operates through the probability that a live-born infant survives. Suppose

$$q_{it} = q(I_{it}), \quad q_I \equiv \frac{\partial q_{it}}{\partial I_{it}} > 0 \quad (42)$$

Conditional on attempting conception and realizing a live birth,

$$E_t \left[u(c_{it}^{1,1}, N_{it} + z_{it}, h_{it}^{1,1}) \right] = q_{it} u(c_{it}^{1,1}, N_{it} + 1, h_{it}^{1,1}) \quad (43)$$

$$+ (1 - q_{it}) u(c_{it}^{1,1}, N_{it}, h_{it}^{1,1}) \quad (44)$$

By the envelope theorem,

$$\frac{\partial V_{it}^{1,1}}{\partial q_{it}} = u(c_{it}^{1,1}, N_{it} + 1, h_{it}^{1,1}) - u(c_{it}^{1,1}, N_{it}, h_{it}^{1,1}) > 0 \quad (45)$$

where the inequality follows from $u_N > 0$. Conditional on attempting conception but realizing no live birth, infant survival is irrelevant, so $\partial V_{it}^{1,0}/\partial q_{it} = 0$. Hence,

$$\frac{\partial (V_{it}^{1,1} - V_{it}^{1,0})}{\partial q_{it}} = u(c_{it}^{1,1}, N_{it} + 1, h_{it}^{1,1}) - u(c_{it}^{1,1}, N_{it}, h_{it}^{1,1}) > 0 \quad (46)$$

The corresponding effect of insurance on the systematic value of attempting conception is

$$\left. \frac{\partial \bar{\Delta}_{it}}{\partial I_{it}} \right|_{\text{Infant}} = p_{it} \left[u(c_{it}^{1,1}, N_{it} + 1, h_{it}^{1,1}) - u(c_{it}^{1,1}, N_{it}, h_{it}^{1,1}) \right] q_I > 0 \quad (47)$$

Using equation (19),

$$\left. \frac{\partial \Pr(b_{it} = 1)}{\partial I_{it}} \right|_{\text{Infant}} = p_{it}^2 f(-\bar{\Delta}_{it}) \left[u(c_{it}^{1,1}, N_{it} + 1, h_{it}^{1,1}) - u(c_{it}^{1,1}, N_{it}, h_{it}^{1,1}) \right] q_I > 0 \quad (48)$$

Thus, if insurance increases infant survival, it raises the expected benefit of attempting conception and therefore increases the probability of a realized live birth.

B.5 Summary

The framework separates the household's behavioral fertility decision from the live-birth outcome observed in the data. Households choose whether to attempt conception according to

$$a_{it}^* = \mathbf{1} \{ \bar{\Delta}_{it} + \varepsilon_{it} \geq 0 \} \quad (49)$$

while a live birth occurs probabilistically conditional on attempting conception. The probability of a realized live birth is

$$\Pr(b_{it} = 1) = p_{it} [1 - F(-\bar{\Delta}_{it})] \quad (50)$$

where

$$\bar{\Delta}_{it} = p_{it} (V_{it}^{1,1} - V_{it}^{1,0}) \quad (51)$$

Insurance affects this systematic value through three channels:

- *Financial channel:* Insurance requires households to pay a premium but reduces the out-of-pocket expense associated with childbirth in the live-birth state. Under the sufficient condition $\kappa C_i \geq \Pi_{it}$, insurance raises the incentive to attempt conception and the probability of a realized live birth. The financial effect is predicted to be larger where baseline childbirth costs

are higher, subject to the conditions discussed above.

- *Maternal-health channel:* Insurance can improve maternal health. Better maternal health lowers the health cost of childbirth but can also raise female labor-market productivity and the opportunity cost of childbearing. The overall effect on fertility is therefore theoretically ambiguous.
- *Infant-survival channel:* Insurance can increase the probability that a live-born infant survives. Because households value an additional surviving child, improved infant survival raises the expected value of attempting conception and predicts a positive fertility response.

These predictions guide the empirical mechanism analysis. Heterogeneity in the effects of Blue Cross by baseline medical costs provides evidence on the financial channel, while effects on maternal and infant mortality provide evidence on whether changes in maternal health or infant survival may account for the fertility response.

C Additional Data Construction

I construct several additional county-level control variables, following Thomasson and Treber [2008], to account for demographic and economic conditions that may be correlated with both the expansion of Blue Cross and fertility outcomes.

To control for county demographic characteristics, I include log total population and the shares of the population that are Black, urban, and foreign-born. Population data come from *Historical, Demographic, Economic, and Social Data: The United States, 1790–2002* (ICPSR 2896) [Haines and Inter-university Consortium for Political and Social Research, 2010]. I use observations from the 1920, 1930, 1940, and 1950 Censuses. For each census year, I calculate the Black, urban, and foreign-born population shares and linearly interpolate these shares between census years. I similarly interpolate total population between census years and use its logarithm as a measure of county size.

To capture county-level economic conditions, I use real retail sales per capita. Retail sales are observed in 1929, 1933, 1935, 1939, and 1948, and I linearly interpolate between observed years to

construct annual values over the sample period. Data for 1929, 1933, 1935, and 1939 come from the *New Deal Studies*, ICPSR file 101199 [Fishback and Kantor, 2018]. The 1948 observation, which provides the endpoint for interpolation through 1947, comes from the *City and County Data Book Consolidated File, County Data 1947–1977*, ICPSR file 7736 [United States Bureau of the Census, 2008]. I adjust retail sales for inflation and divide by the interpolated county population described above to construct real retail sales per capita.