

SOLUTIONS

SINGAPORE POLYTECHNIC 2023/2024 Semester 2 Examination

No.	SOLUTION
A1	<p>Answer: (d)</p> <p>The order of $A_{m \times n} A_{n \times m}^T$ is $m \times m$</p>
A2	<p>Answer: (c)</p> <p>$A \cap B = B$ and $A - B = 1 \rightarrow B = \{3, 6\}, \{3, 9\}, \{6, 9\}$</p> <p>$B \subset C$ and $C \cup A = A \rightarrow C = \{3, 6\}, \{3, 9\}, \{6, 9\}, \{3, 6, 9\}$</p>
A3	Answer: (c)
A4	<p>Answer: (b)</p> <p>Assume moving rightwards = 1 & downwards = 0, we will need 3 '1's and 3 '0's. With 2 missing lines, all possible ways are:</p> <p>111000, 110100, 110010, 110001, 011100, 011010, 011001, 010110, 010101, 001110, 001101, 000111</p> <p>Therefore, there are 12 paths.</p>
A5	<p>Answer: (c)</p> $P(\text{biased} \text{heads}) = \frac{P(\text{biased}) \times P(\text{heads} \text{biased})}{P(\text{biased}) \times P(\text{heads} \text{biased}) + P(\text{fair}) \times P(\text{heads} \text{fair})}$ $= \frac{\frac{1}{3} \times 0.7}{\frac{1}{3} \times 0.7 + \frac{2}{3} \times 0.5} = \frac{7}{17} \approx 0.41$
B1a (i)	$3 \begin{bmatrix} 3 & -2 \\ 4 & 1 \end{bmatrix} + \begin{bmatrix} -2 & 3 \\ 5 & -6 \end{bmatrix} = \begin{bmatrix} 7 & -3 \\ 17 & -3 \end{bmatrix}$
B1a (ii)	$\mathbf{BA}^T = \begin{bmatrix} -2 & 3 \\ 5 & -6 \end{bmatrix} \begin{bmatrix} 3 & 4 \\ -2 & 1 \end{bmatrix} = \begin{bmatrix} -12 & -5 \\ 27 & 14 \end{bmatrix}$

SOLUTIONS

SINGAPORE POLYTECHNIC 2023/2024 Semester 2 Examination

B1b	$\mathbf{CD} = \begin{bmatrix} 3 & 0 & 2 \\ 2 & 0 & -2 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 \\ -1 & \frac{3}{2} & 5 \\ 1 & -\frac{3}{2} & 0 \end{bmatrix} = \begin{bmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$ <p>Since $\mathbf{CD} = 5\mathbf{I}$:</p> $\mathbf{C}^{-1}\mathbf{CD} = 5\mathbf{C}^{-1}\mathbf{I}$ $\Rightarrow \mathbf{D} = 5\mathbf{C}^{-1}$ $\therefore \mathbf{C}^{-1} = \frac{1}{5}\mathbf{D} = \frac{1}{5} \begin{bmatrix} 1 & 1 & 0 \\ -1 & \frac{3}{2} & 5 \\ 1 & -\frac{3}{2} & 0 \end{bmatrix} = \begin{bmatrix} \frac{1}{5} & \frac{1}{5} & 0 \\ -\frac{1}{5} & \frac{3}{10} & 1 \\ \frac{1}{5} & -\frac{3}{10} & 0 \end{bmatrix}$
B2a	$\mathbf{T}_1 = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}; \mathbf{T}_2 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ $\mathbf{C} = \mathbf{T}_2\mathbf{T}_1 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} -1 & 0 & 0 \\ 2 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$
B2b	$\mathbf{P}' = \mathbf{CP} = \begin{bmatrix} -1 & 0 & 0 \\ 2 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & 2 \\ 1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} -1 & 1 & 0 \\ 2 & -3 & -2 \\ 1 & 1 & 1 \end{bmatrix}$
B2c	$\mathbf{T}_1^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}; \mathbf{T}_2^{-1} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ $\mathbf{C}^{-1} = \mathbf{T}_1^{-1}\mathbf{T}_2^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} -1 & 0 & 0 \\ -2 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$
B3a	<p><u>Convert to HEX</u></p> $1100101001.101_2 = 329.A_{16}$ <p><u>Convert to DEC</u></p> $1100101001.101_2 = 329.A_{16} = 3 \times 16^2 + 2 \times 16^1 + 9 \times 16^0 + 10 \times 16^{-1} = 809.625_{10}$

SOLUTIONS

SINGAPORE POLYTECHNIC 2023/2024 Semester 2 Examination

B3b	Integral part:			Fractional part:		
	2	264		2	0.425	
	2	132	0	2	0.85	0
	2	66	0	2	0.7	1
	2	33	0	2	0.4	1
	2	16	1	2	0.8 (rep)	0
	2	8	0	2	0.6	1
	2	4	0	2	0.2	1
	2	2	0	2	0.4	0
	2	1	0	2	0.8 (rep)	0
		0	1	2	0.6	1
				2	0.2	1
				2	0.4	0
	$\therefore 264.425_{10} = 100001000.01\overline{10110}_2$					
	B4a	$U = \{-4, -3, -2, -1, 0, 1, 2, 3, 4, 5\}$ $A = \{-2, -1, 1, 2, 3\}$ $B = \{0, 2\}$				
B4b	$A \cap \overline{B} = \{-2, -1, 1, 3\}$ $\overline{A \cup B} = \{-4, -3, 4, 5\}$					
B4c	<div><div>U</div><div><div><div><div>-4</div><div>-3</div><div>4</div><div>5</div></div><div><div>-2</div><div>-1</div><div>1</div><div>3</div></div><div><div>2</div></div><div><div>0</div></div></div><div><div>A</div><div>B</div></div></div></div>					
B5a (i)	$(p \wedge \neg q) \Rightarrow r \Rightarrow (\mathbf{F} \wedge \neg \mathbf{T}) \Rightarrow \mathbf{F} = (\mathbf{F} \wedge \mathbf{F}) \Rightarrow \mathbf{F} = \mathbf{F} \Rightarrow \mathbf{F} = \mathbf{T}$					
B5a (ii)	$(p \vee r) \Leftrightarrow q \Rightarrow (\mathbf{F} \vee \mathbf{F}) \Leftrightarrow \mathbf{T} = \mathbf{F} \Leftrightarrow \mathbf{T} = \mathbf{F}$					

SOLUTIONS

SINGAPORE POLYTECHNIC 2023/2024 Semester 2 Examination

B5b	<table><tr><th>p</th><th>q</th><th>$p \wedge q$</th><th>$\neg(p \wedge q)$</th><th>$\neg p$</th><th>$\neg q$</th><th>$\neg p \vee \neg q$</th><th>$(\neg(p \wedge q)) \Leftrightarrow (\neg p \vee \neg q)$</th></tr><tr><td>T</td><td>T</td><td>T</td><td>F</td><td>F</td><td>F</td><td>T</td></tr><tr><td>T</td><td>F</td><td>F</td><td>T</td><td>F</td><td>T</td><td>T</td></tr><tr><td>F</td><td>T</td><td>F</td><td>T</td><td>T</td><td>T</td><td>T</td></tr><tr><td>F</td><td>F</td><td>F</td><td>T</td><td>T</td><td>T</td><td>T</td></tr></table>	p	q	$p \wedge q$	$\neg(p \wedge q)$	$\neg p$	$\neg q$	$\neg p \vee \neg q$	$(\neg(p \wedge q)) \Leftrightarrow (\neg p \vee \neg q)$	T	T	T	F	F	F	T	T	F	F	T	F	T	T	F	T	F	T	T	T	T	F	F	F	T	T	T	T
	p	q	$p \wedge q$	$\neg(p \wedge q)$	$\neg p$	$\neg q$	$\neg p \vee \neg q$	$(\neg(p \wedge q)) \Leftrightarrow (\neg p \vee \neg q)$																													
	T	T	T	F	F	F	T																														
	T	F	F	T	F	T	T																														
	F	T	F	T	T	T	T																														
	F	F	F	T	T	T	T																														
$(\neg(p \wedge q)) \Leftrightarrow (\neg p \vee \neg q)$ is a tautology.																																					
B6a	(i) $16! = 2.09 \times 10^{13}$ (ii) $8! \times 2! = 80640$																																				
B6b	(i) $\left\lfloor \frac{800}{3} \right\rfloor - \left\lfloor \frac{150}{3} \right\rfloor + 1 = 266 - 50 + 1 = 217$ (ii) $\left\lfloor \frac{800}{6} \right\rfloor - \left\lfloor \frac{150}{6} \right\rfloor + 1 = 133 - 25 + 1 = 109$ (iii) $ A_3 \cup A_6 = A_3 = 217$																																				
B7a	$\frac{4}{12} \times \frac{3}{11} \times \frac{2}{10} = \frac{1}{55} \approx 0.0182$																																				
B7b	$1 - \frac{1}{55} = \frac{54}{55} \approx 0.982$																																				
B7c	Only 1 combination to give sum of 10: 3, 3, 4. $P(3,3,4) = \frac{8}{12} \times \frac{7}{11} \times \frac{4}{10} = \frac{28}{165} \approx 0.170$																																				
B7d	$P(\text{sum} = 10 \cap \geq 1 \text{ black}) = P(\text{sum} = 10) - P(\text{sum} = 10 \cap \text{all white})$ $= \frac{28}{165} - \frac{2}{12} \times \frac{1}{11} \times \frac{2}{10}$ $= \frac{1}{6} \approx 0.167$																																				

SOLUTIONS

SINGAPORE POLYTECHNIC 2023/2024 Semester 2 Examination

C1a	<table><tr><th>x</th><th>y</th><th>z</th><th>$f(x, y, z)$</th></tr><tr><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>1</td><td>0</td></tr><tr><td>0</td><td>1</td><td>0</td><td>0</td></tr><tr><td>0</td><td>1</td><td>1</td><td>1</td></tr><tr><td>1</td><td>0</td><td>0</td><td>1</td></tr><tr><td>1</td><td>0</td><td>1</td><td>1</td></tr><tr><td>1</td><td>1</td><td>0</td><td>1</td></tr><tr><td>1</td><td>1</td><td>1</td><td>1</td></tr></table> $f(x, y, z) = (x + y + z)(x + y + \bar{z})(x + \bar{y} + z)$	x	y	z	$f(x, y, z)$	0	0	0	0	0	0	1	0	0	1	0	0	0	1	1	1	1	0	0	1	1	0	1	1	1	1	0	1	1	1	1	1
x	y	z	$f(x, y, z)$																																		
0	0	0	0																																		
0	0	1	0																																		
0	1	0	0																																		
0	1	1	1																																		
1	0	0	1																																		
1	0	1	1																																		
1	1	0	1																																		
1	1	1	1																																		
C1b	<table><tr><th>p</th><th>q</th><th>pq</th><th>\bar{p}</th><th>\bar{q}</th><th>\overline{pq}</th><th>$p \odot q$</th></tr><tr><td>0</td><td>0</td><td>0</td><td>1</td><td>1</td><td>1</td><td>1</td></tr><tr><td>0</td><td>1</td><td>0</td><td>1</td><td>0</td><td>0</td><td>0</td></tr><tr><td>1</td><td>0</td><td>0</td><td>0</td><td>1</td><td>0</td><td>0</td></tr><tr><td>1</td><td>1</td><td>1</td><td>0</td><td>0</td><td>0</td><td>1</td></tr></table>	p	q	pq	\bar{p}	\bar{q}	\overline{pq}	$p \odot q$	0	0	0	1	1	1	1	0	1	0	1	0	0	0	1	0	0	0	1	0	0	1	1	1	0	0	0	1	
p	q	pq	\bar{p}	\bar{q}	\overline{pq}	$p \odot q$																															
0	0	0	1	1	1	1																															
0	1	0	1	0	0	0																															
1	0	0	0	1	0	0																															
1	1	1	0	0	0	1																															
C1c	$\begin{aligned} & p \odot (p \odot q) \odot q \\ &= p \odot (pq + \overline{pq}) \odot q && \text{Substitution} \\ &= (p \bullet (pq + \overline{pq}) + \overline{p \bullet (pq + \overline{pq})}) \odot q && \text{Substitution} \\ &= (pq + \overline{p \bullet (\overline{pq} \bullet pq)}) \odot q && \text{De Morgan's Laws} \\ &= (pq + \overline{p \bullet (\bar{p} + \bar{q}) \bullet (p + q)}) \odot q && \text{De Morgan's Laws} \\ &= (pq + \overline{pq}) \odot q && \text{Associative+Idempotent Laws} \\ &= q \odot q && \text{Distributive Laws} \\ &= q \bullet q + \bar{q} \bullet \bar{q} && \text{Substitution} \\ &= 1 && \text{Idempotent Laws} \end{aligned}$																																				

SOLUTIONS

SINGAPORE POLYTECHNIC 2023/2024 Semester 2 Examination

C2a	<p>Let $\mathbf{Q} = \begin{bmatrix} q_{11} & q_{12} & q_{13} \\ q_{12} & q_{22} & q_{23} \\ q_{13} & q_{23} & q_{33} \end{bmatrix}$ (since \mathbf{Q} is a symmetric matrix, we have $q_{ij} = q_{ji}$)</p> $\mathbf{PQP}^T = \begin{bmatrix} x & y & z \end{bmatrix} \begin{bmatrix} q_{11} & q_{12} & q_{13} \\ q_{12} & q_{22} & q_{23} \\ q_{13} & q_{23} & q_{33} \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ $= \begin{bmatrix} q_{11}x + q_{12}y + q_{13}z & q_{12}x + q_{22}y + q_{23}z & q_{13}x + q_{23}y + q_{33}z \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ $= q_{11}x^2 + q_{12}xy + q_{13}xz + q_{12}xy + q_{22}y^2 + q_{23}yz + q_{13}xz + q_{23}yz + q_{33}z^2$ $= q_{11}x^2 + q_{22}y^2 + q_{33}z^2 + 2q_{12}xy + 2q_{13}xz + 2q_{23}yz$ <p>Comparing coefficients with the expression $2x^2 + 5y^2 + 3z^2 + 2xy - 2xz + 8yz$, we have $q_{11} = 2$, $q_{22} = 5$, $q_{33} = 3$, $q_{12} = 1$, $q_{13} = -1$ and $q_{23} = 4$.</p> $\therefore \mathbf{Q} = \begin{bmatrix} 2 & 1 & -1 \\ 1 & 5 & 4 \\ -1 & 4 & 3 \end{bmatrix}$														
C2b	<p>From Clue No. 1 & 2, Jack must be in the 4th place with 2B1G before him and 1B1G after him. From Clue No. 3, Zack must be in the 6th place (3B2G before him). From Clue No. 4 & 5, Amanda must be in the 1st place and Emma in the 5th place. From Clue No. 7, Victor must be in the 3rd place. As such, from Clue No. 6, Nick must be in the 2nd place. Therefore, the sequence of escape is</p> <table border="1" data-bbox="261 1346 572 1592"> <thead> <tr> <th>Sequence</th><th>Name</th></tr> </thead> <tbody> <tr> <td>1</td><td>Amanda</td></tr> <tr> <td>2</td><td>Nick</td></tr> <tr> <td>3</td><td>Victor</td></tr> <tr> <td>4</td><td>Jack</td></tr> <tr> <td>5</td><td>Emma</td></tr> <tr> <td>6</td><td>Zack</td></tr> </tbody> </table>	Sequence	Name	1	Amanda	2	Nick	3	Victor	4	Jack	5	Emma	6	Zack
Sequence	Name														
1	Amanda														
2	Nick														
3	Victor														
4	Jack														
5	Emma														
6	Zack														

SOLUTIONS

SINGAPORE POLYTECHNIC 2023/2024 Semester 2 Examination

C3a	<p>Let A_i be the event that a green ball appears for the first time at ith picking. Yasir can only win if i is an even number.</p> $i = 2 \rightarrow P(A_2) = \frac{7}{12} \times \frac{5}{11} = \frac{35}{132}$ $i = 4 \rightarrow P(A_4) = \frac{7}{12} \times \frac{6}{11} \times \frac{5}{10} \times \frac{5}{9} = \frac{35}{396}$ $i = 6 \rightarrow P(A_6) = \frac{7}{12} \times \frac{6}{11} \times \frac{5}{10} \times \frac{4}{9} \times \frac{3}{8} \times \frac{5}{7} = \frac{5}{264}$ $i = 8 \rightarrow P(A_8) = \frac{7}{12} \times \frac{6}{11} \times \frac{5}{10} \times \frac{4}{9} \times \frac{3}{8} \times \frac{2}{7} \times \frac{1}{6} \times \frac{5}{5} = \frac{1}{792}$ <p>Hence, $P(\text{Yasir wins}) = \frac{35}{132} + \frac{35}{396} + \frac{5}{264} + \frac{1}{792} = \frac{37}{99}$.</p>
C3b (i)	<p>Define the following possible paths from n_1 to n_4:</p> $A = (l_1, l_4) \quad B = (l_2, l_5) \quad C = (l_1, l_3, l_5)$ <p>Inclusion-exclusion formula for 3-event scenario:</p> $P(A \cup B \cup C)$ $= P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$ $= p_1 p_4 + p_2 p_5 + p_1 p_3 p_5 - p_1 p_2 p_4 p_5 - p_1 p_3 p_4 p_5 - p_1 p_2 p_3 p_5 + p_1 p_2 p_3 p_4 p_5$
C3b (ii)	$P(A \cup B \cup C)$ $= P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$ $= p_1 p_4 + p_2 p_5 + p_1 p_3 p_5 - p_1 p_2 p_4 p_5 - p_1 p_3 p_4 p_5 - p_1 p_2 p_3 p_5 + p_1 p_2 p_3 p_4 p_5$ $= 0.65^2 \times 2 + 0.65^3 - 0.65^4 \times 3 + 0.65^5$ $= 0.700$