

Section 5.1-5.3 Applications

5.1 Ex1 Suppose you take out \$180,000 for the purchase of a \$200,000 home. Suppose this loan has a 3% annual interest rate compounded monthly and that you pay \$1500 at the end of each month.

- ① How much do you still owe after 1 year.
- ② When are you done with payments?

n = # of months

a_n = amount we still owe in \$

$$a_{n+1} = (1+r) a_n - 1500$$

monthly interest rate $r = \frac{.03}{12}$

$$f(x) = \left(1 + \frac{.03}{12}\right) x - 1500$$

iteration ($f(x)$, 180000, 12) soln to (1)

\$167,225.30

iteration ($f(x)$, 180000, n) < 0 soln to (2)

what is n ? 143

$$\frac{143}{12} = 11.92 \approx 12 \text{ years.}$$

(see 5.1 #61)

Ex 2 5.2 #126

$$d = 10$$

$n = \#$ of quarters

$$r = \frac{.04}{4} = .01 \text{ periodic (quarterly) interest rate}$$

$n=1$ d amt at beginning of quarter

$$n=2 \quad \underline{(1+r)d + d}$$

$$\textcircled{n=3} \quad \underline{(1+r)((1+r)d + d) + d} = (1+r)^2 d + (1+r)d + d$$

$$= d (1 + (1+r) + (1+r)^2)$$

$$= d \sum_{i=1}^n \underline{(1+r)}^{i-1}$$

$$= d \left(\frac{1 - 1.01^n}{1 - 1.01} \right)^{x-1}$$

$$\sum_{i=1}^n \underline{s}^{i-1} = 1 + s + \dots + s^{n-1}$$
$$= \frac{1 - s^n}{(1 - s)}$$

$$1+r = 1+0.01 = 1.01$$

$$= \frac{d (1.01^n - 1)}{.01}$$

a. after n quarters is like the beginning of the $(n+1)$ st quarter, except w/o that payment.

$$\frac{d (1.01^{n+1} - 1)}{.01} - \textcircled{(n)} \cdot d$$

see problem 5.2 #126

Ex3 5.2 #127

Ibuprofen decreases by $\approx 29\%$ /hr in the body

A new dose every $N=8$ hours

How much drug is in the body after 25 hrs?

$$n = 25 = 3 \cdot 8 + 1 \quad (n = m \cdot N + k)$$

$d = 400\text{mg}$ / dose.

$A(n)$ = amount after n hours

<u>n (hours)</u>	<u>amount</u>
0	d
1	dr
\vdots	
7	dr^7
8	$dr^8 + d$
+1 9	$(dr^8 + d)r$
\vdots	
+7 15	$(dr^8 + d)r^7$
16	$(dr^8 + d)r^8 + d$
17	$((dr^8 + d)r^8 + d)r$
$n = 18$	$((dr^8 + d)r^8 + d)r^2$

~~$r = 0.29$~~

$1 - 0.29 = 0.71$

$$\begin{aligned}
 & d r^2 (1 + r^8 + (r^8)^2) \\
 = & d r^2 (1 + s + s^2) \quad s = r^8 \\
 & \quad \quad \quad \downarrow \quad \#3 \quad \#2 \quad \#1 \\
 & \quad \quad \quad +2 \text{ hrs.}
 \end{aligned}$$

$$n = 18 = 2 \cdot 8 + 2 = m \cdot N + k$$

$$\begin{aligned}
 A(n) &= d \cdot r^k (1 + s + \dots + s^m) \\
 &= d \cdot r^k \frac{(1 - s^{m+1})}{1 - s}, \quad s = r^8
 \end{aligned}$$

$$\begin{aligned}
 \xrightarrow{\text{if } k=0} \quad \lim_{m \rightarrow \infty} & \quad d \cdot (1) \cdot \frac{(1-0)}{1-s} \\
 &= \frac{d}{1-s} \\
 &= \frac{400}{1-(.71)^8} \approx 427 \text{ mg} \\
 & \quad \quad \quad \text{steady state}
 \end{aligned}$$

(decreases until the next dose).

If $M = 2$
 $k = 0$
 $m = 12$
 $d = 266 \text{ mg}$

steady state is 536mg if
 266mg is dosed
 every 2 hr.
 (see Mathematica).