# California Pepper Tree, filfil rafieh

#### GENERAL

Origin : Mediterranean,

sub-tropical, tropical

normal growth Vigour

rate

Humidity verv humid. extremely humid

Propagation sowing and

pricking out, cuttings

Maintenance : moderate

#### CONDITIONS

Urban climate : resistant Dessication : vulnerable Stagnant water: vulnerable Irrigation : hiah

Salinity/ppm : high (4500 ppm)

Hardiness : -6°C

#### SHAPE

Type : tree : 9 m-15 m Height Spread : 7 m-15 m Foliage : semi-evergreen

## **FLOWER**

: yellow Colour Size : 5 cm - 15 cm Period : April - May Smell : aroma, moderate, leaf

### **FRUIT**

Type of fruit : drupe(fleshy/

juicy)

Fruit size : 0.5 cm - 0.7 cm

Toxicity

: edible when processed

Schinus molle, known as the Californian Pepper Tree and filfil-rafieh in Arabic, has its area of distribution in the Peruvian Andean desert, Chile, central America and southeast Argentina. It is not related to the pepper plant (Piper nigrum), but the edible, red fruits are similar to peppercorns and the tree is so called, because of their shape and taste. The evergreen to semi-evergreen tree grows to a height of 12 metres. It has an open, umbrella-shaped crown with pendulous branches. The leaves are alternate and pinnate with a lanceolate shape. The white male and female flowers grow on different trees; the bloom is unimpressive, in contrast to the pinkish-red fruits, which hang in clusters of dozens of berries nearly the whole year round. Propagation can be done by seeds, cuttings and suckers. S. molle has an extensive, but flat root system which can be a problem for hard paving in the proximity. Tolerant of most soils, it requires full sun and is droughttolerant, hardy. Foliage may be damaged by frost and also suffer from insects. The Pepper Tree comes from arid regions, but experience in Arriyadh has shown that it cannot withstand too harsh desert conditions. In King Fahd Road, for example, it disappeared after planting, because it needed more humidity and shelter, i.e. no hot, desiccating winds. However, it can be recommended in specific, sheltered situations in urban areas, in parks as a specimen tree and in pedestrian precincts. Attention should be paid to the possible problems of fruit drop litter and wind damage.







