

*Rosemary, Anthos, huss alban, ikil al jabal*

**GENERAL**

Origin	: temperate
Vigour	: normal growth rate
Humidity	: semi-arid, semi-humid, very humid
Propagation	: sowing and pricking out, cuttings
Maintenance	: moderate

**CONDITIONS**

Urban climate	: resistant
Dessication	: resistant
Stagnant water	: vulnerable
Irrigation	: medium
Salinity/ppm	: moderate (3000 ppm)
Hardiness	: -9°C

**SHAPE**

Type	: shrub, sub-shrub
Height	: 0.5 m-1.5 m
Spread	: 1 m-3 m
Foliage	: evergreen

**FLOWER**

Colour	: pale blue, pale white
Size	: 1 cm
Period	: March - May
Smell	: aroma, strong, pleasantly scented, leaf, spicy

**FRUIT**

Type of fruit	: nutlet
Fruit size	: 0.2 cm
Toxicity	: edible, leaf

*Rosmarinus officinalis* is known as Rosemary in English, and huss alban and ikil al jabal in Arabic. It is native to the Middle East, southern and southeastern Europe. Its climate zone is temperate. Rosemary normally grows close to coastlines. The name is based on this habit: 'ros' means dew, 'marinus' stands for ocean. The evergreen, dwarf-like shrub grows between 50 cm and 1.5 metres high. The shape of the shrub is often round, oval to irregular. The linear leaves are a fresh, dark green. The foliage emits an appealing spicy fragrance in its preferred sunny locations. Rosemary is a very tasty herb. The 1-cm, pale-blue, bee-attracting flowers appear in early spring and are arranged raceme-like on the branches. The fruits are brown nutlets. Rosemary requires medium irrigation, with low salinity. Propagation methods include sowing and pricking, as well as by cuttings. Owing to its natural habitat, Rosemary prefers locations with a good micro-climate and semi-humid conditions and is frost tolerant. Rosemary is best planted in groups, or in massed planting, as a groundcover or a low hedge. It is attractive in rock or steppe gardens, as well as in private gardens. In urban areas it can be used in public open spaces, parks, pedestrian precincts and in containers. *Rosmarinus* off. 'Prostratus' is the creeping variety, and it has the same requirements and habits as the main species. The prostrate form is excellent for covering the tops of walls. In the early 1980s (CE), the plant's first use in Arriyadh was in the Diplomatic Quarter.

