

Pseudo-Boolean Optimization

in memory of P.L. Hammer

Endre Boros

RUTCOR, Rutgers University

April 12, 2013

Outline

- 1 Pseudo-Boolean Optimization
 - Pseudo-Boolean Functions
 - Representations and Bounds
 - Persistencies and Autarkies
 - Graph Cut Models
 - Implication Networks
- 2 Results
 - Computational Results

Pseudo-Boolean Optimization (PBO)

Variables and Literals

- **Variables:** $x_1, x_2, \dots, x_n \in \{0, 1\}; \quad V = \{1, 2, \dots, n\}$
- **Negations:** $\bar{x}_i = 1 - x_i \in \{0, 1\}$ for $i = 1, \dots, n$

Pseudo-Boolean Function (PBF):

$$f : \{0, 1\}^n \rightarrow \mathbb{R}$$

$$f(x_1, \dots, x_n) = \sum_{S \subseteq V} a_S \prod_{i \in S} x_i$$

Unconstrained Binary Optimization (PBO)

$$\min_{(x_1, \dots, x_n) \in \{0, 1\}^n} f(x_1, \dots, x_n)$$

Pseudo-Boolean Optimization (PBO)

Variables and Literals

- **Variables:** $x_1, x_2, \dots, x_n \in \{0, 1\}; \quad V = \{1, 2, \dots, n\}$
- **Negations:** $\bar{x}_i = 1 - x_i \in \{0, 1\}$ for $i = 1, \dots, n$

Pseudo-Boolean Function (PBF):

$$f : \{0, 1\}^n \rightarrow \mathbb{R}$$

$$f(x_1, \dots, x_n) = \sum_{S \subseteq V} a_S \prod_{i \in S} x_i$$

Unconstrained Binary Optimization (PBO)

$$\min_{(x_1, \dots, x_n) \in \{0, 1\}^n} f(x_1, \dots, x_n)$$

Pseudo-Boolean Optimization (PBO)

Variables and Literals

- **Variables:** $x_1, x_2, \dots, x_n \in \{0, 1\}; \quad V = \{1, 2, \dots, n\}$
- **Negations:** $\bar{x}_i = 1 - x_i \in \{0, 1\}$ for $i = 1, \dots, n$

Pseudo-Boolean Function (PBF):

$$f : \{0, 1\}^n \rightarrow \mathbb{R}$$

$$f(x_1, \dots, x_n) = \sum_{S \subseteq V} a_S \prod_{i \in S} x_i$$

Unconstrained Binary Optimization (PBO)

$$\min_{(x_1, \dots, x_n) \in \{0, 1\}^n} f(x_1, \dots, x_n)$$

Pseudo-Boolean Optimization

(Hammer and Rudeanu, 1968)

Pseudo-Boolean Optimization (PBO)

Variables and Literals

- **Variables:** $x_1, x_2, \dots, x_n \in \{0, 1\}; \quad V = \{1, 2, \dots, n\}$
- **Negations:** $\bar{x}_i = 1 - x_i \in \{0, 1\}$ for $i = 1, \dots, n$

Pseudo-Boolean Function (PBF):

$$f : \{0, 1\}^n \rightarrow \mathbb{R}$$

$$f(x_1, \dots, x_n) = \sum_{S \subseteq V} a_S \prod_{i \in S} x_i$$

Unconstrained Binary Optimization (PBO)

$$\min_{(x_1, \dots, x_n) \in \{0, 1\}^n} f(x_1, \dots, x_n)$$

Pseudo-Boolean Optimization

(Hammer and Rudeanu, 1968)

Pseudo-Boolean Optimization (PBO)

Variables and Literals

- **Variables:** $x_1, x_2, \dots, x_n \in \{0, 1\}; \quad V = \{1, 2, \dots, n\}$
- **Negations:** $\bar{x}_i = 1 - x_i \in \{0, 1\}$ for $i = 1, \dots, n$

Pseudo-Boolean Function (PBF):

$$f : \{0, 1\}^n \rightarrow \mathbb{R}$$

$$f(x_1, \dots, x_n) = \sum_{S \subseteq V} a_S \prod_{i \in S} x_i$$

Unconstrained Binary Optimization (PBO)

$$\min_{(x_1, \dots, x_n) \in \{0, 1\}^n} f(x_1, \dots, x_n)$$

Pseudo-Boolean Optimization

(Hammer and Rudeanu, 1968)

Outline

- 1 Pseudo-Boolean Optimization
 - Pseudo-Boolean Functions
 - Representations and Bounds
 - Persistencies and Autarkies
 - Graph Cut Models
 - Implication Networks
- 2 Results
 - Computational Results

Representations and Lower Bounds

Posiforms: Nonnegative (except maybe the constant terms) multi-linear polynomials in $2n$ literals $x_1, \bar{x}_1, \dots, x_n, \bar{x}_n$

Representations and Lower Bounds

Posiforms: Nonnegative (except maybe the constant terms) multi-linear polynomials in $2n$ literals $x_1, \bar{x}_1, \dots, x_n, \bar{x}_n$

$$f = -2 - x_1 - x_2 - x_3 + x_1x_2 + x_1x_3 + x_2x_3 \quad \text{quadratic PBF}$$

Representations and Lower Bounds

Posiforms: Nonnegative (except maybe the constant terms) multi-linear polynomials in $2n$ literals $x_1, \bar{x}_1, \dots, x_n, \bar{x}_n$

$$\begin{aligned} f &= -2 - x_1 - x_2 - x_3 + x_1x_2 + x_1x_3 + x_2x_3 && \text{quadratic PBF} \\ &= -5 + \bar{x}_1 + \bar{x}_2 + \bar{x}_3 + x_1x_2 + x_1x_3 + x_2x_3 && \text{quadratic posiform} \end{aligned}$$

Representations and Lower Bounds

Posiforms: Nonnegative (except maybe the constant terms) multi-linear polynomials in $2n$ literals $x_1, \bar{x}_1, \dots, x_n, \bar{x}_n$

f	$=$	$-2 - x_1 - x_2 - x_3 + x_1x_2 + x_1x_3 + x_2x_3$	quadratic PBF
	$=$	$-5 + \bar{x}_1 + \bar{x}_2 + \bar{x}_3 + x_1x_2 + x_1x_3 + x_2x_3$	quadratic posiform
	$=$	$-4 + \bar{x}_3 + \bar{x}_1\bar{x}_2 + x_1x_3 + x_2x_3$	quadratic posiform

Representations and Lower Bounds

Posiforms: Nonnegative (except maybe the constant terms) multi-linear polynomials in $2n$ literals $x_1, \bar{x}_1, \dots, x_n, \bar{x}_n$

f	$= -2 - x_1 - x_2 - x_3 + x_1x_2 + x_1x_3 + x_2x_3$	quadratic PBF
	$= -5 + \bar{x}_1 + \bar{x}_2 + \bar{x}_3 + x_1x_2 + x_1x_3 + x_2x_3$	quadratic posiform
	$= -4 + \bar{x}_3 + \bar{x}_1\bar{x}_2 + x_1x_3 + x_2x_3$	quadratic posiform
	$= -3 + x_1x_2x_3 + \bar{x}_1\bar{x}_2\bar{x}_3$	cubic posiform

Representations and Lower Bounds

Posiforms: Nonnegative (except maybe the constant terms) multi-linear polynomials in $2n$ literals $x_1, \bar{x}_1, \dots, x_n, \bar{x}_n$

f	$= -2 - x_1 - x_2 - x_3 + x_1x_2 + x_1x_3 + x_2x_3$	quadratic PBF
	$= -5 + \bar{x}_1 + \bar{x}_2 + \bar{x}_3 + x_1x_2 + x_1x_3 + x_2x_3$	quadratic posiform
	$= -4 + \bar{x}_3 + \bar{x}_1\bar{x}_2 + x_1x_3 + x_2x_3$	quadratic posiform
	$= -3 + x_1x_2x_3 + \bar{x}_1\bar{x}_2\bar{x}_3$	cubic posiform

Roof Dual Bound: $C_2(f) \leq f$ (Hammer, Hansen and Simeone, 1984)

$C_2(f)$ = largest C s.t. $f = C + \phi$ for some **quadratic posiform** ϕ .

Representations and Lower Bounds

Posiforms: Nonnegative (except maybe the constant terms) multi-linear polynomials in $2n$ literals $x_1, \bar{x}_1, \dots, x_n, \bar{x}_n$

$$\begin{aligned}
 f &= -2 - x_1 - x_2 - x_3 + x_1x_2 + x_1x_3 + x_2x_3 && \text{quadratic PBF} \\
 &= \textcolor{brown}{-5} + \bar{x}_1 + \bar{x}_2 + \bar{x}_3 + x_1x_2 + x_1x_3 + x_2x_3 && \text{quadratic posiform} \\
 &= \textcolor{green}{-4} + \bar{x}_3 + \bar{x}_1\bar{x}_2 + x_1x_3 + x_2x_3 && \text{quadratic posiform} \\
 &= \textcolor{blue}{-3} + x_1x_2x_3 + \bar{x}_1\bar{x}_2\bar{x}_3 && \text{cubic posiform}
 \end{aligned}$$

Roof Dual Bound: $C_2(f) \leq f$ (Hammer, Hansen and Simeone, 1984)

$C_2(f)$ = largest C s.t. $f = C + \phi$ for some **quadratic posiform** ϕ .

Complete Hierarchy of Bounds: (B, Crama and Hammer, 1990)

$$C_2(f) \leq C_3(f) \leq \dots \leq C_n(f) = \min f$$

Representations and Lower Bounds

Posiforms: Nonnegative (except maybe the constant terms) multi-linear polynomials in $2n$ literals $x_1, \bar{x}_1, \dots, x_n, \bar{x}_n$

$$\begin{aligned}
 f &= -2 - x_1 - x_2 - x_3 + x_1x_2 + x_1x_3 + x_2x_3 && \text{quadratic PBF} \\
 &= \textcolor{red}{-5} + \bar{x}_1 + \bar{x}_2 + \bar{x}_3 + x_1x_2 + x_1x_3 + x_2x_3 && \text{quadratic posiform} \\
 &= \textcolor{green}{-4} + \bar{x}_3 + \bar{x}_1\bar{x}_2 + x_1x_3 + x_2x_3 && \text{quadratic posiform} \\
 &= \textcolor{blue}{-3} + x_1x_2x_3 + \bar{x}_1\bar{x}_2\bar{x}_3 && \text{cubic posiform}
 \end{aligned}$$

Roof Dual Bound: $C_2(f) \leq f$ (Hammer, Hansen and Simeone, 1984)

$C_2(f)$ = largest C s.t. $f = C + \phi$ for some **quadratic posiform** ϕ .

Complete Hierarchy of Bounds: (B, Crama and Hammer, 1990)

$$C_2(f) \leq C_3(f) \leq \dots \leq C_n(f) = \min f$$

Recent generalizations

Bisubmodular functions: (Kolmogorov, 2010; Kahl and Strandmark, 2011)

Outline

- 1 Pseudo-Boolean Optimization
 - Pseudo-Boolean Functions
 - Representations and Bounds
 - **Persistencies and Autarkies**
 - Graph Cut Models
 - Implication Networks
- 2 Results
 - Computational Results

Persistencies and Autarkies

Partial assignment: $y \in \{0, 1\}^S$, $S \subseteq V$

Persistencies and Autarkies

Partial assignment: $y \in \{0, 1\}^S$, $S \subseteq V$

y is a **persistency** for a pseudo-Boolean function f if

$$f(x|y) \leq f(x) \quad \forall \quad x \in \{0, 1\}^V.$$

Persistencies and Autarkies

Partial assignment: $y \in \{0, 1\}^S$, $S \subseteq V$

y is a **persistency** for a pseudo-Boolean function f if

$$f(x|y) \leq f(x) \quad \forall \quad x \in \{0, 1\}^V.$$

y is an **autarky** for a posiform ϕ if

$T(y) = 0$ for all terms T of ϕ for which $\text{Var}(T) \cap S \neq \emptyset$.

Persistencies and Autarkies

Partial assignment: $y \in \{0, 1\}^S$, $S \subseteq V$

y is a **persistency** for a pseudo-Boolean function f if

$$f(x|y) \leq f(x) \quad \forall \quad x \in \{0, 1\}^V.$$

y is an **autarky** for a posiform ϕ if

$T(y) = 0$ for all terms T of ϕ for which $\text{Var}(T) \cap S \neq \emptyset$.

$S = \{\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3\}$ and $y = (\mathbf{1}, \mathbf{0}, \mathbf{1})$ is an autarky of the posiform

$$\phi = \mathbf{x}_1 \mathbf{x}_2 + 5 \bar{\mathbf{x}}_1 \mathbf{x}_3 x_6 + 4 \bar{\mathbf{x}}_2 \bar{\mathbf{x}}_3 x_7 + 4 \bar{\mathbf{x}}_1 x_4 + 5 \mathbf{x}_2 x_5 + 6 x_4 x_5$$

Persistencies and Autarkies

Partial assignment: $y \in \{0, 1\}^S$, $S \subseteq V$

y is a **persistency** for a pseudo-Boolean function f if

$$f(x|y) \leq f(x) \quad \forall \quad x \in \{0, 1\}^V.$$

y is an **autarky** for a posiform ϕ if

$T(y) = 0$ for all terms T of ϕ for which $\text{Var}(T) \cap S \neq \emptyset$.

Every autarky of a posiform ϕ is a persistency of the function $f = \phi$.

Persistencies and Autarkies

Partial assignment: $y \in \{0, 1\}^S$, $S \subseteq V$

y is a **persistency** for a pseudo-Boolean function f if

$$f(x|y) \leq f(x) \quad \forall \quad x \in \{0, 1\}^V.$$

y is an **autarky** for a posiform ϕ if

$T(y) = 0$ for all terms T of ϕ for which $\text{Var}(T) \cap S \neq \emptyset$.

Every autarky of a posiform ϕ is a persistency of the function $f = \phi$.

Given y it is

- **easy** to test if y is an autarky for a given posiform ϕ ;
- **hard** to test if y is a persistency for a given PBF f .

Persistencies and Autarkies

Partial assignment: $y \in \{0, 1\}^S$, $S \subseteq V$

y is a **persistency** for a pseudo-Boolean function f if

$$f(x|y) \leq f(x) \quad \forall \quad x \in \{0, 1\}^V.$$

y is an **autarky** for a posiform ϕ if

$T(y) = 0$ for all terms T of ϕ for which $\text{Var}(T) \cap S \neq \emptyset$.

Every autarky of a posiform ϕ is a persistency of the function $f = \phi$.

Given y it is

- **easy** to test if y is an autarky for a given posiform ϕ ;
- **hard** to test if y is a persistency for a given PBF f .

Basic facts about persistencies and autarkies

Every persistency of a function f is an autarky for some posiform ϕ representing f .

Basic facts about persistencies and autarkies

Every persistency of a function f is an autarky for some posiform ϕ representing f .

If f has persistencies $y^1 \in \{0, 1\}^{S_1}$ and $y^2 \in \{0, 1\}^{S_2}$, then it also has a persistency $y^3 \in \{0, 1\}^{S_1 \cup S_2}$.

Basic facts about persistencies and autarkies

Every persistency of a function f is an autarky for some posiform ϕ representing f .

If f has persistencies $y^1 \in \{0, 1\}^{S_1}$ and $y^2 \in \{0, 1\}^{S_2}$, then it also has a persistency $y^3 \in \{0, 1\}^{S_1 \cup S_2}$.

Every posiform ϕ has a unique maximal subset $S = S(\phi)$ for which it has an autarky $y \in \{0, 1\}^S$.

Basic facts about persistencies and autarkies

Every persistency of a function f is an autarky for some posiform ϕ representing f .

If f has persistencies $y^1 \in \{0, 1\}^{S_1}$ and $y^2 \in \{0, 1\}^{S_2}$, then it also has a persistency $y^3 \in \{0, 1\}^{S_1 \cup S_2}$.

Every posiform ϕ has a unique maximal subset $S = S(\phi)$ for which it has an autarky $y \in \{0, 1\}^S$.

For a **quadratic** function f it is “**easy**” to **find** the unique maximal persistent set $S(f)$ provable by quadratic posiforms. (B. and Hammer, 1990)

Outline

- 1 Pseudo-Boolean Optimization
 - Pseudo-Boolean Functions
 - Representations and Bounds
 - Persistencies and Autarkies
 - Graph Cut Models
 - Implication Networks
- 2 Results
 - Computational Results

Network Model for Submodular QPBO (Hammer, 1965)

- A QPBF is submodular IFF all quadratic coefficients are nonpositive. *(Doit Yourself, anytime)*
- To a submodular QPBF f associate a network G_f as follows

=

- There is a one-to-one correspondence between values of f and $s - t$ cut values of G_f . *(Hammer, 1965)*
- **Graph cuts (325000)** in computer vision: (Greig, Porteous, and Seheult, 1989), (Boykov, Veksler and Zabih, 1998)

Network Model for Submodular QPBO (Hammer, 1965)

s

3

1

2

t

- A QPBF is submodular IFF all quadratic coefficients are nonpositive. (*Doit Yourself, anytime*)
- To a submodular QPBF f associate a network G_f as follows

$$f = 4 - x_1 + 7x_2 + x_3 - 3x_1x_2 - x_1x_3 - 2x_2x_3$$

$$=$$

- There is a one-to-one correspondence between values of f and $s - t$ cut values of G_f . (*Hammer, 1965*)
- **Graph cuts** (**325000**) in computer vision: (Greig, Porteous, and Seheult, 1989), (Boykov, Veksler and Zabih, 1998)

Network Model for Submodular QPBO (Hammer, 1965)

s

- A QPBF is submodular IFF all quadratic coefficients are nonpositive. (*Doit Yourself, anytime*)
- To a submodular QPBF f associate a network G_f as follows

$$\begin{aligned} f &= 4 - x_1 + 7x_2 + x_3 - 3x_1x_2 - x_1x_3 - 2x_2x_3 \\ &= 4\bar{x}_1 + 5x_2 + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3 \\ &= \end{aligned}$$

2

1

t

- There is a one-to-one correspondence between values of f and $s - t$ cut values of G_f . (*Hammer, 1965*)
- **Graph cuts (325000)** in computer vision: (Greig, Porteous, and Seheult, 1989), (Boykov, Veksler and Zabih, 1998)

Network Model for Submodular QPBO (Hammer, 1965)

s

- A QPBF is submodular IFF all quadratic coefficients are nonpositive. (*Doit Yourself, anytime*)
- To a submodular QPBF f associate a network G_f as follows

3

1

2

$$f = 4 - x_1 + 7x_2 + x_3 - 3x_1x_2 - x_1x_3 - 2x_2x_3$$


$$= 4\bar{x}_1 + 5x_2 + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3$$

t

$$= 4s\bar{x}_1 + 5x_2\bar{t} + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3$$

- There is a one-to-one correspondence between values of f and $s - t$ cut values of G_f . (*Hammer, 1965*)
- **Graph cuts (325000)** in computer vision: (Greig, Porteous, and Seheult, 1989), (Boykov, Veksler and Zabih, 1998)

Network Model for Submodular QPBO (Hammer, 1965)




- A QPBF is submodular IFF all quadratic coefficients are nonpositive. *(Doit Yourself, anytime)*
- To a submodular QPBF f associate a network G_f as follows

$$\begin{aligned}
 f &= 4 - x_1 + 7x_2 + x_3 - 3x_1x_2 - x_1x_3 - 2x_2x_3 \\
 &= 4\bar{x}_1 + 5x_2 + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3 \\
 &= 4s\bar{x}_1 + 5x_2\bar{t} + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3
 \end{aligned}$$

- There is a one-to-one correspondence between values of f and $s - t$ cut values of G_f . *(Hammer, 1965)*
- **Graph cuts (325000)** in computer vision: (Greig, Porteous, and Seheult, 1989), (Boykov, Veksler and Zabih, 1998)

Network Model for Submodular QPBO (Hammer, 1965)




- A QPBF is submodular IFF all quadratic coefficients are nonpositive. *(Doit Yourself, anytime)*
- To a submodular QPBF f associate a network G_f as follows

$$\begin{aligned}
 f &= 4 - x_1 + 7x_2 + x_3 - 3x_1x_2 - x_1x_3 - 2x_2x_3 \\
 &= 4\bar{x}_1 + 5x_2 + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3 \\
 &= 4s\bar{x}_1 + 5x_2\bar{t} + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3
 \end{aligned}$$

- There is a one-to-one correspondence between values of f and $s - t$ cut values of G_f . *(Hammer, 1965)*
- **Graph cuts (325000)** in computer vision: (Greig, Porteous, and Seheult, 1989), (Boykov, Veksler and Zabih, 1998)

Network Model for Submodular QPBO (Hammer, 1965)




- A QPBF is submodular IFF all quadratic coefficients are nonpositive. *(Doit Yourself, anytime)*
- To a submodular QPBF f associate a network G_f as follows

$$\begin{aligned}
 f &= 4 - x_1 + 7x_2 + x_3 - 3x_1x_2 - x_1x_3 - 2x_2x_3 \\
 &= 4\bar{x}_1 + 5x_2 + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3 \\
 &= 4s\bar{x}_1 + 5x_2\bar{t} + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3
 \end{aligned}$$

- There is a one-to-one correspondence between values of f and $s - t$ cut values of G_f . *(Hammer, 1965)*
- **Graph cuts (325000)** in computer vision: (Greig, Porteous, and Seheult, 1989), (Boykov, Veksler and Zabih, 1998)

Network Model for Submodular QPBO (Hammer, 1965)




- A QPBF is submodular IFF all quadratic coefficients are nonpositive. *(Doit Yourself, anytime)*
- To a submodular QPBF f associate a network G_f as follows

$$\begin{aligned}
 f &= 4 - x_1 + 7x_2 + x_3 - 3x_1x_2 - x_1x_3 - 2x_2x_3 \\
 &= 4\bar{x}_1 + 5x_2 + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3 \\
 &= 4s\bar{x}_1 + 5x_2\bar{t} + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3
 \end{aligned}$$

- There is a one-to-one correspondence between values of f and $s - t$ cut values of G_f . *(Hammer, 1965)*
- **Graph cuts (325000)** in computer vision: (Greig, Porteous, and Seheult, 1989), (Boykov, Veksler and Zabih, 1998)

Network Model for Submodular QPBO (Hammer, 1965)




- A QPBF is submodular IFF all quadratic coefficients are nonpositive. *(Doit Yourself, anytime)*
- To a submodular QPBF f associate a network G_f as follows

$$\begin{aligned}
 f &= 4 - x_1 + 7x_2 + x_3 - 3x_1x_2 - x_1x_3 - 2x_2x_3 \\
 &= 4\bar{x}_1 + 5x_2 + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3 \\
 &= 4s\bar{x}_1 + 5x_2\bar{t} + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3
 \end{aligned}$$

- There is a one-to-one correspondence between values of f and $s - t$ cut values of G_f . *(Hammer, 1965)*
- Graph cuts (325000) in computer vision: (Greig, Porteous, and Seheult, 1989), (Boykov, Veksler and Zabih, 1998)

Network Model for Submodular QPBO (Hammer, 1965)




- A QPBF is submodular IFF all quadratic coefficients are nonpositive. *(Doit Yourself, anytime)*
- To a submodular QPBF f associate a network G_f as follows

$$\begin{aligned}
 f &= 4 - x_1 + 7x_2 + x_3 - 3x_1x_2 - x_1x_3 - 2x_2x_3 \\
 &= 4\bar{x}_1 + 5x_2 + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3 \\
 &= 4s\bar{x}_1 + 5x_2\bar{t} + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3
 \end{aligned}$$

- There is a one-to-one correspondence between values of f and $s - t$ cut values of G_f . *(Hammer, 1965)*
- Graph cuts (325000) in computer vision: (Greig, Porteous, and Seheult, 1989), (Boykov, Veksler and Zabih, 1998)

Network Model for Submodular QPBO (Hammer, 1965)



- A QPBF is submodular IFF all quadratic coefficients are nonpositive. *(Doit Yourself, anytime)*
- To a submodular QPBF f associate a network G_f as follows

$$\begin{aligned}
 f &= 4 - x_1 + 7x_2 + x_3 - 3x_1x_2 - x_1x_3 - 2x_2x_3 \\
 &= 4\bar{x}_1 + 5x_2 + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3 \\
 &= 4s\bar{x}_1 + 5x_2\bar{t} + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3
 \end{aligned}$$

- There is a one-to-one correspondence between values of f and $s - t$ cut values of G_f . *(Hammer, 1965)*

$$f(\mathbf{0}, \mathbf{1}, \mathbf{0}) = C(\{\mathbf{s}, \mathbf{2}\}, \{\mathbf{1}, \mathbf{3}, \mathbf{t}\}) = 11$$

- Graph cuts (**325000**) in computer vision: (Greig, Porteous, and Seheult, 1989), (Boykov, Veksler and Zabih, 1998)

Network Model for Submodular QPBO (Hammer, 1965)

- A QPBF is submodular IFF all quadratic coefficients are nonpositive. *(Doit Yourself, anytime)*
- To a submodular QPBF f associate a network G_f as follows

$$\begin{aligned}
 f &= 4 - x_1 + 7x_2 + x_3 - 3x_1x_2 - x_1x_3 - 2x_2x_3 \\
 &= 4\bar{x}_1 + 5x_2 + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3 \\
 &= 4s\bar{x}_1 + 5x_2\bar{t} + 3x_1\bar{x}_2 + \bar{x}_1x_3 + 2x_2\bar{x}_3
 \end{aligned}$$

- There is a one-to-one correspondence between values of f and $s - t$ cut values of G_f . *(Hammer, 1965)*

$$f(\mathbf{0}, \mathbf{1}, \mathbf{0}) = C(\{\mathbf{s}, \mathbf{2}\}, \{\mathbf{1}, \mathbf{3}, \mathbf{t}\}) = 11$$

- **Graph cuts (325000)** in computer vision: (Greig, Porteous, and Seheult, 1989), (Boykov, Veksler and Zabih, 1998)

Outline

- 1 Pseudo-Boolean Optimization
 - Pseudo-Boolean Functions
 - Representations and Bounds
 - Persistencies and Autarkies
 - Graph Cut Models
 - Implication Networks
- 2 Results
 - Computational Results

QPBF \longrightarrow Posiform

$$f = 10 - 2x_1 - 6x_2 + 2x_1x_2 - 2x_1x_3 + 4x_2x_3$$

QPBF \longrightarrow Posiform

$$\begin{aligned} f &= 10 - 2x_1 - 6x_2 + 2x_1x_2 - 2x_1x_3 + 4x_2x_3 \\ &= 10 - 2x_1 - 6x_2 + 2x_1x_2 - 2x_1(1 - \bar{x}_3) + 4x_2x_3 \end{aligned}$$

QPBF \longrightarrow Posiform

$$\begin{aligned} f &= 10 - 2x_1 - 6x_2 + 2x_1x_2 - 2x_1x_3 + 4x_2x_3 \\ &= 10 - 2x_1 - 6x_2 + 2x_1x_2 - 2x_1(1 - \bar{x}_3) + 4x_2x_3 \\ &= 10 - 4x_1 - 6x_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \end{aligned}$$

QPBF \longrightarrow Posiform

$$\begin{aligned} f &= 10 - 2x_1 - 6x_2 + 2x_1x_2 - 2x_1x_3 + 4x_2x_3 \\ &= 10 - 2x_1 - 6x_2 + 2x_1x_2 - 2x_1(1 - \bar{x}_3) + 4x_2x_3 \\ &= 10 - 4x_1 - 6x_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \\ &= 10 - 4(1 - \bar{x}_1) - 6(1 - \bar{x}_2) + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \end{aligned}$$

QPBF \longrightarrow Posiform


$$\begin{aligned} f &= 10 - 2x_1 - 6x_2 + 2x_1x_2 - 2x_1x_3 + 4x_2x_3 \\ &= 10 - 2x_1 - 6x_2 + 2x_1x_2 - 2x_1(1 - \bar{x}_3) + 4x_2x_3 \\ &= 10 - 4x_1 - 6x_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \\ &= 10 - 4(1 - \bar{x}_1) - 6(1 - \bar{x}_2) + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \\ &= 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 = \phi \end{aligned}$$

QPBF \longrightarrow Posiform

$$\begin{aligned}f &= 10 - 2\mathbf{x}_1 - 6\mathbf{x}_2 + 2\mathbf{x}_1\mathbf{x}_2 - 2\mathbf{x}_1\mathbf{x}_3 + 4\mathbf{x}_2\mathbf{x}_3 \\&= 10 - 2x_1 - 6x_2 + 2x_1x_2 - 2x_1(1 - \bar{x}_3) + 4x_2x_3 \\&= 10 - 4x_1 - 6x_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \\&= 10 - 4(1 - \bar{x}_1) - 6(1 - \bar{x}_2) + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \\&= 4\bar{x}_1 + 6\bar{x}_2 + 2\mathbf{x}_1\mathbf{x}_2 + 2\mathbf{x}_1\bar{x}_3 + 4\mathbf{x}_2\mathbf{x}_3 = \phi\end{aligned}$$


$$\mathbf{C}(\phi) = \mathbf{0} \quad \text{and} \quad \mathbf{S}(\phi) = \emptyset$$

Implication Networks





$$f = 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3$$

Implication Networks




$$f = 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3$$

Implication Networks




$$f = 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3$$

Implication Networks




$$f = 4\bar{x}_1 + 6\bar{x}_2 + \mathbf{2x_1x_2} + 2x_1\bar{x}_3 + 4x_2x_3$$

Implication Networks




$$f = 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3$$

Implication Networks




$$f = 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3$$

Implication Networks




$$f = 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3$$

Flows and Posiform Transformations




$$f = 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3$$

Flows and Posiform Transformations




$$\begin{aligned}
 f &= 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \\
 &= 3\bar{x}_1 + 5\bar{x}_2 + 2x_1x_2 + x_1\bar{x}_3 + 3x_2x_3 \\
 &\quad + (\bar{x}_1 + x_1\bar{x}_3 + x_3x_2 + \bar{x}_2)
 \end{aligned}$$

Flows and Posiform Transformations




$$\begin{aligned}
 f &= 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \\
 &= 3\bar{x}_1 + 5\bar{x}_2 + 2x_1x_2 + x_1\bar{x}_3 + 3x_2x_3 \\
 &\quad + (\bar{x}_1 + x_1\bar{x}_3 + x_3x_2 + \bar{x}_2) \\
 &= 3\bar{x}_1 + 5\bar{x}_2 + 2x_1x_2 + x_1\bar{x}_3 + 3x_2x_3 \\
 &\quad + (1 + \bar{x}_1x_3 + \bar{x}_3\bar{x}_2)
 \end{aligned}$$

Flows and Posiform Transformations




$$\begin{aligned}
 f &= 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \\
 &= 3\bar{x}_1 + 5\bar{x}_2 + 2x_1x_2 + x_1\bar{x}_3 + 3x_2x_3 \\
 &\quad + (\bar{x}_1 + x_1\bar{x}_3 + x_3x_2 + \bar{x}_2) \\
 &= 3\bar{x}_1 + 5\bar{x}_2 + 2x_1x_2 + x_1\bar{x}_3 + 3x_2x_3 \\
 &\quad + (1 + \bar{x}_1x_3 + \bar{x}_3\bar{x}_2) \\
 &= 1 + 3\bar{x}_1 + 5\bar{x}_2 + 2x_1x_2 + x_1\bar{x}_3 + 3x_2x_3 + \bar{x}_1x_3 + \bar{x}_3\bar{x}_2
 \end{aligned}$$

Flows and Posiform Transformations




$$\begin{aligned}
 f &= 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \\
 &= 1 + 3\bar{x}_1 + 5\bar{x}_2 + 2x_1x_2 + x_1\bar{x}_3 + 3x_2x_3 + \bar{x}_1x_3 + \bar{x}_3\bar{x}_2
 \end{aligned}$$

Flows and Posiform Transformations




$$\begin{aligned}
 f &= 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \\
 &= 1 + 3\bar{x}_1 + 5\bar{x}_2 + 2x_1x_2 + x_1\bar{x}_3 + 3x_2x_3 + \bar{x}_1x_3 + \bar{x}_3\bar{x}_2 \\
 &= 4 + 2\bar{x}_2 + 2\bar{x}_1\bar{x}_2 + 2\bar{x}_1x_3 + 2x_2x_3 + 2\bar{x}_2\bar{x}_3
 \end{aligned}$$

Persistencies and Decompositions




$$\begin{aligned}
 f &= 10 - 2x_1 - 6x_2 + 2x_1x_2 - 2x_1x_3 + 4x_2x_3 \\
 &= 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \\
 &= 4 + 2\bar{x}_2 + 2\bar{x}_1\bar{x}_2 + 2\bar{x}_1x_3 + 2x_2x_3 + 2\bar{x}_2\bar{x}_3
 \end{aligned}$$

Persistencies and Decompositions




$$\begin{aligned}
 f &= 10 - 2x_1 - 6x_2 + 2x_1x_2 - 2x_1x_3 + 4x_2x_3 \\
 &= 4\bar{x}_1 + 6\bar{x}_2 + 2x_1x_2 + 2x_1\bar{x}_3 + 4x_2x_3 \\
 &= 4 + 2\bar{x}_2 + 2\bar{x}_1\bar{x}_2 + 2\bar{x}_1x_3 + 2x_2x_3 + 2\bar{x}_2\bar{x}_3
 \end{aligned}$$

Persistencies and Decompositions




- Strong persistency: $x_2 = 1, x_3 = 0$
- Weak persistency: $x_1 = 1$ (or $x_1 = 0$)
- This method computes the **unique maximal set** of autark variables justifiable by any quadratic posiform of the given quadratic PBF (B. and Hammer, 1990).

Persistencies and Decompositions



- Strong persistency: $x_2 = 1, x_3 = 0$
- Weak persistency: $x_1 = 1$ (or $x_1 = 0$)
- This method computes the **unique maximal set** of autark variables justifiable by any quadratic posiform of the given quadratic PBF (B. and Hammer, 1990).

Persistencies and Decompositions



- Strong persistency: $x_2 = 1, x_3 = 0$
- Weak persistency: $x_1 = 1$ (or $x_1 = 0$)
- This method computes the **unique maximal set** of autark variables justifiable by any quadratic posiform of the given quadratic PBF (B. and Hammer, 1990).

Outline

- 1 Pseudo-Boolean Optimization
 - Pseudo-Boolean Functions
 - Representations and Bounds
 - Persistencies and Autarkies
 - Graph Cut Models
 - Implication Networks
- 2 Results
 - Computational Results

Via Minimization in VLSI Design

		Percentage of Variables Fixed by					
Problem	n	Persistency		Probing		ALL TOOLS	Time (sec)
		(strong)	(weak)	(forc)	(equal)		
via.c1y	829	93.6%	6.4%	0%	0%	100%	0.03
via.c2y	981	94.7%	5.3%	0%	0%	100%	0.06
via.c3y	1328	94.6%	5.4%	0%	0%	100%	0.09
via.c4y	1367	96.4%	3.6%	0%	0%	100%	0.09
via.c5y	1203	93.1%	6.9%	0%	0%	100%	0.08
via.c1n	828	57.4%	9.6%	32.4%	0.6%	100%	0.49
via.c2n	980	12.4%	4.4%	83.1%	0.1%	100%	7.14
via.c3n	1327	6.8%	5.7%	87.3%	0.2%	100%	18.17
via.c4n	1366	11.1%	1.3%	87.6%	0%	100%	23.08
via.c5n	1202	3.4%	1.4%	95.0%	0.2%	100%	17.13

¹S. Homer and M. Peinado. Design and performance of parallel and distributed approximation algorithms for maxcut. Journal of Parallel and Distributed Computing 46 (1997) 48-61.

Vertex Cover in Planar Graphs

	Averages for 100 graphs in each of the 4 groups			
	Variables Fixed (%)		Time (sec)	
n	A. D. N. ²	QUBO ³	A. D. N. ²	QUBO ³
1000	68.4	100	4.06	0.05
2000	67.4	100	12.24	0.16
3000	65.5	100	30.90	0.27
4000	62.7	100	60.45	0.53

²Alber, Dorn, Niedermeier. Experimental evaluation of a tree decomposition based algorithm for vertex cover on planar graphs. Discrete Applied Mathematics 145 (2005) 219-231; 750 GHz, Linux PC, 720 MB

³Pentium 4, 2.8 GHz, Windows XP, 512 MB

Jumbo Vertex Cover in Planar Graphs

Vertices	Computing Times (min) ⁴		
	Planar Density		
	10%	50%	90%
50,000	0.7	2.3	0.9
100,000	2.9	10.2	3.9
250,000	19.5	69.8	26.3
500,000	79.3	277.3	106.9

QUBO fixed all variables for all problems!

⁴Averages over 3 experiments on a Xeon 3.06 GHz, XP, 3.5 GB RAM.

One Dimensional Ising Models

σ	Number of Spins	Average Computing Time (s)		
		Branch, Cut & Price ⁵	BiqMaq ⁵	QUBO⁶
2.5	100	699	68	1
	150	92 079	388	3
	200	N/A	993	9
	250	N/A	6 567	14
	300	N/A	34 572	21
3.0	100	256	59	1
	150	13 491	293	2
	200	61 271	1 034	3
	250	55 795	3 594	4
	300	55 528	8 496	5

⁵F. Rendl, G. Rinaldi, A. Wiegele. (2007). Solving max-cut to optimality by intersecting semidefinite and polyhedral relaxations.

⁶ALL problems were solved by QUBO.

Larger One Dimensional Ising Models

σ	n	Average of 3 Problems	
		Variables not fixed	QUBO Time (s) ⁷
2.5	500	5	13
	750	22	30
	1000	24	53
	1250	20	81
	1500	32	124
3.0	500	0	4
	750	0	12
	1000	0	23
	1250	0	37
	1500	0	59

⁷Pentium M, 1.6 GHz 760 MB RAM

Open Ends

- How can we extend the above results for general PBF-s?
- How can we find autarkies for a given posiform?
- Quadratization via graph stability (Ebenegger, Hammer, de Werra, 1984): Can we map persistencies?
- Quadratization in higher dimension: given f , find quadratic g such that

$$f(x) = \min_{y \in \{0,1\}^m} g(x, y) \text{ for all } x \in \{0,1\}^n$$

Does it exist? How many new variables do we need?
(Hammer, 1975)

Which quadratization procedure works the best?
(Hammer, 1975)

Open Ends

- How can we extend the above results for general PBF-s?
- How can we find autarkies for a given posiform?
- Quadratization via graph stability (Ebenegger, Hammer, de Werra, 1984): Can we map persistencies?
- Quadratization in higher dimension: given f , find quadratic g such that

$$f(x) = \min_{y \in \{0,1\}^m} g(x, y) \text{ for all } x \in \{0,1\}^n$$

• How can we extend the above results for general PBF-s?

• How can we find autarkies for a given posiform?

• Quadratization via graph stability (Ebenegger, Hammer, de Werra, 1984): Can we map persistencies?

• Quadratization in higher dimension: given f , find quadratic g such that

Open Ends

- How can we extend the above results for general PBF-s?
- How can we find autarkies for a given posiform?
- Quadratzation via graph stability (Ebenegger, Hammer, de Werra, 1984): Can we map persistencies?
- Quadratzation in higher dimension: given f , find quadratic g such that

$$f(x) = \min_{y \in \{0,1\}^m} g(x, y) \text{ for all } x \in \{0,1\}^n$$

- Does it exists? How many new variables do we need? (Rosenberg, 1975), ...
- Which quadratic form provides us with the min

Open Ends

- How can we extend the above results for general PBF-s?
- How can we find autarkies for a given posiform?
- Quadratization via graph stability (Ebenegger, Hammer, de Werra, 1984): Can we map persistencies?
- Quadratization in higher dimension: given f , find quadratic g such that

$$f(x) = \min_{y \in \{0,1\}^m} g(x, y) \text{ for all } x \in \{0,1\}^n$$

- Does it exists? How many new variables do we need? (Rosenberg, 1975), ...
- Which quadratization provides us with the most persistencies?

Open Ends

- How can we extend the above results for general PBF-s?
- How can we find autarkies for a given posiform?
- Quadratization via graph stability (Ebenegger, Hammer, de Werra, 1984): Can we map persistencies?
- Quadratization in higher dimension: given f , find quadratic g such that

$$f(x) = \min_{y \in \{0,1\}^m} g(x, y) \text{ for all } x \in \{0,1\}^n$$

- Does it exists? How many new variables do we need? (Rosenberg, 1975), ...
- Which quadratization provides us with the most persistencies?

Open Ends

- How can we extend the above results for general PBF-s?
- How can we find autarkies for a given posiform?
- Quadratization via graph stability (Ebenegger, Hammer, de Werra, 1984): Can we map persistencies?
- Quadratization in higher dimension: given f , find quadratic g such that

$$f(x) = \min_{y \in \{0,1\}^m} g(x, y) \text{ for all } x \in \{0,1\}^n$$

- Does it exists? How many new variables do we need? (Rosenberg, 1975), ...
- Which quadratization provides us with the most persistencies?

References

- Hammer, P.L. Some network flow problems solved with pseudo-Boolean programming. *Operations Research* **13** (1965) 388-399.
- Hammer, P.L. and S. Rudeanu. *Boolean Methods in Operations Research and Related Areas*. (Springer-Verlag, Berlin, Heidelberg, New York, 1968.)
- Hammer, P.L., P. Hansen and B. Simeone. Roof duality, complementation and persistency in quadratic 0 – 1 optimization. *Mathematical Programming* **28** (1984), pp. 121-155.
- Boros E. and P.L. Hammer. A max-flow approach to improved roof-duality in quadratic 0 – 1 minimization. RUTCOR Research Report RRR 15-1989, RUTCOR, March 1989.
- Boros, E., Y. Crama, and P.L. Hammer. Upper bounds for quadratic 0 – 1 maximization. *Operations Research Letters*, **9** (1990), 73-79,
- Billionnet, A. and A. Sutter. Persistency in quadratic 0 – 1 optimization. *Math. Programming* **54** (1992), no. 1, Ser. A, pp. 115–119.

References Cont'd

- Boros, E., Y. Crama and P.L. Hammer. Chvátal cuts and odd cycle inequalities in quadratic 0 – 1 optimization. *SIAM Journal on Discrete Mathematics*, **5** (1992), 163-177.
- Boros, E. and P.L. Hammer, Pseudo-Boolean Optimization, *Discrete Applied Mathematics*, **123** (2002) 155–225.
- Boros, E., P.L. Hammer, G. Tavares, and R. Sun. A max-flow approach to improved lower bounds for quadratic 0 – 1 minimization. *Discrete Optimization*, **5/2** (2007) 501-529.
- E. Boros, Y-H. Chen, A. Fix, A.G. Gruber, J. Schwartz, R. Zabih: A graph cut algorithm for higher order Markov random fields. IEEE Conference Proceedings for ICCV 2011, Barcelona, Spain (2011), pp. 1020-1027.
- A. Fix, Y-H. (Joyce) Chen, E. Boros, and R. Zabih: A graph cut algorithm based on approximating the signed bi-form graph, European Conference on Computer Vision, 2012, Part I, LNCS 7572, pp. 385-398.