

- bury, A. Schilling, D. L. Venton, *J. Med. Chem.* **1997**, *40*, 4006–4012.
 c) Y. F. Hsieh, N. Gordon, F. Regnier, N. Afeyan, S. A. Martin, G. J. Vella, *Mol. Div.* **1996**, *2*, 189–196.
 [6] See, for example, a) R. W. Nelson, J. R. Krone, A. L. Bieber, P. Williams, *Anal. Chem.* **1995**, *67*, 1153–1158; b) D. C. Schriemer, L. Li, *Anal. Chem.* **1996**, *68*, 3382–3387; c) R. Wieboldt, J. Zweigenbaum, J. Henion, *Anal. Chem.* **1997**, *69*, 1683–1691; d) R. B. van Breemen, C.-R. Huang, D. Nikolic, C. P. Woodbury, Y.-Z. Zhao, D. L. Venton, *Anal. Chem.* **1997**, *69*, 2159–2164.
 [7] K.-I. Kasai, Y. J. Oda, *Chromatogr.* **1986**, *376*, 33–47.
 [8] See, for example, a) H. Ibrahim, A. F. Aubry, *Anal. Biochem.* **1995**, *229*, 272–277; b) J. Renard, C. Vidal Madjar, C. Lapresle, *J. Colloid Interface Sci.* **1995**, *174*, 61–67; c) M. Bjorklund, M. T. W. Hearn, *J. Chromatogr. A* **1996**, *145*–162; d) S. Lin, L. R. Drake, G. D. Rayson, *Anal. Chem.* **1996**, *68*, 4087–4093, and references therein.
 [9] D. R. Bundle, E. Eichler, M. A. J. Gidney, M. Meldal, A. Ragauskas, B. W. Sigurskjold, B. Sinnott, D. C. Watson, M. Yaguchi, N. M. Young, *Biochemistry* **1994**, *33*, 5172–5182.
 [10] E. Altman, D. R. Bundle, *Meth. Enzymol.* **1994**, *242*, 243–253.
 [11] U. Bahr, A. Pfenninger, M. Karas, B. Stahl, *Anal. Chem.* **1997**, *69*, 4530–4535.
 [12] D. C. Schriemer, L. Li, *Anal. Chem.* **1996**, *68*, 3382–3387.

A Convenient and General Method for Pd-Catalyzed Suzuki Cross-Couplings of Aryl Chlorides and Arylboronic Acids**

Adam F. Littke and Gregory C. Fu*


In memory of George Büchi

The Suzuki reaction has emerged as an extremely powerful method for the cross-coupling of aryl bromides, iodides, and triflates with arylboronic acids.^[1] As compared with the Stille cross-coupling,^[2] the Suzuki reaction possesses the practical advantages that the boron-containing by-products, unlike the tin-containing by-products of the Stille process, are nontoxic and are easily separated from the desired product.^[3] One current limitation in the scope of the Suzuki reaction is its inefficiency when aryl chlorides are employed as substrates. Although there have been several accounts of Suzuki reactions of electron-poor aryl chlorides,^[4–6] to the best of our knowledge there has been only one report of a coupling of an electron-neutral or an electron-rich aryl chloride (41 % yield).^[5] In view of the increased availability and decreased

expense of aryl chlorides relative to aryl bromides and iodides,^[4a] this methodological gap becomes all the more significant. Herein we describe a solution to this problem: specifically, the development of conditions under which Suzuki cross-couplings of an array of electronically and sterically different aryl chlorides and arylboronic acids proceed in excellent yield with commercially available reagents (see Table 3).

In early experiments we established that 4-chlorotoluene and phenylboronic acid are efficiently cross-coupled in the presence of 1.5 % [Pd₂(dba)₃] (dba = dibenzylideneacetone), 3.6 % *Pr*Bu₃, and two equivalents of Cs₂CO₃ (dioxane, 80 °C; 86 % yield by GC after 5.0 h; Table 1, entry 9). We observe little or no coupling in the absence of phosphane (entry 1) or in the presence of triarylphosphanes (entries 2–5) and

Table 1. Effect of phosphane on the palladium-catalyzed Suzuki cross-coupling of aryl chlorides.



Entry	Phosphane	Yield [%] (GC)
1	–	0
2	PPh ₃	0
3	BINAP ^[a]	0
4	dppf ^[b]	0
5	P(<i>o</i> -tol) ₃	10
6	Ph ₂ P(CH ₂) ₃ PPh ₂	0
7	Cy ₂ P(CH ₂) ₂ PCy ₂	0
8	PCy ₃	75
9	<i>Pr</i> Bu ₃	86

[a] BINAP = 2,2'-bis(diphenylphosphanyl)-1,1'-binaphthyl. [b] dppf = 1,1'-bis(diphenylphosphanyl)ferrocene.

chelating phosphanes (entries 6 and 7). Of the ligands that we have examined to date, only PCy₃ approaches *Pr*Bu₃ in effectiveness (entries 8 and 9). The use of a phosphane: Pd ratio between 1 and 1.5 appears to be optimal with *Pr*Bu₃ as the ligand, an observation that suggests that the active palladium catalyst bears a single *Pr*Bu₃ group. Thus, we believe that the steric bulk and the electron-richness of *Pr*Bu₃ are critical for this unprecedented reactivity.^[4, 7]

Among the bases listed in Table 2, Cs₂CO₃ has proved to be the base of choice (entry 6).^[8, 9] In the absence of base (entry 1) or in the presence of other bases commonly used in Suzuki reactions (entries 2–5), cross-coupling proceeds with diminished efficiency. In contrast to most Suzuki cross-couplings,^[1] only one equivalent of base is required for our process (95 % GC yield after five hours with 1.2 equiv Cs₂CO₃). We have found [Pd₂(dba)₃] and dioxane to be best among the palladium sources (for example, Pd(OAc)₂) and solvents (for example, THF) that we have surveyed thus far.

A wide array of electronically and structurally diverse aryl chlorides and arylboronic acids can be cross-coupled very efficiently under these conditions (1.5 % [Pd₂(dba)₃], 3.6 % *Pr*Bu₃, 1.2 equiv Cs₂CO₃).^[10] Thus, with respect to the aryl chloride, both electron-neutral 4-chlorotoluene and electron-

[*] Prof. Dr. G. C. Fu, A. F. Littke
 Department of Chemistry
 Massachusetts Institute of Technology
 Cambridge, Massachusetts 02139 (USA)
 Fax: (+1) 617-258-7500
 E-mail: gcf@mit.edu

[**] Support has been provided by the Alfred P. Sloan Foundation, the American Cancer Society, Bristol–Myers Squibb, the Camille and Henry Dreyfus Foundation, Eli Lilly, Firmenich, Glaxo Wellcome, the National Science Foundation (Young Investigator Award, with funding from Merck, Pharmacia & Upjohn, DuPont, Bayer, and Novartis), the Natural Sciences and Engineering Research Council of Canada (predoctoral fellowship to A.F.L.), Pfizer, Procter & Gamble, and the Research Corporation.

Supporting information for this article is available on the WWW under <http://www.wiley-vch.de/home/angewandte/> or from the author.

Table 2. Effect of base on the palladium-catalyzed Suzuki cross-coupling of aryl chlorides.

Entry	Base	Yield [%] ^[a]	Entry	Base	Yield [%] ^[a]
1	–	3	4	NEt ₃	50
2	Na ₂ CO ₃	23	5	K ₃ PO ₄	83
3	K ₂ CO ₃	29	6	Cs ₂ CO ₃	86

[a] Determined by gas chromatography.

Table 3. Scope of the [Pd₂(dba)₃]/PtBu₃/Cs₂CO₃-catalyzed Suzuki cross-coupling of aryl chlorides.

Entry	X	Y	Yield [%]
1	4-Me	H	87
2	4-MeCO	H	91
3	4-MeO	H	89
4	4-NH ₂	H	92
5	4-Me	4-CF ₃	86
6	4-Me	4-OMe	82
7	2-Me	H	90
8	2-Me	2-Me	87

poor 4-chloroacetophenone react with PhB(OH)₂ to provide the biaryl product in excellent yield (Table 3, entries 1 and 2). More remarkable is our observation that even extremely electron-rich aryl chlorides, including 4-chloroanisole and 4-chloroaniline, couple cleanly with PhB(OH)₂ in the presence of this catalyst system (entries 3 and 4).

The cross-coupling also proceeds smoothly regardless of the electronic character of the boronic acid component. More electron-rich boronic acids have been reported to be less reactive partners in other Suzuki reactions.^[1] With our catalyst system, trifluoromethyl- and methoxy-substituted benzeneboronic acids couple with 4-chlorotoluene to produce the desired biaryls in excellent yields under comparable reaction conditions (Table 3, entry 5: 5 h at 80 °C; entry 6: 9 h at 80 °C).

Finally, this [Pd₂(dba)₃]/PtBu₃-catalyzed cross-coupling is tolerant of *ortho* substitution in both the aryl halide and the arylboronic acid (Table 3, entries 7 and 8). For example, sterically hindered 2,2'-dimethyl-1,1'-biphenyl is formed in 87% yield after seven hours at 80 °C.

In summary, we have developed a general method for the Suzuki cross-coupling of aryl chlorides with arylboronic acids. Our catalyst system, which employs readily available components (1.5% [Pd₂(dba)₃], 3.6% PtBu₃, 1.2 equiv Cs₂CO₃), couples effectively a broad spectrum of substrates, to provide the desired biaryls in excellent yields. Because of the accessibility of aryl chlorides and the relatively innocuous nature of the boron-containing by-products of Suzuki reactions, we anticipate that this method is likely to find widespread use in synthetic organic chemistry. Future work will be

directed at further optimization of this process and on the development of related coupling reactions.^[11]

Received: August 18, 1998 [Z12301 IE]

German version: *Angew. Chem.* **1998**, *110*, 3586–3587

Keywords: cross-coupling • homogeneous catalysis • palladium • Suzuki reaction

- [1] For a review, see N. Miyaura, A. Suzuki, *Chem. Rev.* **1995**, *95*, 2457–2483.
- [2] For a review, see V. Farina, V. Krishnamurthy, W. J. Scott, *Org. React.* **1997**, *50*, 1–652.
- [3] For discussions of the toxicity of triorganotin compounds, see a) S. J. De Mora, *Tributyltin: Case Study of an Environmental Contaminant*, Cambridge University Press, Cambridge, **1996**; b) I. J. Boyer, *Toxicology* **1989**, *55*, 253–298.
- [4] The low reactivity of aryl chlorides in cross-coupling reactions is generally ascribed to their reluctance to oxidatively add to Pd⁰. Aryl halides that bear an electron-withdrawing group oxidatively add to Pd⁰ more readily than do the corresponding unsubstituted aryl halides. For discussions, see a) V. V. Grushin, H. Alper, *Chem. Rev.* **1994**, *94*, 1047–1062; b) references [1, 2].
- [5] a) S. Gronowitz, A.-B. Hörnfeldt, V. Kristjansson, T. Musil, *Chem. Scr.* **1986**, *26*, 305–309; b) W. J. Thompson, J. H. Jones, P. A. Lyle, J. E. Thies, *J. Org. Chem.* **1988**, *53*, 2052–2055; c) M. B. Mitchell, P. J. Wallbank, *Tetrahedron Lett.* **1991**, *32*, 2273–2276; d) M. Uemura, H. Nishimura, K. Kamikawa, K. Nakayama, Y. Hayashi, *Tetrahedron Lett.* **1994**, *35*, 1909–1912; e) M. Beller, H. Fischer, W. A. Herrmann, K. Ofele, C. Brossmer, *Angew. Chem.* **1995**, *107*, 1992–1993; *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 1848–1849; f) S. Saito, M. Sakai, N. Miyaura, *Tetrahedron Lett.* **1996**, *37*, 2993–2996; g) W. Shen, *Tetrahedron Lett.* **1997**, *38*, 5575–5578; h) W. A. Herrmann, C.-P. Reisinger, M. Spiegler, *J. Organomet. Chem.* **1998**, *557*, 93–96; i) F. Firooznia, C. Gude, K. Chan, Y. Satoh, *Tetrahedron Lett.* **1998**, *39*, 3985–3988.
- [6] For nickel-catalyzed cross-couplings of aryl chlorides and arylboronic acids, see a) S. Saito, S. Oh-tani, N. Miyaura, *J. Org. Chem.* **1997**, *62*, 8024–8030; b) A. F. Indolese, *Tetrahedron Lett.* **1997**, *38*, 3513–3516; c) reference [5i].
- [7] a) For reports of enhanced reactivity in palladium-catalyzed cross-couplings when PtBu₃ is used as a ligand, see M. Nishiyama, T. Yamamoto, Y. Koie, *Tetrahedron Lett.* **1998**, *39*, 617–620; T. Yamamoto, M. Nishiyama, Y. Koie, *Tetrahedron Lett.* **1998**, *39*, 2367–2370; b) For other reports that deal with the use of electron-rich phosphanes in palladium-catalyzed cross-couplings, see M. Portnoy, D. Milstein, *Organometallics* **1993**, *12*, 1655–1664; M. Portnoy, D. Milstein, *Organometallics* **1993**, *12*, 1665–1673; M. Portnoy, Y. Ben-David, D. Milstein, *Organometallics* **1993**, *12*, 4734–4735; N. P. Reddy, M. Tanaka, *Tetrahedron Lett.* **1997**, *38*, 4807–4810; reference [4a].
- [8] a) For reports of the use of Cs₂CO₃ in Suzuki reactions, see H. E. Katz, *J. Org. Chem.* **1987**, *52*, 3932–3934; C. R. Johnson, M. P. Braun, *J. Am. Chem. Soc.* **1993**, *115*, 11014–11015; b) For reports of the use of Cs₂CO₃ in other palladium-catalyzed cross-couplings, see J. P. Wolfe, S. L. Buchwald, *Tetrahedron Lett.* **1997**, *38*, 6359–6362; T. Satoh, Y. Kawamura, M. Miura, M. Nomura, *Angew. Chem.* **1997**, *109*, 1820–1822; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 1740–1742.
- [9] Very recently we have found that CsF is also an extremely effective base.
- [10] General experimental conditions: A solution of aryl chloride (0.833 mmol; in dioxane (0.5–0.7 mL)) and a solution of PtBu₃ (0.030 mmol; in dioxane (0.3 mL)) were added in turn to a Schlenk tube charged with [Pd₂(dba)₃] (0.0125 mmol), boronic acid (0.875 mmol), Cs₂CO₃ (1.00 mmol), and a magnetic stir bar under an atmosphere of nitrogen. The Schlenk tube was stirred at 80 °C for five hours. The reaction mixture was then cooled to room temperature, diluted with Et₂O, filtered through a pad of Celite, concentrated, and purified by flash chromatography.
- [11] Note added in proof (November 2, 1998): Buchwald et al. have reported a versatile method for Suzuki cross-coupling of aryl chlorides: D. W. Old, J. P. Wolfe, S. L. Buchwald, *J. Am. Chem. Soc.* **1998**, *120*, 9722–9723.