

Research Letter

Syntheses of 5-Thio-D-Mannose from Petrochemicals and a Disaccharide Analog Containing It

Hideya Yuasa, Masatake Jouyabu, Nobuyuki Mitsuhashi, and Hironobu Hashimoto

Graduate School of Bioscience and Biotechnology, Tokyo Institute of Technology, 4259 Nagatsuta-cho, Midori-ku, Yokohama 226-8501, Japan

Correspondence should be addressed to Hideya Yuasa, hyuasa@bio.titech.ac.jp

Received 8 April 2008; Accepted 11 May 2008

Recommended by Robin Polt

Per-*O*-acetyl-5-thio-DL-mannose was synthesized from petrochemicals in six steps and 9% overall yield. It was then derivatized into glycosyl trichloroacetimidate and subjected to glycosidation reaction with a mannosyl acceptor to give a separatable mixture of disaccharides with 5-thio-D- and L-mannosides. This is the first synthesis of an enantiomerically pure 5-thiosugar derivative from racemic chemicals. The D-glycoside was derivatized into methyl (5-thio- α -D-mannopyranosyl)-2-*O*- α -D-mannopyranoside 6-phosphate as a potential inhibitor of a golgi α -1,2-mannosidase.

Copyright © 2008 Hideya Yuasa et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

5-thiosugars are aldopyranose analogs with a sulfur atom in the pyranose ring [1–3] and it has been demonstrated that oligosaccharide analogs with a 5-thiosugar at the nonreducing end are exoglycosidase-resistant [4–6]. The resistance to hydrolases is a property desired for potential drugs [7], allowing for a practical duration of potency, hence a number of 5-thiosugars have been synthesized [2] and their behaviors against glycosidases studied [8–10]. Although most of 5-thiosugars have been synthesized from natural carbohydrates in not less than 10 steps, a few trials were made for the construction of the thiopyran structure through hetero Diels-Alder reaction from petrochemicals [11–13]. With this strategy, however, 5-thiosugars are obtained necessarily in racemic form and no analogs for mammalian monosaccharides have been synthesized. One of the racemic 5-thiosugars obtained in this method is 5-thiomannuronate, a potential intermediate for the synthesis of 5-thio-D-mannose. 5-thio-D-mannose is the only 5-thiosugar that has been isolated from nature [14], awaiting studies for its biological significance. We thus report here on the synthesis of racemic 5-thiomannose (5SMan) from the product of a hetero Diels-Alder reaction reported by Prabhakaran et al. [12] and then on the glycosidation reaction of the 1-*O*-trichloroacetimidate

derivative of 5SMan, which gave a diastereomeric mixture of 5SMan-containing disaccharides, and they were easily separated by column chromatography. As a result, we obtained an enantiomerically pure 5-thiosugar derivative from petrochemicals for the first time. To make the synthesis more significant and advantageous, we derivatized the disaccharide analog into the 6-*O*-phosphate derivative, 5SMan α 1,2Man6P, as a potential inhibitor of a golgi α -1,2-mannosidase. The native Man α 1,2Man6P structure is contained in the biosynthetic intermediates of the *N*-glycans specifically expressed on lysosomal hydrolases [15]. Mannose-6-phosphates (Man6P) at the nonreducing ends of the *N*-glycans, exposed by the action of a golgi α -1,2-mannosidase toward the Man α 1,2Man6P structures [16], are specifically recognized by the cargo proteins bound for lysosomes [17]. Thus, the nondigestive property of 5SMan would allow 5SMan α 1,2Man6P to be a potential blocking agent against the distribution of lysosomal enzymes into lysosomes, though the mannosidase is unspecific for the phosphate group.

2. Results and Discussion

Ethyl 1,4-di-*O*-acetyl-5-thiomannopyranosyluronate **3** was synthesized from 1,4-diacetoxyl-1,3-butadiene **1** and ethyl thioacetate **2**, generated in situ from the anthracene

adduct, through hetero Diels-Alder and osmium oxidation reactions in a reported method [11, 12]. Prior to the reduction of the carboxylate moiety, compound **3DL** was subjected to Fischer methanolysis to give methyl (methyl 5-thiomannopyranosid)uronate **4DL** in 62%. **4DL**: R_f = 0.23 (CHCl₃-MeOH, 9 : 1), ¹H NMR (270 MHz, CDCl₃) δ 4.53 (d, 1H, J = 3.6 Hz, H-1), 4.22 (t, 1H, J = 3.6 Hz, H-2), 4.16 (t, 1H, J = 10 Hz, H-4), 3.79 (s, 1H, CO₂CH₃), 3.72 (dd, 1H, J = 3.6, 10 Hz, H-3), 3.67 (t, 1H, J = 10 Hz, H-5), 3.49 (s, 3H, OCH₃); ¹³C NMR (67.8 MHz, CDCl₃) δ 170.6, 87.2, 71.9, 70.8, 56.5, 53.0, 43.8; ESI (M + Na)⁺ 261. As attempted reduction of **4DL** with LiAlH₄ ended in decomposition, **4DL** was treated with NaBH₄, which has been occasionally successful in the reduction of sugar esters [18], to give methyl 5-thiomannopyranoside **5DL** in 77%. Spectral data of its tetraacetate were consistent with those reported for D-enantiomer [19]. Acetylation followed by acetylation of compound **5** produced per-O-acetyl-5-thiomannopyranose **6DL** in 85% yield, whose NMR data coincided with those reported for its D-enantiomer [20]. We thus obtained the protected racemic 5-thiomannose **6DL** from petrochemicals in six steps and 9% overall yield. The synthesis of the corresponding 5-thio-D-mannose derivative **6D** has been achieved in ten steps from D-mannose with 14% overall yield [20], showing that the merit of this study is the fewer synthetic steps and the use of petrochemicals for raw material. Although the low yield is a downside, the most impractical aspect of this method resided in the production of racemic compounds. To minimize the demerits, we next studied the use of the racemic 5-thiomannose derivative **6DL** for the synthesis of a disaccharide analog.



Per-O-acetyl-5-thiomannose **6DL** was derivatized to 1-O-trichloroacetimidate **7DL** in the same manner as that of D-enantiomer [21] in 80% yield. The glycosidation reaction of **7DL** toward a 3-O-benzyl-protected mannopyranoside **8** [22] was conducted with the conditions reported for the synthesis of a similar 5-thio-D-mannose-containing disaccharide, in which the 3-O-benzoyl group caused the predominant production of an orthoester derivative [21]. Use of the benzyl group for 3-O-protection prevented the orthoester formation and methyl 2-O-(5-thio- α -D and L-mannopyranosyl) mannopyranoside derivatives (**9D** : **9L** = 1 : 1) were obtained in 78% yield from the racemic glycosyl donor **7DL**. The diastereomers were easily separated by silica gel column chromatography. The stereochemistry of the diastereomers was confirmed by comparison of R_f values, $[\alpha]_D^{25}$, and ¹H NMR **9D**: R_f = 0.43 (hexane-ethyl acetate, 1 : 1); $[\alpha]_D^{25}$ +67.7° (c 0.73, CHCl₃); +66.8° for the same product synthesized from D-enantiomer of **7**; ¹H NMR (270 MHz, CDCl₃) δ 7.50-7.26 (m, 10H, Ph), 5.68 (s, 1H, PhCH), 5.67 (dd, 1H, J = 2.8, 4.0 Hz, H-2'), 5.45 (t, 1H, J = 10 Hz, H-4'), 5.37 (dd, 1H, J = 2.8, 10 Hz, H-3'), 5.00 (d, 1H, $J_{1',2'}$ 4.0 Hz, H-1'), 4.91-4.60 (dd, 2H, J = 12.2 Hz, PhCH₂), 4.66 (d, 1H, $J_{1,2}$ 1.3 Hz, H-1'), 4.88 (d, 1H, J = 12.2 Hz, PhCHH), 4.66 (d, 1H, J = 2.3 Hz, H-1), 4.62 (d, 1H, J = 12.2 Hz, PhCHH), 4.30 (dd, 1H, J = 5.6, 12.2 Hz, H-6a'), 4.26 (dd, 1H, J = 4.6, 10.2 Hz, H-6a), 4.17 (t, 1H, J = 9.6 Hz, H-4), 4.10 (dd, 1H,

J = 2.3, 3.0 Hz, H-2), 4.09 (dd, 1H, J = 3.6, 12.2 Hz, H-6b'), 3.97 (dd, 1H, J = 3.0, 9.6 Hz, H-3), 3.91 (t, 1H, J = 10.2 Hz, H-6b), 3.77 (ddd, 1H, J = 4.6, 9.6, 10.2, H-5), 3.45 (ddd, 1H, J = 3.6, 5.6, 10 Hz, H-5'), 3.37 (s, 3H, OCH₃), 2.12, 2.08, 2.04, 2.00 (s \times 4, 12H, COCH₃); ¹³C NMR (67.8 MHz, CDCl₃) δ 170.5, 169.75, 169.70, 169.5, 138.4, 137.5, 128.8, 128.2, 128.1, 127.4, 127.3, 126.1, 101.5, 101.0, 83.3, 79.5, 76.0, 75.8, 73.4, 70.6, 70.3, 69.5, 68.7, 63.8, 62.1, 54.8, 39.3, 21.0, 20.7, 20.6; HR-ESMS calcd for C₃₅H₄₂O₁₄SNa (M + Na)⁺ 741.2194, found 741.2282. **9L**: R_f = 0.35 (hexane-ethyl acetate, 1:1); $[\alpha]_D^{25}$ -72.9° (c 0.79, CHCl₃); ¹H NMR (270 MHz, CDCl₃) δ 7.52-7.26 (m, 10H, Ph), 5.69 (s, 1H, PhCH), 5.45-5.40 (m, 3H, H-2', H-3', H-4'), 4.84 (d, 1H, J = 12.2 Hz, PhCHH), 4.70 (d, 1H, J = 12.2 Hz, PhCHH), 4.69 (d, 1H, J = 3.3 Hz, H-1'), 4.68 (d, 1H, J = 1.5 Hz, H-1), 4.29 (dd, 1H, J = 1.5, 3.6 Hz, H-2), 4.25 (dd, 1H, J = 4.5, 10 Hz, H-6a), 4.13 (t, 1H, J = 9.6 Hz, H-4), 4.04 (dd, 1H, J = 4.6, 11.9 Hz, H-6a'), 3.96 (dd, 1H, J = 3.6, 9.6 Hz, H-3), 3.90 (t, 1H, J = 10 Hz, H-6b), 3.80-3.72 (m, 2H, H-5, H-5'), 3.59 (dd, 1H, J = 3.3, 11.9 Hz, H-6b'), 3.36 (s, 3H, OCH₃), 2.18, 2.01, 2.002, 1.996 (s \times 4, 12H, COCH₃); ¹³C NMR (67.8 MHz, CDCl₃) δ 170.5, 170.2, 169.7, 169.6 (C = O), 138.2, 137.6, 128.9, 128.3, 128.1, 127.6, 126.1 (Ph), 101.5 (PhCH), 98.3 (C-1), 80.4 (C-1'), 78.8 (C-3), 74.4, 74.1, 72.9 (PhCH₂), 71.5, 70.2, 68.9, 68.7, 64.0, 61.3 (C-6'), 54.9 (OCH₃), 38.7 (C-5'), 21.1, 20.7, 20.65, 20.6 (COCH₃); HR-ESMS calcd for C₃₅H₄₂O₁₄SNa (M + Na)⁺ 741.2194, found 741.2242. with those of **9D*** synthesized from 5-thio-D-mannopyranosyl donor **7D**. This is the first synthesis of an enantiomerically pure 5-thiosugar derivative from racemic petrochemicals.

As stated in the introductory section, we modified the synthesized disaccharide **9D** into 6-O-phosphate. The benzylidene group of compound **9D** was deprotected by hydrolysis (99%) to give compound **10**, which was then subjected to phosphorylation conditions giving regioselectively the di-O-benzyl 6-O-phosphate **11** in 58% yield. Deacetylation (**12** in 91%) followed by debenzylation gave compound **13** in 83% yield. **13**: R_f = 0.20 (iPrOH-H₂O, 3:1); $[\alpha]_D^{25}$ +45.1° (c 1.0, H₂O); ¹H NMR (270 MHz, D₂O) δ 8.46 (d, 1H, J = 1.5 Hz, H-1), 4.72 (d, 1H, $J_{1,2}$ 4.0 Hz, H-1'), 4.20 (dd, 1H, J = 2.8, 4.0 Hz, H-2'), 3.89-3.50 (m, 3H, H-2, H-6a, H-6b), 3.80 (dd, 1H, J = 3.3, 11.9 Hz, H-6a'), 3.75-3.50 (m, 6H, H-3, H-4, H-5, H-3', H-4', H-6b'), 3.26 (s, 3H, OCH₃), 2.98 (m, 1H, H-5'); ¹³C NMR (67.8 MHz, D₂O) δ 100.2, 86.8, 77.7, 72.5, 72.34, 72.2, 72.0, 70.6, 70.1, 66.7, 63.3, 60.9, 55.2, 44.6; ³¹P NMR (109.4 MHz, D₂O) δ 3.75.

3. Conclusion

We achieved the synthesis of an enantiomerically pure 5-thio-D-mannose-containing disaccharide derivative from racemic raw materials for the first time. To pursue the advantage of 5-thiosugar, we derivatized the disaccharide into 5SMan α 1,2Man6P, the analog of the partial structure of an intermediate oligosaccharide of the N-glycans on lysosomal enzymes. 5-thiopyranosides are in general



SCHEME 2: a. AcCl/MeOH ; b. $\text{NaBH}_4/\text{H}_2\text{O}$; c. (i) $\text{Ac}_2\text{O}-\text{Py}$, (ii) Ac_2O , H_2SO_4 ; d. (i) $\text{H}_2\text{NNH}_2 \cdot \text{AcOH}/\text{DME}$, (ii) K_2CO_3 , CCl_3CN ; e. **8**, $\text{TESOTf}/\text{CH}_2\text{Cl}_2$; f. 80% AcOH ; g. (i) $(\text{BnO})_2\text{PN}(\text{iPr})_2$, 1H-tetrazole, (ii) 30% H_2O_2 ; h. $\text{Et}_3\text{N}-\text{CH}_3\text{OH}$; i. 10% $\text{Pd}-\text{C}$, H_2 , $\text{AcOH}-\text{EtOH}$.

glycosidase-resistant and the analog is a potential inhibitor of a golgi α -1,2-mannosidase that cleaves $\text{Man}\alpha 1,2\text{Man}6\text{P}$ into $\text{Man}6\text{P}$.

Acknowledgments

This work was supported by the Grant of the 21st Century COE Program and a Grant-in-Aid for Scientific Research (no. 19655056) from Japanese Ministry of Education, Culture, Sports, Science, and Technology.

References

- [1] H. Yuasa and H. Hashimoto, "Replacing the ring oxygen of carbohydrates with sulfur: its biological and chemical consequences," *Reviews on Heteroatom Chemistry*, vol. 19, pp. 35–65, 1999.
- [2] I. Robina, P. Vogel, and Z. J. Witczak, "Synthesis and biological properties of monothiosaccharides," *Current Organic Chemistry*, vol. 5, no. 12, pp. 1177–1214, 2001.
- [3] I. Robina and P. Vogel, "The synthesis of disaccharides, oligosaccharides and analogues containing thiosugars," *Current Organic Chemistry*, vol. 6, no. 5, pp. 471–491, 2002.
- [4] H. Yuasa, O. Hindsgaul, and M. M. Palcic, "Chemical-enzymatic synthesis of 5'-thio-N-acetylactosamine: the first disaccharide with sulfur in the ring of the nonreducing sugar," *Journal of the American Chemical Society*, vol. 114, no. 14, pp. 5891–5892, 1992.
- [5] S. Mehta, J. S. Andrews, B. D. Johnston, B. Svensson, and B. M. Pinto, "Synthesis and enzyme inhibitor activity of novel glycosidase inhibitors containing sulfur and selenium," *Journal of the American Chemical Society*, vol. 117, no. 39, pp. 9783–9790, 1995.
- [6] H. Hashimoto, M. Kawanishi, and H. Yuasa, "Synthesis of methyl 5'-thio- α -isomaltoside via an acyclic monothioacetal

- and its behavior toward glucoamylase," *Chemistry: A European Journal*, vol. 2, no. 5, pp. 556–560, 1996.
- [7] C. M. Bladon, *Pharmaceutical Chemistry: Therapeutic Aspects of Biomolecules*, John Wiley & Sons, Chichester, UK, 2002.
- [8] H. Hashimoto, T. Fujimori, and H. Yuasa, "Synthesis of 5-thio-L-fucose and its inhibitory effect on fucosidase," *Journal of Carbohydrate Chemistry*, vol. 9, no. 5, pp. 683–694, 1990.
- [9] O. Tsuruta, H. Yuasa, and H. Hashimoto, "*p*-nitrophenyl 1,5-dithio- α -L-fucopyranoside: a novel sulfur based fucosidase inhibitor," *Bioorganic & Medicinal Chemistry Letters*, vol. 6, no. 16, pp. 1989–1992, 1996.
- [10] H. Yuasa, Y. Nakano, and H. Hashimoto, "Importance of ring sulfur and anomeric hydroxyl for inhibitory activity of 5-thiofucose toward α -fucosidases," *Carbohydrate Letters*, vol. 2, pp. 23–26, 1996.
- [11] D. Adam, A. A. Freer, N. W. Isaacs, G. W. Kirby, A. Littlejohn, and M. S. Rahman, "Synthesis of a thiashikimic acid derivative," *Journal of the Chemical Society, Perkin Transactions 1*, no. 10, pp. 1261–1264, 1992.
- [12] J. Prabhakaran, H. Lhermitte, J. Das, T. K. Sasi-Kumar, and D. S. Grierson, "The synthesis of a sulfone containing analogue of the esperamicin-A1 aglycone: a Hetero Diels-Alder approach," *Synlett*, no. 5, pp. 658–662, 2000.
- [13] M. Heras, M. Gulea, S. Masson, and C. Philouze, "A Hetero-Diels-Alder approach to phosphonothiashikimic acid and new thiaglycosides," *European Journal of Organic Chemistry*, no. 1, pp. 160–172, 2004.
- [14] R. J. Capon and J. K. MacLeod, "5-thio-D-mannose from the marine sponge *Clathria pyramida* (Lendenfeld). The first example of a naturally occurring 5-thiosugar," *Journal of the Chemical Society, Chemical Communications*, no. 15, pp. 1200–1201, 1987.
- [15] M. E. Taylor and K. Drickamer, *Introduction to Glycobiology*, Oxford University Press, Oxford, UK, 2003.
- [16] D. A. Lazzarino and C. A. Gabel, "Mannose processing is an important determinant in the assembly of phosphorylated high mannose-type oligosaccharides," *The Journal of Biological Chemistry*, vol. 264, no. 9, pp. 5015–5023, 1989.
- [17] J. J. Distler, J. Guo, G. W. Jourdain, O. P. Srivastava, and O. Hindsgaul, "The binding specificity of high and low molecular weight phosphomannosyl receptors from bovine testes," *The Journal of Biological Chemistry*, vol. 266, no. 32, pp. 21687–21692, 1991.
- [18] M. L. Wolfrom and K. Anno, "Sodium borohydride as a reducing agent in the sugar series," *Journal of the American Chemical Society*, vol. 74, no. 22, pp. 5583–5584, 1952.
- [19] H. Yuasa and H. Hashimoto, "Factors influencing stereoselectivity of sulfur oxidation: substituent effects on the oxidation of 5-thioglycopyranose derivatives," *Tetrahedron*, vol. 49, no. 40, pp. 8977–8998, 1993.
- [20] H. Yuasa, Y. Izukawa, and H. Hashimoto, "Synthesis of 5-thio-D-mannose," *Journal of Carbohydrate Chemistry*, vol. 8, no. 5, pp. 753–763, 1989.
- [21] B. D. Johnston and B. M. Pinto, "Synthesis of heteroanalogues of disaccharides as potential inhibitors of the processing mannosidase class I enzymes," *Carbohydrate Research*, vol. 310, no. 1–2, pp. 17–25, 1998.
- [22] Z. Szurmai, L. Balatoni, and A. Lipták, "Synthesis of some partially substituted methyl α -D- and phenyl 1-thio- α -D-mannopyranosides for the preparation of manno-oligosaccharides," *Carbohydrate Research*, vol. 254, pp. 301–309, 1994.