

Synthesis of 1,1-Organodiboronates via Rh(I)Cl-Catalyzed Sequential Regioselective Hydroboration of 1-Alkynes

Kohei Endo,* Munenao Hirokami, Takanori Shibata*

Department of Chemistry and Biochemistry, Advanced School of Science and Engineering, Waseda University, Shinjuku, Tokyo 169-8555, Japan

Fax +81(3)52863487; E-mail: kendo@aoni.waseda.jp; E-mail: tshibata@waseda.jp

Received 21 January 2009


Abstract: A Rh(I)Cl–DPPB-complex-catalyzed sequential hydroboration of aryl alkynes and aliphatic alkynes was achieved. The reaction proceeded with almost perfect regioselectivity to afford 1,1-organodiboronate compounds in moderate to good yield.

Key words: rhodium, hydroborations, alkynes, regioselectivity

1,1-Organodimetallic compounds are interesting synthetic intermediates and provide unique reactivity.^{1,2} We focused on the easy access to 1,1-organodiboronates, the borons of which attach to an sp^3 carbon atom, as the new approach to the preparation of 1,1-organodimetallic compounds.³ The boronate compounds can be readily converted into alcohols, amines, and carboxylic acids, and also used for the formation of C–C bond; thus 1,1-organodiboronates would be fascinating synthetic intermediates for the further functionalization.⁴ The precedents reported the formation of multiborylated compounds as a regioisomeric mixture via hydroboration or diboration.⁵ Another approach is the oxidative borylation of alkenylboronates affording 1,1-organodiboronate as a side product.⁶ Hiya-ma et al. reported the synthesis of 1,1-organodiborylated cyclopropanes and alkenes from 1,1-diboromo compounds.⁴ Here describes the one-step synthesis of 1,1-organodiboronates via Rh(I)Cl-catalyzed sequential and regioselective hydroboration.

In general, the hydroboration of 1-alkynes gives alkenylboronates; the sequential regioselective hydroboration of alkenylboronates using pinacolborane providing 1,1-organodiboronates has never been reported.⁷ The conventional reports concerning the hydroboration of C–C multiple bond provides a regioisomeric mixture and reduction product as well as tri- and tetraboronates; thus the control of the regioselectivity and the inhibition of overreaction via oxidative borylation are difficult to achieve. We focused on the Rh-catalyzed hydroboration, which generates Rh–C bond after the insertion of olefinic moiety into Rh–H bond (Scheme 1).

The generation of α -Rh,B intermediate is expected, which is the typical feature of the carbometallation reaction of alkenylboronates.¹ At one hand, the predominant generation of π -benzylic rhodium intermediate (β -Rh,B interme-



Scheme 1 Regioselective Rh-catalyzed hydroboration

diate) derived from vinylarenes would be another regioselectivity to 1,2-organodiboronates.

The numerous numbers of studies for Rh-catalyzed hydroboration of alkenes provide the efficient synthetic approaches to organoboronate compounds.^{8,9} In contrast, there are a few reports concerning on Rh-catalyzed hydroboration of 1-alkynes providing alkenylboronates.¹⁰ We hypothesized the Rh-catalyzed sequential and regioselective hydroboration of 1-alkynes to form 1,1-organodiboronates via the generation of α -Rh,B intermediate. The examination of various reaction conditions realized the regioselective formation of 1,1-organodiboronates. The reaction of **1a** and pinacolborane (HBpin, commercially available, 2.4 equiv)¹¹ was carried out in the presence of [RhCl(cod)]₂ (5 mol% Rh) and ligand (6 mol%) in DCE at room temperature (Scheme 2 and Table 1).

Among the various phosphine ligands, DPPB provided the high catalytic activity to give the desired 1,1-organodiboronate **2a** in 67% yield along with the formation of reduction product, 2-phenethylboronate, as a byproduct (entry 4). The appreciated feature of the present reaction is the almost perfect regioselectivity; the formation of α -Rh,B intermediate surpasses the formation of π -benzylic rhodium intermediate. The reaction in the presence of cationic and basic Rh complexes decreased the yield of product (entries 9 and 10).




Scheme 2 Sequential hydroboration of 1-alkyne

Table 1 Screening of Reaction Conditions from **1a** to **2a**

Entry	Rh complex ^a	Ligand ^b	Time (h)	Yield (%) ^c
1	[RhCl(cod)] ₂	DPPM ^d	48	11
2	[RhCl(cod)] ₂	DPPE ^e	24	4
3	[RhCl(cod)] ₂	DPPP ^f	48	25
4	[RhCl(cod)] ₂	DPPB ^g	24	67
5	[RhCl(cod)] ₂	DPPPent ^h	30	21
6	[RhCl(cod)] ₂	XANTPHOS ⁱ	24	6
7	[RhCl(cod)] ₂	Ph ₃ P ^j	24	21
8	[RhCl(cod)] ₂	P(OPh) ₃ ^j	48	40
9	[Rh(cod) ₂]BF ₄	DPPB	24	43
10	[Rh(OH)(cod)] ₂	DPPB	24	53

^a Amount of Rh used: 5 mol%.^b Amount of ligand used: 6 mol%.^c The yield was determined with 1,1,2,2-tetrachloroethane by the integration ratio of ¹H NMR analysis.^d 1,1-Bis(diphenylphosphino)methane.^e 1,2-Bis(diphenylphosphino)ethane.^f 1,3-Bis(diphenylphosphino)propane.^g 1,4-Bis(diphenylphosphino)butane.^h 1,5-Bis(diphenylphosphino)pentane.ⁱ 9,9-Dimethyl-4,5-bis(diphenylphosphino)xanthene.^j Amount of ligand used: 12 mol%.

Under the optimum reaction conditions in hand, the sequential hydroboration of terminal alkynes was examined (Table 2). The reaction of phenylacetylene (**1a**) gave the desired product **2a** in 62% isolated yield (entry 1). The reaction using Rh complex (1 mol% Rh) afforded the product **2a** in 58% yield. The use of aryethynes bearing electron-donating group on benzene ring gave the similar results; *p*- and *o*-tolylacetylenes (**1b,c**) provided the desired products **2b** and **2c** in 57% and 75% yield, respectively (entries 2 and 3). The reaction of *p*-, *m*-, and *o*-ethynylanisoles (**1d–f**) afforded the diboronates **2d–f** in moderate yield (entries 4–6). The use of aryethyne bearing halide substituent on benzene ring decreased the yield of product; 4-bromophenylacetylene (**1g**) provided the diborylated compound **2g** in 40% yield (entry 7). The reaction of 1-ethynynaphthalene (**1h**) gave the product **2h** in 68% yield (entry 8). 4-Ethynylbiphenyl (**1i**) provided the product **2i** in 55% yield (entry 9). The aliphatic alkynes could participate in the reaction. The reaction of **1j** gave the product **2j** in 59% yield (entry 10). Propargyl ether derivative **1k** afforded the product **2k** in 51% yield along with the formation of unidentified byproducts (entry 11). The bulky alkyne, 3,3-dimethylbut-1-yne (**1l**), decreased the yield of product **2l** (entry 12). The reaction of alkenylboronate was carried out (Scheme 3). The hydroboration of **3** with HBpin (1.2 equiv) in the presence of [RhCl(cod)]₂ (5 mol% Rh) and DPPB (6 mol%) gave **2a** in 75% yield. Thus, the present reaction from alkynes as substrate is considered to proceed via alkenylboronates. It is notable that the use of alkenylboronate **3** is in favor of

**Scheme 3** Rh(I)Cl-catalyzed hydroboration of alkenylboronate

the generation of α -Rh,B intermediate rather than π -benzylic β -Rh,B intermediate.¹²

The synthetic applications were examined for the present novel type of 1,1-organodiboronate. The selective homologation reaction enables the synthesis of a wide variety of boronate compounds.^{8k} The selective monohomologation of **2a** using chloromethylithium gave the corresponding product **4** in 64% yield (Scheme 4). The reaction using dichloromethylithium derivatives afforded the corresponding alkenylboronates **5a** and **5b**, respectively, in good yield (Scheme 5). This is the useful approach to the synthesis of multisubstituted alkenylboronates, which can participate in the Suzuki cross-coupling




























**Scheme 4** Homologation of **2a** using chloromethylithium**Scheme 5** Homologation of **2a** using dichloromethylithium derivatives**Scheme 6** Homologation of **2a** for allenylboronate synthesis and sequential oxidation

Table 2 Synthesis of Various 1,1-Organodiboronates^a

Entry	1-Alkyne	Product	Yield (%) ^b
1	1a 	2a 	62, 58 ^c (17, 16 ^c)
2	1b 	2b 	57 (24)
3	1c 	2c 	75 (16)
4	1d 	2d 	61 (23)
5	1e 	2e 	63 (19)
6	1f 	2f 	64 (18)
7	1g 	2g 	40 (24)
8	1h 	2h 	68 (14)
9	1i 	2i 	55 (20)
10	1j 	2j 	59 (12)
11 ^d	1k 	2k 	51
12 ^d	1l 	2l 	26

^a The reaction of alkyne and HBpin (2.4 equiv) was carried out in the presence of [RhCl(cod)]₂ (5 mol% Rh) and DPPB (6 mol%) in DCE at r.t. for 24 h.

^b The isolated yields are described. The yields of the reduction product of alkenylboronate are described in parentheses.

^c The yield using [RhCl(cod)]₂ (1 mol% Rh) and DPPB (1.2 mol%) is described.

^d Unidentified side products were obtained.

reaction. The reaction using 3-chloro-3-methylbut-1-ynyllithium provided the allenylboronate **6** in moderate yield (Scheme 6).^{4d} The allenylboronate **6** bearing an additional boryl group alpha to the C–C double bond was successively oxidized under the typical conditions to give the enone derivative **7**. This result suggested that the predominant oxidation of allenylboronate moiety proceeded to form enone bearing boronate moiety at α -carbon and subsequent isomerization and protonation took place to furnish enone **7** in 86% yield.

In conclusion, we succeeded in the sequential regioselective hydroboration of terminal alkynes for the facile synthesis of 1,1-organodiboronates in the presence of [RhCl(cod)]₂ and DPPB. The formation of α -Rh,B intermediate surpasses the formation of β -Rh,B intermediate which is the favorable regioselectivity of the Rh-catalyzed hydroboration of vinylarenes. The demonstration of synthetic utility of 1,1-organodiboronate showed the homologation reaction for the synthesis of 1,2-organodiboronate, di- and trisubstituted alkenylboronates, and allenylboro-

nate. The continuous studies are in progress, and the synthetic applications are in due course in our laboratory.

Typical Procedure of Rh(I)-Catalyzed Sequential Hydroboration

To a mixture of [RhCl(cod)]₂ (12.3 mg, 0.025 mmol, 5 mol% Rh), DPPB (25.6 mg, 0.06 mmol, 6 mol%) in DCE (1 mL) were added phenylacetylene (**1a**, 1.0 mmol, 102 mg) and pinacolborane (2.4 mmol, 308 mg, 2.4 equiv) at r.t. The reaction mixture was stirred at r.t. for 24 h and passed through a pad of SiO₂ with Et₂O (50 mL). The crude mixture was concentrated to dryness. The purification by SiO₂ column chromatography (5% EtOAc in hexane as eluent) gave the product **2a** in 62% yield (0.62 mmol) as colorless oil.

2-Phenyl-1,1-bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)ethane (**2a**)

Colorless oil. ¹H NMR (400 MHz, C₆D₆): δ = 7.32 (m, 2 H), 7.14 (m, 2 H), 7.03 (m, 1 H), 3.24 (d, *J* = 8.3 Hz, 2 H), 1.51 (t, *J* = 8.3 Hz, 1 H), 1.04 (s, 24 H). ¹³C NMR (100 MHz, C₆D₆): δ = 145.1, 128.8, 125.7, 82.9, 32.1, 24.9, 24.7. ¹¹B NMR (128 MHz, C₆D₆): δ = 34.2. IR (neat): 2978, 1454, 1319, 1139, 971, 851 cm⁻¹. HRMS–FAB (+): *m/z* calcd for C₂₀H₃₃B₂O₄⁺: 359.2559 [M + H]⁺; found: 359.2567 [M + H]⁺.

1,1-Bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-*p*-tolyl-ethane (**2b**)

Colorless oil. ¹H NMR (400 MHz, C₆D₆): δ = 6.98 (m, 2 H), 6.70 (m, 2 H), 2.97 (d, *J* = 8.0 Hz, 2 H), 1.84 (s, 3 H), 1.25 (t, *J* = 8.0 Hz, 1 H), 0.77 (s, 24 H). ¹³C NMR (100 MHz, C₆D₆): δ = 124.1, 134.6, 129.0, 128.7, 82.9, 31.7, 27.7, 24.6, 21.0. ¹¹B NMR (128 MHz, C₆D₆): δ = 33.7. IR (neat): 2978, 1379, 1319, 1140, 970, 854 cm⁻¹. HRMS–FAB (+): *m/z* calcd for C₂₁H₃₄B₂O₄⁺: 372.2643; found: 372.2629.

1,1-Bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-*o*-tolyl-ethane (**2c**)

Pale yellow oil. ¹H NMR (400 MHz, C₆D₆): δ = 7.21 (m, 1 H), 6.77 (m, 3 H), 2.93 (d, *J* = 8.5 Hz, 2 H), 1.95 (s, 3 H), 1.26 (t, *J* = 8.5 Hz, 1 H), 0.78 (s, 24 H). ¹³C NMR (100 MHz, C₆D₆): δ = 142.9, 136.2, 130.2, 128.7, 125.9, 125.8, 82.9, 29.0, 24.8, 19.5. ¹¹B NMR (128 MHz, C₆D₆): δ = 34.5. IR (neat): 2978, 1464, 1315, 1141, 971, 852 cm⁻¹. HRMS–FAB (+): *m/z* calcd for C₂₁H₃₅B₂O₄⁺: 373.2716 [M + H]⁺; found: 373.2733 [M + H]⁺.

2-*p*-Methoxyphenyl-1,1-bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)ethane (**2d**)

Colorless oil. ¹H NMR (400 MHz, C₆D₆): δ = 7.24 (m, 2 H), 6.77 (m, 2 H), 3.33 (s, 3 H), 3.22 (d, *J* = 8.4 Hz, 2 H), 1.50 (t, *J* = 8.4 Hz, 1 H), 1.07 (s, 12 H), 1.06 (s, 12 H). ¹³C NMR (100 MHz, C₆D₆): δ = 158.3, 137.2, 129.6, 113.9, 82.9, 54.8, 31.3, 24.9, 24.8. ¹¹B NMR (128 MHz, C₆D₆): δ = 34.0. IR (neat): 2978, 1463, 1301, 1140, 972, 854 cm⁻¹. HRMS–FAB (+): *m/z* calcd for C₂₁H₃₄B₂O₅: 388.2592; found: 388.2552.

2-*m*-Methoxyphenyl-1,1-bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)ethane (**2e**)

Pale yellow oil. ¹H NMR (400 MHz, C₆D₆): δ = 6.81 (m, 1 H), 6.70 (m, 2 H), 6.41 (m, 1 H), 3.08 (s, 3 H), 2.97 (d, *J* = 8.4 Hz, 2 H), 1.75 (t, *J* = 8.4 Hz, 1 H), 0.79 (s, 12 H), 0.78 (s, 12 H). ¹³C NMR (100 MHz, C₆D₆): δ = 160.2, 146.7, 129.3, 121.1, 114.2, 111.6, 82.9, 54.6, 32.3, 24.9, 24.8. ¹¹B NMR (128 MHz, C₆D₆): δ = 34.0. IR (neat): 2978, 1489, 1317, 1140, 972, 850 cm⁻¹. HRMS–FAB (+): *m/z* calcd for C₂₁H₃₄B₂O₅: 388.2592; found: 388.2554.

2-*o*-Methoxyphenyl-1,1-bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)ethane (**2f**)

Colorless oil. ¹H NMR (400 MHz, C₆D₆): δ = 6.99 (m, 1 H), 6.54 (m, 1 H), 6.27 (m, 1 H), 5.99 (m, 1 H), 2.91 (d, *J* = 8.0 Hz, 2 H), 2.75 (s, 3 H), 1.28 (t, *J* = 8.0 Hz, 1 H), 0.55 (s, 12 H), 0.54 (s, 12 H). ¹³C NMR (100 MHz, C₆D₆): δ = 158.1, 133.2, 130.0, 126.9, 120.2, 110.2, 82.8, 54.6, 26.9, 24.9, 24.8. ¹¹B NMR (128 MHz, C₆D₆): δ = 33.8. IR (neat): 2978, 1493, 1317, 1140, 972, 852 cm⁻¹. HRMS–FAB (+): *m/z* calcd for C₂₁H₃₄B₂O₅: 388.2592; found: 388.2594.

2-*p*-Bromophenyl-1,1-bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)ethane (**2g**)

Pale yellow oil. ¹H NMR (400 MHz, C₆D₆): δ = 7.27 (m, 2 H), 6.93 (m, 2 H), 3.03 (d, *J* = 8.4 Hz, 2 H), 1.35 (t, *J* = 8.4 Hz, 1 H), 1.02 (s, 24 H). ¹³C NMR (100 MHz, C₆D₆): δ = 144.0, 131.3, 130.6, 119.5, 83.0, 31.4, 24.9, 24.7. ¹¹B NMR (128 MHz, C₆D₆): δ = 34.4. IR (neat): 2978, 1486, 1317, 1072, 971, 852 cm⁻¹. HRMS–FAB (+): *m/z* calcd for C₂₀H₃₂B₂BrO₄⁺: 437.1665 [M + H]⁺; found: 437.1678 [M + H]⁺.

2-(Naphthalen-1-yl)-1,1-bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)ethane (**2h**)

Colorless oil. ¹H NMR (400 MHz, C₆D₆): δ = 8.12 (m, 1 H), 7.58 (m, 2 H), 7.47 (m, 1 H), 7.23 (m, 1 H), 7.15 (m, 2 H), 3.63 (d, *J* = 8.0 Hz, 2 H), 1.64 (t, *J* = 8.0 Hz, 1 H), 0.98 (s, 12 H), 0.97 (s, 12 H). ¹³C NMR (100 MHz, C₆D₆): δ = 140.9, 134.4, 132.6, 128.8, 126.7, 125.8, 125.6, 125.5, 125.4, 124.5, 83.0, 28.9, 24.9, 24.8. ¹¹B NMR (128 MHz, C₆D₆): δ = 33.8. IR (neat): 2978, 1378, 1323, 1140, 972, 849 cm⁻¹. HRMS–FAB (+): *m/z* calcd for C₂₄H₃₄B₂O₄: 408.2643; found: 408.2614.

2-*p*-Biphenyl-1,1-bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)ethane (**2i**)

Pale yellow solid; mp 87 °C. ¹H NMR (400 MHz, C₆D₆): δ = 7.43 (m, 6 H), 7.15 (m, 3 H), 3.29 (d, *J* = 8.3 Hz, 2 H), 1.56 (t, *J* = 8.3 Hz, 1 H), 1.07 (s, 12 H), 1.06 (s, 12 H). ¹³C NMR (100 MHz, C₆D₆): δ = 144.3, 141.9, 138.8, 129.2, 128.9, 127.3, 127.1, 127.0, 83.0, 31.8, 24.9, 24.7. ¹¹B NMR (128 MHz, C₆D₆): δ = 34.1. IR (neat): 2977, 1486, 1319, 971, 842 cm⁻¹. HRMS–FAB (+): *m/z* calcd for C₂₆H₃₆B₂O₄: 434.2800; found: 434.2795.

5-Phenyl-1,1-bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pentane (**2j**)

Colorless oil. ¹H NMR (400 MHz, C₆D₆): δ = 7.27 (m, 2 H), 7.14 (m, 3 H), 2.58 (t, *J* = 7.6 Hz, 2 H), 1.60 (m, 4 H), 1.32 (m, 2 H), 1.21 (s, 24 H), 0.72 (t, *J* = 7.6 Hz, 1 H). ¹³C NMR (100 MHz, C₆D₆): δ = 142.9, 128.4, 128.1, 125.4, 82.9, 36.7, 32.1, 31.3, 25.5, 24.8, 24.5. ¹¹B NMR (128 MHz, C₆D₆): δ = 33.9. IR (neat): 2978, 1371, 1315, 1142, 970, 850 cm⁻¹. HRMS–FAB (+): *m/z* calcd for C₂₃H₃₈B₂O₄: 400.2956; found: 400.2942.

3-(*tert*-Butyldimethylsiloxy)-3-phenyl-1,1-bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)propane (**2k**)

Pale yellow oil. ¹H NMR (400 MHz, C₆D₆): δ = 7.51 (m, 2 H), 7.17 (m, 2 H), 7.04 (m, 1 H), 4.87 (t, *J* = 7.6 Hz, 1 H), 2.47 (m, 1 H), 2.34 (m, 1 H), 1.30 (t, *J* = 7.6 Hz, 1 H), 1.08 (m, 24 H), 0.99 (s, 9 H), 0.19 (s, 3 H), 0.00 (m, 3 H). ¹³C NMR (100 MHz, C₆D₆): δ = 146.4, 128.3, 127.2, 126.6, 84.84, 84.82, 77.6, 38.1, 26.2, 25.1, 24.9, 24.8, 24.7, 18.5, −4.3, −4.5. ¹¹B NMR (128 MHz, C₆D₆): δ = 34.0. IR (neat): 2978, 1389, 1319, 1092, 970, 849 cm⁻¹. HRMS–FAB (+): *m/z* calcd for C₂₇H₄₇B₂O₅⁺: 501.3373 [M − 1]⁺; found: 501.3397 [M − 1]⁺.

3,3-Dimethyl-1,1-bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)butane (**2l**)

White solid; mp 67 °C. ¹H NMR (400 MHz, C₆D₆): δ = 1.53 (d, *J* = 6.8 Hz, 2 H), 1.23 (s, 24 H), 0.84 (s, 9 H), 0.74 (t, *J* = 6.8 Hz, 1

H). ^{13}C NMR (100 MHz, C_6D_6): δ = 82.3, 39.2, 31.5, 29.1, 24.7, 24.6. ^{11}B NMR (128 MHz, C_6D_6): δ = 33.9. IR (neat): 2978, 1389, 1309, 1144, 968, 866 cm^{-1} . ESI-HRMS (+): m/z calcd for $\text{C}_{18}\text{H}_{36}\text{B}_2\text{NaO}_4^+$: 361.2692 $[\text{M} + \text{Na}]^+$; found: 361.2686 $[\text{M} + \text{Na}]^+$.

Acknowledgment

K.E. thanks the Teijin Pharma Award in Synthetic Organic Chemistry, Japan and Waseda University Grant for Special Research Projects.

Reference and Notes

- (1) (a) *Handbook of Functionalized Organometallics*, Vol. 1 and 2; Knochel, P., Ed.; Wiley-VCH: Weinheim, **2005**. (b) Hirai, A.; Nakamura, M.; Nakamura, E. *J. Am. Chem. Soc.* **1999**, *121*, 8665. (c) Normant, J. F. *Acc. Chem. Res.* **2001**, *34*, 640. (d) Dembitsky, V. M.; Ali, H. A.; Srebnik, M. *Appl. Organomet. Chem.* **2003**, *17*, 327.
- (2) The 1,1-organodiboronate attached to sp^3 -carbon center was obtained as a byproduct, see: McIsaac, D. I.; Geier, S. J.; Vogels, C. M.; Decken, A.; Westcott, S. A. *Inorg. Chim. Acta* **2006**, *359*, 2771.
- (3) The diboration reaction of alkynylboronate or alkenylboronate was reported, which provides a mixture of 1,1,1- and 1,1,2-alkenyltriborylated or 1,1,1- and 1,1,2-alkyltriborylated compounds along with the formation of other types of diborylated compounds, see: Nguyen, P.; Coapes, R. B.; Woodward, A. D.; Taylor, N. J.; Burke, J. M.; Howard, J. A. K.; Marder, T. B. *J. Organomet. Chem.* **2002**, *652*, 77.
- (4) (a) Hata, T.; Kitagawa, H.; Masai, H.; Kurahashi, T.; Shimizu, M.; Hiyama, T. *Angew. Chem. Int. Ed.* **2001**, *40*, 790. (b) Kurahashi, T.; Hata, T.; Masai, H.; Kitagawa, H.; Shimizu, M.; Hiyama, T. *Tetrahedron* **2002**, *58*, 6381. (c) Shimizu, M.; Nakamaki, C.; Shimono, K.; Schelper, M.; Kurahashi, T.; Hiyama, T. *J. Am. Chem. Soc.* **2005**, *127*, 12506. (d) Shimizu, M.; Schelper, M.; Nagao, I.; Shimoto, K.; Kurahashi, T.; Hiyama, T. *Chem. Lett.* **2006**, *35*, 1222.
- (5) The hydroboration of alkynes using $\text{BH}_3\cdot\text{THF}$ complex was reported to give a mixture of monoborylated, 1,1-diborylated, and 1,2-diborylated compounds along with the formation of reduction product, see: (a) Pasto, D. J. *J. Am. Chem. Soc.* **1964**, *86*, 3039. (b) Rao, V. V. R.; Agarwal, S. K.; Mehrotra, I.; Devaprabhakara, D. *J. Organomet. Chem.* **1979**, *166*, 9.
- (6) The diboration reaction gave the regioisomeric mixture of products, see: Coapes, R. B.; Souza, F. E. S.; Thomas, R. L.; Hall, J. J.; Marder, T. B. *Chem. Commun.* **2003**, 614.
- (7) (a) Brown reported the sequential hydroboration of alkynes using 9-BBN to obtain 1,1-organodiborane intermediates, which was readily converted into primary alcohol via

protonolysis and oxidation; see: Brown, H. C.; Scouten, C. G.; Liotta, R. *J. Am. Chem. Soc.* **1979**, *101*, 96.

- (b) Matteson reported the stepwise synthesis of 1,1-organodiboronate via the hydroboration of 1-hexyne using BCl_3 and TMS. The subsequent reaction with 1,3-propanediol afforded the corresponding 1,1-organodiboronate, see: Soundararajan, R.; Matteson, D. S. *Organometallics* **1995**, *14*, 4157.
- (8) Rh(I)-catalyzed hydroboration of alkenes, see: (a) Hayashi, T. In *Comprehensive Asymmetric Catalysis*, Vol. 1; Jacobsen, E. N.; Phaltz, A.; Yamamoto, H., Eds.; Springer: New York, **1999**, 349. (b) Hayashi, T.; Matsumoto, Y.; Ito, Y. *J. Am. Chem. Soc.* **1989**, *111*, 3426. (c) Westcott, S. A.; Blom, H. P.; Marder, T. B.; Baker, R. T. *J. Am. Chem. Soc.* **1992**, *114*, 8863. (d) Brown, J. M.; Hulmes, D. E.; Layzell, T. P. *J. Chem. Soc., Chem. Commun.* **1993**, 1673. (e) Tucker, C. E.; Davidson, J.; Knochel, P. *J. Org. Chem.* **1992**, *57*, 3482. (f) Doucet, H.; Fernandez, E.; Layzell, T. P.; Brown, J. M. *Chem. Eur. J.* **1999**, *5*, 1320. (g) Chen, A.; Ren, L.; Crudden, C. M. *J. Org. Chem.* **1999**, *64*, 9704. (h) Demay, S.; Volant, F.; Knochel, P. *Angew. Chem. Int. Ed.* **2001**, *40*, 1235. (i) Murata, M.; Kawakita, K.; Asana, T.; Watanabe, S.; Masuda, Y. *Bull. Chem. Soc. Jpn.* **2002**, *75*, 825. (j) Crudden, C. M.; Edwards, D. *Eur. J. Org. Chem.* **2003**, 4695. (k) Crudden, C. M.; Hleba, Y. B.; Chen, A. C. *J. Am. Chem. Soc.* **2004**, *126*, 9200. (l) Segarra, A. M.; Daura-Oller, E.; Claver, C.; Poblet, J. M.; Bo, C.; Fernández, E. *Chem. Eur. J.* **2004**, *10*, 6456. (m) Moteki, S. A.; Wu, D.; Chandra, K. L.; Reddy, D. S.; Takacs, J. M. *Org. Lett.* **2006**, *8*, 3097. (n) Carroll, A.-M.; O'Sullivan, T. P.; Guiry, P. J. *Adv. Synth. Catal.* **2005**, *347*, 609. (o) Edwards, D. R.; Hleba, Y. B.; Lata, C. J.; Calhoun, L. A.; Crudden, C. M. *Angew. Chem. Int. Ed.* **2007**, *46*, 7799.
 - (9) (a) Endo, K.; Hirokami, M.; Shibata, T. *Organometallics* **2008**, *27*, 5390. (b) Endo, K.; Hirokami, M.; Takeuchi, K.; Shibata, T. *Synlett* **2008**, 3231.
 - (10) (a) Lee, T.; Baik, C.; Jung, I.; Song, K. H.; Kim, S.; Kim, D.; Kang, S. O.; Ko, J. *Organometallics* **2004**, *23*, 4569. (b) Ohmura, T.; Yamamoto, Y.; Miyaura, N. *J. Am. Chem. Soc.* **2000**, *122*, 4990.
 - (11) Both of the commercially available HBpin and purified HBpin according to the literature gave the almost same results; the impurities in HBpin was suggested by: Shimada, S.; Batsanov, A. S.; Howard, J. A. K.; Marder, T. B. *Angew. Chem. Int. Ed.* **2001**, *40*, 2168.
 - (12) The regioselectivity was confirmed by ^1H NMR, HMQC, and HMBC analyses. The crude products did not include 1,2-diboronate. The oxidation of product **2a** under $\text{H}_2\text{O}_2/\text{NaOH}$ conditions did not give 1-phenyl-1,2-ethanediol derived from 1,2-diboronate at all (unidentified products were obtained). The oxidation of 1,2-diboration product was reported, see: Trudeau, S.; Morgan, J. B.; Shrestha, M.; Morken, J. P. *J. Org. Chem.* **2005**, *70*, 9538.

Copyright of Synlett is the property of Georg Thieme Verlag Stuttgart and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.