

Research Article

Directed Metalation of Heterocycles, 5-Methoxy-2-phenyloxazol-4-yl lithium: An Approach to α,β -Dehydroamino Acids

Sosale Chandrasekhar and Malempati Srimannarayana

Department of Organic Chemistry, Indian Institute of Science, Bangalore 560 012, India

Correspondence should be addressed to Sosale Chandrasekhar, sosale@orgchem.iisc.ernet.in

Received 1 July 2009; Accepted 16 September 2009

Academic Editor: Kirpal Bisht

Copyright © 2010 S. Chandrasekhar and M. Srimannarayana. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

5-Methoxy-2-phenyloxazole was deprotonated at C₄ (by *n*-BuLi or LDA, in THF at -78°C). The resulting anion was generally unreactive to alkylation (except methylation with MeI-TMEDA) but added to PhCHO and Me₂CHCHO. The alcohols thus produced dehydrated and ring opened in acid, to the corresponding α,β -dehydroamino acids in moderate overall yields.

1. Introduction


Directed metalation (sometimes also “*ortho*-metalation”) is now firmly established as an important synthetic strategy in aromatic chemistry [1]. This involves deprotonation at a site *ortho* to a directing group such as OMe or NMe₂; this is assisted either by coordination of the counterion of the attacking base with the heteroatom lone pairs, or by electron withdrawal by the heteroatoms (or both). Thus, the formation of the anionic product is facilitated both kinetically and thermodynamically, with its subsequent electrophilic reactions leading to products not easily accessed otherwise.

Interestingly, however, heteroaromatic examples of directed metalation have been relatively scarce [2]. In particular, we have been interested in the possibility that the analogous reaction in the case of 5-methoxy-2-phenyloxazole (**1** \rightarrow **I**, Scheme 1) would define a novel approach to α -amino acids, via hydrolysis of the resulting C₄ substituted derivatives (Scheme 1). The 5-methoxy group would then not only function as a directing group but also maintain C₅ as a masked carboxyl centre. We report in what follows preliminary results from our study.


2. Results and Discussion

5-Methoxy-2-phenyloxazole (**1**) [3, 4] was treated with an equivalent of *n*-butyllithium (*n*-BuLi) in THF at -78°C for 3 hours, and the reaction mixture quenched with D₂O. ¹H NMR showed evidence of deuterium incorporation (88%), as seen by the disappearance of the C₄-H resonance at δ 5.80 (cf. **2a**). When the deprotonation was followed by treatment with methyl iodide, no reaction was observed; however, when the methylation was conducted in the presence of one equivalent of *N,N,N',N'*-tetramethylethylenediamine (TMEDA, -70 to $-40^{\circ}\text{C}/3\text{ h}$), 5-methoxy-4-methyloxazole (**2b**) was formed in 63% yield. These reactions also occurred in lower yields with lithium diisopropylamide (LDA, *vide infra*) as base [5].

These results clearly indicate the formation and intermediacy of 5-methoxy-2-phenyloxazol-4-yl lithium (**I**). Intriguingly, however, this did not react with longer chain alkylating agents, for example, *n*-BuI, *n*-C₇H₁₅Br, PhCH₂Br, and allyl bromide. (**I** could also be generated with LDA and methylated with MeI and added hexamethylphosphoramide (HMPA).) It seems likely that TMEDA and HMPA act by chelating the lithium counterion of **I**, thus breaking down aggregates into more reactive monomeric and dimeric forms.



SCHEME 1: The directed metalation strategy as applied to 5-methoxy-2-phenyloxazole (**1**), shown reacting with *n*-BuLi to form the oxazolylithium **I**; this reacted with D₂O to form **2a** (R = D, 88%) and with MeI-TMEDA to form **2b** (R = Me, 63%).



		Yield (%)	
	R	3	5
a	Ph	54	52
b	Me ₂ CH	65	78

SCHEME 2: Addition of the oxazolylithium **I** to benzaldehyde and iso-butylaldehyde to form the alcohols **3**, and their acid catalyzed rearrangement, ring cleavage, and hydrolysis to furnish the corresponding *N*-benzoyl dehydroamino acids **5**.

However, the oxazolylithium **I** was found to add to benzaldehyde and iso-butylaldehyde to furnish the expected alcohols **3**, in moderate yields (Scheme 2). Interestingly, **3** underwent rearrangement with ring cleavage and hydrolysis in dilute acid, to form the corresponding *N*-benzoyl α,β -dehydroamino acids **5** in fair yields [5].

A probable mechanism is shown, with the double bond geometry in **5** being known to be variable [6, 7]. (The corresponding esters **4** were isolated and characterized spectrally. 4-Alkylideneoxazol-5-ones reportedly fragment similarly [8].) Dehydroamino acids have gained importance in recent times, particularly as immediate precursors for homochiral amino acids [9, 10].

2.1. Experimental: General Remarks. ν_{\max} are in cm⁻¹ and *J* are in Hz; δ_{H} were recorded at 300 MHz and δ_{C} at 75 MHz. **Metalation and methylation of oxazole 1.** A stirred mixture of *n*-BuLi (1.6 mmol in 1.0 mL hexane) and TMEDA (1.6 mmol) at -78°C was treated dropwise with a solution of

the oxazole [3, 4] (1 mmol in 3.0 mL THF). After 3 hours the mixture was warmed to -40°C, treated with MeI (1.6 mmol), and allowed to reach room temperature. Work-up with EtOAc and satd. NH₄Cl, followed by chromatography (SiO₂ eluting with 20% EtOAc-hexane), led to the 4-Me derivative **1b** (63%); ν_{\max} 1663; δ_{H} 2.12 (s, 3H), 3.99 (s, 3H), 7.37–7.44 (m, 3H), 7.89–7.93 (m, 2H); δ_{C} 10.1, 61.1, 113.1, 125.3, 127.7, 128.6, 129.4, 152.0, 155.0; HRMS 190.0856 (Calcd. for C₁₁H₁₁NO₂ + H 190.0868). *iso*-Butylaldehyde adduct **3b**: ν_{\max} 3410, 2957, 1654; δ_{H} 0.88 (d, 3H, *J* 6.6), 1.07 (d, 3H, *J* 6.6), 2.09–2.16 (m, 1H), 2.53–2.55 (m, 1H), 4.00 (s, 3H), 4.26–4.31 (m, 1H), 7.38–7.41 (m, 3H), 7.89–7.91 (m, 2H); δ_{C} 18.3, 18.8, 33.6, 60.8, 71.4, 118.3, 125.5, 128.6, 129.6, 152.2, 154.7; HRMS 270.1110 (Calcd. for C₁₄H₁₇NO₃ + Na 270.1106). *N*-Benzoyl 2-amino-4-methylpent-2-enoic acid (**5b**): ν_{\max} 3267, 1697, 1640; δ_{H} 1.09 (d, 6H, *J* 6.6), 2.60–2.80 (m, 1H), 6.73 (d, 1H, *J* 10.5), 7.44–7.54 (m, 3H), 7.79–7.87 (m, 2H); δ_{C} 21.5, 28.4, 122.8, 127.4, 128.6, 132.0, 132.2, 133.8, 147.2, 166.4, 168.9, 170.8; HRMS 256.0960 (Calcd. for C₁₃H₁₅NO₃ + Na 256.0950).

3. Conclusions

We have demonstrated the feasibility of the directed metalation strategy in the case of a heterocycle that is a potential amino acid synthon. The resulting carbanion is generally unreactive towards several alkylating agents (except MeI). However, addition to aldehydes, followed by rearrangement and hydrolysis in acid, led to α,β -dehydroamino acids in fair yields in two cases.

Acknowledgment

The authors are grateful to CSIR (New Delhi) for generous fellowship support to the second author.

References

- [1] V. Snieckus, "Directed ortho metalation. Tertiary amide and O-carbamate directors in synthetic strategies for polysubstituted aromatics," *Chemical Reviews*, vol. 90, no. 6, pp. 879–933, 1990.
- [2] R. Chinchilla, C. Nájera, and M. Yus, "Metalated heterocycles and their applications in synthetic organic chemistry," *Chemical Reviews*, vol. 104, no. 5, pp. 2667–2722, 2004.
- [3] G. Labbé, A.-M. Ilisiu, W. Dehaen, and S. Toppet, "Synthesis and thermolysis of 5-azido-4-formyloxazoles," *Journal of the Chemical Society, Perkin Transactions 1*, no. 19, pp. 2259–2261, 1993.
- [4] A. G. Griesbeck, S. Bondock, and J. Lex, "Synthesis of erythro- α -amino- β -hydroxy carboxylic acid esters by diastereoselective photocycloaddition of 5-methoxyoxazoles with aldehydes," *Journal of Organic Chemistry*, vol. 68, no. 26, pp. 9899–9906, 2003.
- [5] M. Srimannarayana, *Topics in synthetic methodology: from heterocycles to hydride transfers*, Ph.D. thesis, Indian Institute of Science, Bangalore, India, 2009.
- [6] K. Brocklehurst, R. P. Bywater, R. A. Palmer, and R. Patrick, "The crystal structure of the stable isomer of α -benzamidocinnamic acid: the influence of *cis-trans*-isomerism on the kinetics of the hydrolysis of the products of interaction of α -chymotrypsin with the isomeric 4-benzylidene-2-phenyl- Δ^2 -oxazolin-5-ones," *Journal of the Chemical Society D*, no. 12, pp. 632–633, 1971.
- [7] V. Busetti, D. Ajò, and A. Vittadini, "Structure of (Z)-N-benzoyl- α,β -dehydroleucine," *Acta Crystallographica Section C*, vol. 42, no. 9, pp. 1178–1181, 1986.
- [8] H. Takagaki, S. Tanabe, M. Asaoka, and H. Takei, "Reaction of 2-phenyl-5-trimethylsiloxyoxazole with carbonyl compounds or acetals: synthesis of α,β -dehydro- α -amino acid derivatives," *Chemistry Letters*, vol. 8, no. 4, pp. 347–350, 1979.
- [9] K. F. W. Hekking, D. C. J. Waalboer, M. A. H. Moelands, F. L. van Delft, and F. P. J. T. Rutjes, "A ring-closing metathesis approach to cyclic α,β -dehydroamino acids," *Advanced Synthesis & Catalysis*, vol. 350, no. 1, pp. 95–106, 2008.
- [10] F. Giacomina, A. Meetsma, L. Panella, L. Lefort, A. H. M. de Vries, and J. G. de Vries, "High enantioselectivity is induced by a single monodentate phosphoramidite ligand in iridium-catalyzed asymmetric hydrogenation," *Angewandte Chemie International Edition*, vol. 46, no. 9, pp. 1497–1500, 2007.