

## Research Article

# A Novel and Efficient Synthesis of Intermediates for the Preparation of Fexofenadine

Jiangang Mao,<sup>1</sup> Haining Gu,<sup>2</sup> and Pengfei Zhang<sup>1</sup>

<sup>1</sup> College of Material, Chemistry and Chemical Engineering, Hangzhou Normal University, Xiasha Campus, Hangzhou, Zhejiang 310036, China

<sup>2</sup> Analytical Center, Zhejiang University, Xixi Campus, Hangzhou, Zhejiang 310036, China

Correspondence should be addressed to Pengfei Zhang, pfzhang@hznu.edu.cn

Received 22 April 2008; Accepted 26 June 2008

An efficient and environmentally friendly synthetic approach for the preparation of fexofenadine was developed. The method involves hydrogenation and oxygenation reaction and has good atom utilization, all the starting material is nontoxic.

Copyright © 2008 Jiangang Mao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

Fexofenadine (**4**), the carboxylic acid metabolite of terfenadine and the actual pharmacologically active compound [1–3], is a selective histamine H1 receptor antagonist, clinically effective in the treatment of seasonal allergic rhinitis, and chronic idiopathic urticaria as a first-line therapeutic agent [3–5]. Several synthetic routes to fexofenadine have been reported [6–15]. However, a review of the literature, including patents, indicated the absence of both environmentally benign and relatively low-cost method for its synthesis. A summary of known representative procedures is shown in Schemes 1 [14, 15] and 2 [16, 17], these methods utilize either reaction of poor regioselectivity or unwieldy and expensive reagents. Although, Scheme 2 [16, 17] is a relatively low-cost and convenient operation method and also solves the problem of purification caused by the formation of the ortho isomer (step 1, Scheme 2), the mercury byproduct is harmful and difficult to remove. In connection with our systematic studies on the process of fexofenadine, we have developed an environmentally friendly modified method for synthesis intermediates toward its preparation (Scheme 3).

Hydrogenation of alkyne **5** on a modified Lindlar catalyst system (5% Pd/CaCO<sub>3</sub> and a small amount of tertiary amine) to give alkene **7** (95% yield). Alkene **7** was obtained only 70% yield if the commercial Lindlar catalyst obtained from the Catalysis Institute of Zhejiang University was used because of the production of saturate ester as a byproduct. Alkene **7** was subsequently oxidized to the methyl ester **6**

with oxygen at room temperature, using the catalytic system of Mn(OAc)<sub>2</sub>, NaBH<sub>4</sub>, and Schiff-base ligand (**B**) [18, 19]; subsequently, hydrolysis of methyl ester **6** with hydrochloric acid afforded fexofenadine **4**.


In summary, this synthesis has good atom utilization, the starting material is inexpensive and readily available commercially, the new method has the advantage of good yield and of being environmental benign. This procedure seems to be amenable to pharmaceutical manufacture.

## 2. Experimental Section


Melting points are uncorrected. Infrared spectra were obtained on an IR-408 instrument. <sup>1</sup>H NMR spectra were recorded on a Bruker 400 (400 MHz) instrument using CDCl<sub>3</sub> as the solvent with TMS as internal standard. All other chemicals were reagent grade.

### 2.1. Synthesis of 4-{4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-butenyl}- $\alpha,\alpha$ -dimethyl-benzeneacetic acid methyl ester (**7**)

A solution of compound **5** (4.45 g, 8.99 mmol) in 300 mL ethanol and 0.38 g of Lindlar catalyst (palladium:CaCO<sub>3</sub> 1:20) was added to a 500-mL autoclave under a nitrogen atmosphere, and quinoline (1.0 mL) was slowly added dropwise to the stirred mixture at 20–25°C. Subsequently,



SCHEME 1




SCHEME 2

hydrogen was introduced into the autoclave at a  $\text{H}_2$  pressure of 0.2 Mpa, the reaction mixture was stirred under 0.2 MPa pressure of hydrogen at  $30^\circ\text{C}$  for 10 hours before releasing hydrogen, and then filtered by suction. The filtrate was concentrated to give 4.02 g (90% yield) of alkene **7** as a white solid, mp  $118\text{--}123^\circ\text{C}$ . IR: 3374, 3055, 2945, 1729, 1597, 1447,  $1147\text{ cm}^{-1}$ ;  $^1\text{H}$  NMR( $\text{CDCl}_3$ ):  $\delta$  7.47 (d, 4H), 7.29–7.14 (m, 10H), 6.40 (d, 1H), 5.60 (m, 1H), 3.63 (s, 3H), 2.95 (d, 2H), 2.52–2.41 (m, 5H), 1.98–1.94 (m, 2H), 1.56 (s, 6H), 1.51–1.46 (m, 4H);  $^{13}\text{C}$  NMR:  $\delta$  177.2, 146.1, 143.0, 136.0, 130.5, 129.3, 128.8, 128.3, 127.0, 126.2, 125.5, 79.5, 58.6, 54.1, 52.3, 46.3, 44.2, 26.6, 26.5, 26.4; MS (EI)  $m/z$ : 498  $[\text{M} + 1]^+$ ; HRMS (EI):  $[\text{M} + 1]^+$  Calcd for  $\text{C}_{33}\text{H}_{39}\text{NO}_3$ : 498.3003. Found: 498.2999. Anal. Calcd for  $\text{C}_{33}\text{H}_{39}\text{NO}_3$ : C, 79.64; H, 7.90; N, 2.82. Found: C, 79.62; H, 7.93; N, 2.84.

## 2.2. Synthesis of 4-[1-Hydroxy-4-[4-(hydroxy-diphenylmethyl)-1-piperidinyl]-butyl]- $\alpha,\alpha$ -dimethyl-benzeneacetic acid methyl ester (**6**)

$\text{Mn}(\text{OAc})_2$  (0.11 g, 0.64 mmol), the Schiff-base ligand **B** [18, 19] (0.20 g, 0.64 mmol), and  $\text{NaBH}_4$  (0.62 g, 16.18 mmol) were added to a stirred solution of compound **7** (4.02 g, 8.09 mmol) in 150 mL of a mixture of toluene and ethanol (1:1). The mixture was stirred at room temperature under oxygen pressure of 0.1 MPa for 4 hours, the progress of reaction was monitored by TLC. The ethanol was removed under reduced pressure and the organic layer was washed with water, dried over  $\text{MgSO}_4$ , and filtered under vacuum.



SCHEME 3

The filtrate was evaporated in vacuo to give crude product **6**, which was chromatographed over silica gel (1:5 hexanes-acetone, v/v) to give pure methyl ester **6** (3.65 g, 90%) as a white powder, mp 157–158°C. IR 3540, 3062, 2931, 2808, 1713, 1445, 1148  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR( $\text{CDCl}_3$ ):  $\delta$  7.51–7.13 (Ar, 14H), 4.59 (d, 1H), 3.15 (d, 1H), 2.98 (d, 1H), 2.50–2.40 (m, 4H), 2.17 (t, 1H), 2.00 (t, 1H), 1.90–1.48 (m, 13H);  $^{13}\text{C}$  NMR:  $\delta$  177.2, 145.9, 144.0, 142.7, 127.9, 126.2, 125.6, 125.5, 79.0, 76.6, 58.6, 54.5, 53.1, 51.9, 46.0, 41.0, 39.5, 26.3, 25.8, 25.7; MS  $m/z$  515( $\text{M}^+$ ). Anal. Calcd for  $\text{C}_{33}\text{H}_{41}\text{NO}_4$ : C, 76.86; H, 8.01; N, 2.72. Found: C, 76.88; H, 8.03; N, 2.69.

## Acknowledgments

The authors thank the National Natural Science Foundation of China (no. 20376016) and the Natural Science Foundation of Zhejiang Province (no. R406378) for financial support of this research.

## References

- [1] A. Graul and J. Castañer, "Fexofenadine hydrochloride," *Drugs of the Future*, vol. 21, no. 10, pp. 1017–1021, 1996.
- [2] A. Markham and A. J. Wagstaff, "Fexofenadine," *Drug*, vol. 55, no. 2, pp. 269–274, 1998.
- [3] K. Simpson and B. Jarvis, "Fexofenadine: a review of its use in the management of seasonal allergic rhinitis and chronic idiopathic urticaria," *Drugs*, vol. 59, no. 2, pp. 301–321, 2000.
- [4] T. Uno, N. Yasui-Furukori, T. Takahata, K. Sugawara, and T. Tateishi, "Liquid chromatographic determination of fexofenadine in human plasma with fluorescence detection," *Journal of Pharmaceutical and Biomedical Analysis*, vol. 35, no. 4, pp. 937–942, 2004.
- [5] R. Tacke, T. Schmid, M. Penka, C. Burschka, W. Bains, and J. Warneck, "Syntheses and pharmacological properties of the histaminic  $\text{H}_1$  antagonists sila-terfenadine-A, sila-terfenadine-B, disila-terfenadine, and sila-fexofenadine: a study on C/Si bioisosterism," *Organometallics*, vol. 23, no. 21, pp. 4915–4923, 2004.
- [6] H. Gross, A. Rieche, E. Höft, and E. Beyer, "Dichloromethyl methyl ether," *Organic Syntheses, Collective*, vol. 5, p. 365, 1973.
- [7] C. H. Senanayake, Q. K. Fang, and S. H. Wilkinson, "Preparation of terfenadine and derivatives," WO patent 9833789, 1998.
- [8] C. H. Senanayake, Q. K. Fang, and S. H. Wilkinson, "Preparation of terfenadine and derivatives," *Chemical Abstracts*, vol. 129, p. 161498r, 1998.
- [9] R. C. H. King and M. A. Kaminski, "4-Diphenylmethyl piperidine derivatives and process for their preparation," WO patent 9321156, 1993.
- [10] R. C. H. King and M. A. Kaminski, "4-(diphenylmethyl) piperidine derivatives and process for their preparation," *Chemical Abstracts*, vol. 120, p. 217287e, 1994.
- [11] C. Schroeder, R. Huddleston, and R. Charles, "Process for the production of the piperidine derivative fexofenadine," WO patent 102,776(1994). A1, 1994.
- [12] C. Schroeder, R. Huddleston, and R. Charles, "Preparation of fexofenadine and related compounds," *Chemical Abstracts*, vol. 138, p. 55873k, 2003.
- [13] S. H. Kawai, R. J. Hambalek, and G. Just, "A facile synthesis of an oxidation product of terfenadine," *Journal of Organic Chemistry*, vol. 59, no. 9, pp. 2620–2622, 1994.
- [14] A. A. Carr, J. E. Dolfini, and G. J. Wright, "Piperidine derivatives," US patent 4254129, 1981.
- [15] A. A. Carr, J. E. Dolfini, and G. J. Wright, "Piperidine derivatives with antihistamine action," *Chemical Abstracts*, vol. 194, p. 156759e, 1981.
- [16] G. Castaldi, G. Barreca, and D. Magron, "Process for preparation of 4-[1-hydroxy-4-(4-(hydroxydiphenylmethyl)-1-piperidinyl)butyl]- $\alpha$ ,  $\alpha$ -dimethylbenzeneacetic acid," US patent 2002/0198233, 2002.
- [17] G. Castaldi, G. Barreca, and D. Magron, "A process for the preparation of 4-[1-hydroxy-4-(4-(hydroxydiphenylmethyl)-1-piperidinyl)-butyl]- $\alpha$ ,  $\alpha$ -dimethylbenzeneacetic acid," *Chemical Abstracts*, vol. 138, p. 4522, 2002.
- [18] N. H. Lee, J. C. Byun, J. S. Baik, C.-H. Han, and S. Han, "Development of Mn(III)(schiff base) complexes for the catalyst of olefin oxygenation to alcohols in the presence of  $\text{NaBH}_4$ ," *Bulletin of the Korean Chemical Society*, vol. 23, no. 10, pp. 1365–1366, 2002.
- [19] N. H. Lee, J. S. Baik, and S. Han, "Development of manganese(III) acetate along with schiff-base ligands as the catalyst for the oxygenation of olefins in the  $\text{O}_2/\text{NaBH}_4$  system," *Bulletin of the Korean Chemical Society*, vol. 25, no. 10, pp. 1455–1456, 2004.