

Research Letter

Three-Component One-Pot Synthesis of Novel Benzo[*b*]1,8-naphthyridines Catalyzed by Bismuth(III) Chloride

**Tangali R. Ravikumar Naik, Halehatty S. Bhojya Naik,
Halehatty R. Prakasha Naik, and P. J. Bindu**

*Department of PG Studies and Research in Industrial Chemistry, School of Chemical Sciences, Kuvempu University,
Shankaraghatta, Karnataka 577451, India*

Correspondence should be addressed to Halehatty S. Bhojya Naik, hsb_naik@rediffmail.com

Received 21 July 2008; Accepted 23 November 2008

Recommended by Pierre Esteves

A novel and efficient three-component one-pot synthesis of benzo[*b*]1,8-naphthyridines by 2-amino-4-methylquinoline, aromatic aldehydes, and malononitrile was done. The reaction was catalyzed by an acidic Bismuth(III) chloride, functionalized Bismuth(III) chloride, at room temperature to give various benzo[*b*]1,8-naphthyridines in high yields. The Bismuth(III) chloride is an environmentally friendly catalyst.

Copyright © 2008 Tangali R. Ravikumar Naik et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction


One of the most important reactions in organic chemistry for the formation of carbon—carbon bonds—is the multicomponent reactions (MCRs) and much effort has been devoted to the development of this reactions in recent years [1, 2].

Among the nitrogen heterocycles, naphthyridines and their derivatives represent an important class of organic molecules that attract the interest of both synthetic and medicinal chemists. Functionalized naphthyridines have found applications as pharmaceuticals, fungicides, bactericides, herbicides, and insecticides as well as useful synthetic blocks in the preparation of several alkaloids [3–6]. Many syntheses of naphthyridines are known, but due to their importance, the development of new synthetic approaches remains an active research area [7, 8].

Various procedures are available for the synthesis of benzo[*b*]1,8-naphthyridines. However, there are several disadvantages associated with these methodologies including unsatisfactory yields, long conversion times and inflammable organic solvents. In recent years, we have initiated our efforts on the development of simple methods for the synthesis of naphthyridines derivatives [9–11]. Hence, now we report herein a useful, general approach towards the

formation of benzo[*b*]1,8-naphthyridines in one-pot operation (Scheme 1). This method permits the condensation of aromatic 2-amino-4-methylquinoline 1, aldehyde 2, and malononitrile 3 using Lewis acid catalysts under mild conditions to afford diverse benzo[*b*]1,8-naphthyridines (4a–h) and is also amenable to small library production using solid-phase techniques.

In addition, there has been an intense interest in the selection of different catalysts in development of new methods for MCRs. The utility of different metal salts such as mercuric salts, lead salts, zinc chloride, cadmium chloride, and indium chloride, lanthanide compounds, InBr_3 , ZrCl_4 [12, 13], as potential catalyst in variety of synthetic reactions was recognized in recent years. However, the expensive-ness and toxicity of some metal salts hampers their wide applications in organic synthesis. On the other hand, in recent years, economically affordable ecofriendly catalysts received some interest in various organic reactions. The application of some of these catalysts such as Cu(II) salts [14], Fe(II)/Fe(III) salts [15], nickel salts, and Bismuth(III) salts [16, 17] as catalysts in organic synthesis, has been investigated extensively. Bismuth trichloride has received considerable attention as an inexpensive, nontoxic, and readily available catalyst for various organic transformations,

SCHEME 1: The synthesis of benzo[*b*]1,8-naphthyridines.

affording the corresponding products in excellent yields with high selectivity (Scheme 1).

Equimolar of various substituted aromatic aldehyde 2 reacted well with 2-amino-4-methylquinoline 1 and malononitrile 3 in presence of 10 mol% of BiCl_3 catalyst in CH_3CN solvent to give the corresponding substituted naphthyridine 4a-h in 93–97% yields. In each experiment, molar ratio 1:1:1.5 of the three components 1, 2, and 3 were used as reactants. The method is very simple and it can be used as derivatives of substituted 2-amino-4-methylquinoline 1, different substituted aromatic aldehyde 2, and malononitrile 3 to prepare different substituted naphthyridine derivatives (Table 1).

2. Experimental

Melting points were recorded on an open capillary tube with a Buchi melting point apparatus and are uncorrected. Elemental analyses were carried out using Perkin-Elmer 24°C CHN-analyzer. IR spectra were recorded on an FT-IR infrared spectrophotometer. ^1H -NMR spectra were obtained using a 300 MHz on a Bruker spectrometer (chemical shifts in δ ppm). Mass spectra were recorded on an LC MS Mass spectrometer.

2.1. General Procedure

A mixture of 2-amino-4-methylquinoline (1 mmol), aromatic aldehyde (1.5 mmol), and malononitriles (1 mmol) in anhydrous CH_3CN (15 mL) was stirred at room temp for 30 minutes. BiCl_3 (20 mol%) was added over a period of 20 minutes. The resulting mixture was stirred for 4–5 hours. After completion of the reaction as indicated by TLC, the reaction mixture was diluted with water (30 mL) and extracted with ethyl acetate (3×15 mL). The organic layer was separated, dried (Na_2SO_4), and concentrated, and the resulting product was recrystallized to afford pure benzo[*b*]1,8-naphthyridines 4a. The same procedure was used for the synthesis of 4b–h. The physicochemical data for the synthesized compounds are as shown below.

4a: M.p. 160–163°C. IR (KBr): 3251; 1723; cm^{-1} . ^1H NMR ($\text{DMSO } d_6$): 1.20 (s, 3H, CH_3), 6.33 (s, 2H, NH_2), 6.80–7.40 (m, 5Harom), 7.45 (d, 1H, $J = 8.5$), 7.51 (d, 1H, $J = 8.5$), 7.70–7.72 (m, 2Harom), 21.5, 89.2, 118.5 (CN), 121.4, 124.5, 126.7, 127.5, 127.6, 127.7, 129.4, 129.5, 129.6, 129.5, 129.4, 138.3, 138.8, 146.8, 152.6, 160.6, 162.5 (NH_2), mass m/z :310.

4b: M.p. 162–165°C. IR (KBr): 3251; 1723; cm^{-1} . ^1H NMR ($\text{DMSO } d_6$): 1.26 (s, 3H, CH_3), 1.80 (s, 3H, CH_3), 6.30 (s, 2H, NH_2), 6.68–7.25 (m, 5Harom), 7.36 (d, 1H, $J = 8.5$), 7.42–7.45 (m, 3Harom), 21.3, 24.9, 89.0, 118.4 (CN), 121.3, 124.2, 126.8, 127.3, 127.6, 127.4, 129.2, 129.5, 129.3, 129.4, 129.7, 138.0, 138.7, 146.7, 152.4, 160.3, 162.4 (NH_2), mass m/z :324.

4c: M.p. 171–173°C. IR (KBr): 3251; 1723; cm^{-1} . ^1H NMR ($\text{DMSO } d_6$): 1.22 (s, 3H, CH_3), 3.10 (s, 3H, OCH_3), 6.25 (s, 2H, NH_2), 6.70–7.13 (m, 5Harom), 7.26 (d, 1H, $J = 8.5$), 7.33–7.40 (m, 3Harom), 20.8, 25.4, 88.9, 118.3 (CN), 121.5, 124.5, 126.7, 127.6, 127.6, 127.8, 129.4, 129.7, 129.8, 129.8, 129.9, 138.3, 138.5, 146.8, 152.5, 160.4, 162.7 (NH_2), mass m/z :340.

4d: M.p. 178–180°C. IR (KBr): 3251; 1723; cm^{-1} . ^1H NMR ($\text{DMSO } d_6$): 1.15 (s, 3H, CH_3), 6.20 (s, 2H, NH_2), 6.80–7.36 (m, 5Harom), 7.40 (d, 1H, $J = 8.5$), 7.58–7.60 (m, 3Harom), 21.7, 89.5, 118.2 (CN), 121.3, 124.4, 126.5, 127.3, 127.6, 127.5, 129.5, 129.6, 129.5, 129.5, 129.7, 132.5(Cl), 138.5, 138.8, 146.8, 152.6, 160.5, 162.7 (NH_2), mass m/z :344.

4e: M.p. 170–173°C. IR (KBr): 3251; 1723; cm^{-1} . ^1H NMR ($\text{DMSO } d_6$): 1.23 (s, 3H, CH_3), 1.65 (s, 3H, CH_3), 1.93 (s, 3H, CH_3), 6.20 (s, 2H, NH_2), 6.70–6.92 (m, 5Harom), 7.08 (d, 1H, $J = 8.5$), 7.20–7.30 (m, 3Harom), 20.5, 23.5, 24.7, 89.4, 118.4 (CN), 121.5, 124.4, 126.4, 127.5, 127.6, 127.7, 129.4, 129.5, 129.6, 129.8, 129.9, 138.5, 138.6, 146.8, 152.6, 160.5, 162.9 (NH_2), mass m/z :338.

4f: M.p. 162–164°C. IR (KBr): 3251; 1723; cm^{-1} . ^1H NMR ($\text{DMSO } d_6$): 1.16 (s, 3H, CH_3), 1.95 (s, 3H, CH_3), 6.20 (s, 2H, NH_2), 6.66–6.90 (m, 5Harom), 6.95 (d, 1H, $J = 8.5$), 6.98–7.20 (m, 3Harom), 21.6, 25.2, 88.7, 118.0 (CN), 121.5, 124.6, 126.7, 127.5, 127.6, 127.6, 129.5, 129.6, 129.6, 129.7, 129.8, 138.5, 138.7, 146.8, 152.6, 160.5, 162.6 (NH_2), mass m/z :324.

TABLE 1: One-pot synthesis of benzo[*b*]1,8-naphthyridines.

Entry	R	R ₁	R ₂	Product ^a	Yield (%) ^b
1	H	H	H	4a	93
2	CH ₃	H	H	4b	95
3	OCH ₃	H	H	4c	95
4	Cl	H	H	4d	97
5	CH ₃	CH ₃	H	4e	93
6	H	H	CH ₃	4f	95
7	H	H	OCH ₃	4g	96
8	H	H	Cl	4h	96

^aAll the products were characterised by IR, ¹H NMR, and mass spectra.^bYields of isolated products.

4g: M.p. 171–173°C. IR (KBr): 3251; 1723; cm⁻¹. ¹H NMR (DMSO d₆): 1.21 (s, 3H, CH₃), 3.00 (s, 3H, OCH₃), 6.20 (s, 2H, NH₂), 6.71–7.04 (m, 5Harom), 7.10 (d, 1H, *J* = 8.5), 7.29–7.35 (m, 3Harom), 21.2, 26.3, 89.3, 117.9 (CN), 121.5, 124.2, 126.6, 127.5, 127.5, 127.6, 129.4, 129.4, 129.5, 129.6, 129.6, 138.7, 138.9, 146.7, 152.6, 160.5, 162.7 (NH₂), mass *m/z*: 340.

4h: M.p. 176–179°C. IR (KBr): 3251; 1723; cm⁻¹. ¹H NMR (DMSO d₆): 1.15 (s, 3H, CH₃), 6.20 (s, 2H, NH₂), 6.75–7.20 (m, 5Harom), 7.32 (d, 1H, *J* = 8.5), 7.35–7.38 (m, 3Harom), 21.7, 89.2, 118.3 (CN), 121.3, 124.2, 126.5, 127.3, 127.4, 127.4, 129.2, 129.3, 129.3, 129.3, 129.3, 131.5(Cl), 138.0, 138.7, 146.7, 152.4, 160.3, 162.2 (NH₂), mass *m/z*: 344.

Elemental Analysis

4a. C₂₀H₁₄N₄ Calc: C, 77.40; H, 4.55; N, 18.05; Found: C, 77.38; H, 4.53; N, 18.03.

4b. C₂₁H₁₆N₄ Calc: C, 77.76; H, 4.97; N, 17.27; Found: C, 77.74; H, 4.96; N, 17.24.

4c. C₂₁H₁₆N₄O Calc: C, 74.10; H, 4.74; N, 16.46; Found: C, 74.11; H, 4.72; N, 16.43.

4d. C₂₀H₁₃ClN₄ Calc: C, 69.67; H, 3.80; N, 16.25; Found: C, 69.65; H, 3.78; N, 16.23.

4e. C₂₂H₁₈N₄ Calc: C, 78.08; H, 5.36; N, 16.56; Found: C, 78.06; H, 5.34; N, 16.53.

4f. C₂₁H₁₆N₄ Calc: C, 77.76; H, 4.97; N, 17.27; Found: C, 77.75; H, 4.95; N, 17.25.

4g. C₂₁H₁₆N₄O Calc: C, 74.10; H, 4.74; N, 16.46; Found: C, 74.07; H, 4.73; N, 16.47.

4h. C₂₀H₁₃ClN₄ Calc: C, 69.67; H, 3.80; N, 16.25; Found: C, 69.68; H, 3.79; N, 16.24.

3. Conclusion

In summary, we have established a new methodology, based on a three-component reaction to obtain new substituted benzo[*b*]1,8-naphthyridines derivatives catalyzed by BiCl₃. The simplicity of this elegant protocol and accessibility of the starting materials allowed us to prepare these new benzo[*b*]1,8-naphthyridines derivatives that should have wide applicability in heterocyclic and medicinal chemistry.

Acknowledgments

The authors would like to thank UGC, New Delhi, for awarding Rajiv Gandhi Research Fellowship and SIFC, IISC, Bangalore for ¹H NMR and Mass Spectral Studies.

References

- [1] Y. K. Chen and P. J. Walsh, "A one-pot multicomponent coupling reaction for the stereocontrolled synthesis of (*Z*)-trisubstituted allylic alcohols," *Journal of the American Chemical Society*, vol. 126, no. 12, pp. 3702–3703, 2004.
- [2] I. Bae, H. Han, and S. Chang, "Highly efficient one-pot synthesis of *N*-sulfonylamidines by Cu-catalyzed three-component coupling of sulfonyl azide, alkyne, and amine," *Journal of the American Chemical Society*, vol. 127, no. 7, pp. 2038–2039, 2005.
- [3] V. P. Litvinov, S. V. Roman, and V. D. Dyachenko, "Naphthyridines. Structure, physicochemical properties and general methods of synthesis," *Russian Chemical Reviews*, vol. 69, no. 3, pp. 201–220, 2000.
- [4] G. B. Barlin and W.-L. Tan, "Potential antimalarials. I. 1,8-naphthyridines," *Australian Journal of Chemistry*, vol. 37, no. 5, pp. 1065–1073, 1984.
- [5] D. Ramesh and B. Sreenivasulu, "Synthesis and antimicrobial activity of 2-aryl-3-(2-methyl-1,8-h naphthyridin-3-yl)-thiazolidin-4-ones," *Indian Journal of Heterocyclic Chemistry*, vol. 15, no. 4, pp. 363–366, 2006.
- [6] B. Bachowska and T. Zujewska, "Chemistry and applications of benzonaphthyridines," *Arkivoc*, vol. 2001, no. 6, pp. 77–84, 2001.
- [7] K. Mogilaiah and J. U. Rani, "Microwave-assisted solvent-free Friedlander synthesis of 1,8-naphthyridines using ammonium acetate as catalyst," *Indian Journal of Chemistry Section B*, vol. 45, no. 4, pp. 1051–1053, 2006.
- [8] K. Mogilaiah, M. Prashanthi, and S. Kavitha, "Lithium chloride as an efficient catalyst for Friedlander synthesis of 1,8-naphthyridines via the use of microwave irradiation and pestle/mortar," *Indian Journal of Chemistry Section B*, vol. 45, no. 1, pp. 302–304, 2006.
- [9] T. R. R. Naik, H. S. B. Naik, M. Raghavedra, and S. G. K. Naik, "Synthesis of thieno[2,3-*b*]benzo[1,8]naphthyridine-2-carboxylic acids under microwave irradiation and interaction with DNA studies," *Arkivoc*, vol. 2006, no. 15, pp. 84–94, 2006.
- [10] T. R. R. Naik, H. S. B. Naik, and S. R. G. K. Naik, "One pot solvent-free synthesis of 2*H*-pyrano, 2*H*-thiopyrano, 2*H*-selenopyrano[2,3-*b*]-1,8-naphthyridin-2-ones on solid phase catalyst under microwave irradiation," *Journal of Sulfur Chemistry*, vol. 28, no. 4, pp. 393–400, 2007.
- [11] R. N. Butler, "Comparative reactions of nitrogen compounds with the isoelectronic series mercury(II), thallium(III), and lead(IV) acetates. Principles of oxidation reactions," *Chemical Reviews*, vol. 84, no. 3, pp. 249–276, 1984.
- [12] Q. Sun, Y. Yi, Z. Ge, T. Cheng, and R. Li, "A highly efficient solvent-free synthesis of dihydropyrimidinones catalyzed by zinc chloride," *Synthesis*, no. 7, pp. 1047–1051, 2004.
- [13] A. Manjula, B. Vittal Rao, and P. Neelakantan, "An inexpensive protocol for biginelli reaction," *Synthetic Communications*, vol. 34, no. 14, pp. 2665–2671, 2004.
- [14] J. S. Yadav, B. V. S. Reddy, P. Sridhar, et al., "Green protocol for the Biginelli three-component reaction: Ag₃PW₁₂O₄₀ as a novel, water-tolerant heteropolyacid for the synthesis

- of 3,4-dihydropyrimidinones,” *European Journal of Organic Chemistry*, vol. 2004, no. 3, pp. 552–557, 2004.
- [15] I. Mohammadpoor-Baltork, A. R. Khosropour, and H. Aliyan, “Efficient conversion of epoxides to 1,3-dioxolanes catalyzed by bismuth(III) salts,” *Synthetic Communications*, vol. 31, no. 22, pp. 3411–3416, 2001.
- [16] H. Firouzabadi, I. Mohammadpoor-Baltork, and S. Kolagar, “A rapid, selective, and efficient method for deprotection of silyl ethers catalyzed by bismuth(III) salts,” *Synthetic Communications*, vol. 31, no. 6, pp. 905–909, 2001.
- [17] S. H. Mashraqui and M. A. Karnik, “Bismuth nitrate pentahydrate: a convenient reagent for the oxidation of Hantzsch 1,4-dihydropyridines,” *Synthesis*, no. 5, pp. 713–714, 1998.