

THE EULER CIRCLE COMPLEX ANALYSIS SAMPLE PROBLEM

SIDDHARTH KOTHARI

This document contains my solutions to select The Euler Circle Complex Analysis Problems. The first one is same one as in the Poster.

1. SAMPLE PROBLEM 1: JENSEN'S FORMULA

Question 1.1. *Prove the following version of Jensen's Formula: Let f be an entire function that is not identically zero, and let $\nu(t)$ be the number of zeros of f with modulus $< t$, counted with multiplicity. Then $\log |f(0)| = -\int_0^r \frac{\nu(t)}{t} dt + \frac{1}{2\pi} \int_0^{2\pi} \log |f(re^{i\theta})| d\theta$.¹*

Proof. Let D denote the set of zeros of f . Since f is entire and not identically zero, f has only finitely many zeros in $\overline{B_r(0)}$. Indeed, if $\#(D \cap \overline{B_r(0)}) = \infty$, then the Bolzano-Weierstrass theorem and the fact that $\overline{B_r(0)}$ is closed would imply that D would have a limit point in $\overline{B_r(0)}$. By the identity theorem, f would then be zero on $\overline{B_r(0)}$, a contradiction.

Let the zeros be a_1, \dots, a_n (counted with multiplicity). Arrange the elements of the set $\{|a_j| : 1 \leq j \leq n\}$ as $r_1 < \dots < r_k$ where $k \leq n$. For notational convenience, we define $r_{k+1} = r$. Further, let m_j denote the number of zeros of f with modulus r_j . By Jensen's formula (as applied to f on $\overline{B_r(0)}$), it suffices to show that:

$$\sum_{j=1}^n \log \frac{\rho}{|a_j|} = \int_0^r \frac{\nu(t)}{t} dt.$$

First, note that

$$\int_0^r \frac{\nu(t)}{t} dt = \sum_{j=1}^k \int_{r_j}^{r_{j+1}} \frac{\nu(t)}{t} dt = \sum_{j=1}^k M_j \int_{r_j}^{r_{j+1}} \frac{1}{t} dt = \sum_{j=1}^k M_j \log \frac{r_{j+1}}{r_j} = \log \left(\prod_{j=1}^k \left(\frac{r_{j+1}}{r_j} \right)^{M_j} \right),$$

where $M_j = \sum_{a=1}^j m_a$. This is because for $r_j < t < r_{j+1}$, $\nu(t) = M_j$ by the definition of ν . Further,

$$\prod_{j=1}^k \left(\frac{r_{j+1}}{r_j} \right)^{M_j} = \left(\frac{r_2}{r_1} \right)^{m_1} \left(\frac{r_3}{r_2} \right)^{m_1+m_2} \left(\frac{r_4}{r_3} \right)^{m_1+m_2+m_3} \dots \left(\frac{r}{r_k} \right)^{m_1+\dots+m_k} = \prod_{j=1}^k \left(\frac{r}{r_j} \right)^{m_j}.$$

Therefore,

$$\int_0^r \frac{\nu(t)}{t} dt = \sum_{j=1}^k m_j \log \frac{r}{r_j} = \sum_{j=1}^n \log \frac{r}{|a_j|},$$

as desired. ■

¹Course Description: <https://eulercircle.com/classes/complex-analysis/>. Complex Analysis was spread pover two quarters: Winter and Spring 2025.

2. SAMPLE PROBLEM 2: THE CAUCHY INTEGRAL FORMULA

Question 2.1. *Prove the differential version of the Cauchy integral formula.*

Proof. By the induction hypothesis, we have:

$$\frac{f^{(n-1)}(z_0 + h) - f^{(n-1)}(z_0)}{h} = \frac{(n-1)!}{2\pi i h} \left(\int_{\Gamma} \frac{f(z)}{(z - z_0 - h)^n} dz - \int_{\Gamma} \frac{f(z)}{(z - z_0)^n} dz \right).$$

Simplifying,

$$\begin{aligned} \int_{\Gamma} \frac{f(z)}{h} \left(\frac{1}{(z - z_0 - h)^n} - \frac{1}{(z - z_0)^n} \right) dz &= \int_{\Gamma} \frac{f(z)}{h} \left(\frac{(z - z_0)^n - (z - z_0 - h)^n}{(z - z_0 - h)^n (z - z_0)^n} \right) dz \\ &= \int_{\Gamma} \frac{(z - z_0)^n - \sum_{i=0}^n \binom{n}{i} (z - z_0)^i (-h)^{n-i}}{(z - z_0 - h)^n (z - z_0)^n} \frac{f(z)}{h} dz \\ &= \int_{\Gamma} \frac{\sum_{i=0}^{n-1} \binom{n}{i} (z - z_0)^i h^{n-i-1}}{(z - z_0 - h)^n (z - z_0)^n} f(z) dz. \end{aligned}$$

Therefore, we are left with

$$f^{(n)}(z_0) = \int_{\Gamma} \frac{n}{(z - z_0 - h)^n (z - z_0)} f(z) dz + \int_{\Gamma} \frac{\sum_{i=0}^{n-2} \binom{n}{i} (z - z_0)^i h^{n-i-1}}{(z - z_0 - h)^n (z - z_0)^n} f(z) dz.$$

Let M be the maximum value of $|f(z)|$ over Γ . Then,

$$\left| \frac{\sum_{i=0}^{n-2} \binom{n}{i} (z - z_0)^i h^{n-i-1}}{(z - z_0 - h)^n (z - z_0)^n} f(z) \right| \leq \frac{M f(h)}{\rho^n (\rho - h)^n},$$

where

$$f(h) := \sup \left\{ \left| \sum_{i=0}^{n-2} \binom{n}{i} (z - z_0)^i h^{n-i-1} \right| : z \in \Gamma \right\}.$$

Clearly $f(h)$ goes to zero as h tends to 0, so the ML inequality tells us that

$$\lim_{h \rightarrow 0} \left| \int_{\Gamma} \frac{\sum_{i=0}^{n-2} \binom{n}{i} (z - z_0)^i h^{n-i-1}}{(z - z_0 - h)^n (z - z_0)^n} f(z) dz \right| = 0 \text{ as } \lim_{h \rightarrow 0} \frac{2\pi M f(h)}{\rho^{n-1} (\rho - h)^n} = 0.$$

Next, we deal with the first integral. First, note the decomposition

$$\frac{1}{(z - z_0 - h)^n (z - z_0)} = \frac{1}{(z - z_0)^{n+1}} + \frac{(z - z_0)^n - (z - z_0 - h)^n}{(z - z_0)^{n+1} (z - z_0 - h)^n} = \frac{1}{(z - z_0)^{n+1}} + \frac{\sum_{i=0}^{n-1} \binom{n}{i} (z - z_0)^i h^{n-i}}{(z - z_0)^n (z - z_0 - h)^n}.$$

Hence,

$$\int_{\Gamma} \frac{n}{(z - z_0 - h)^n (z - z_0)} f(z) dz = \int_{\Gamma} \frac{n}{(z - z_0)^{n+1}} f(z) dz + \int_{\Gamma} \frac{\sum_{i=0}^{n-1} \binom{n}{i} (z - z_0)^i h^{n-i}}{(z - z_0)^n (z - z_0 - h)^n} n f(z) dz.$$

Invoking the ML inequality again for the second integral

$$\left| \int_{\Gamma} \frac{\sum_{i=0}^{n-1} \binom{n}{i} (z - z_0)^i h^{n-i}}{(z - z_0)^n (z - z_0 - h)^n} n f(z) dz \right| \leq \frac{M n g(h)}{\rho^n (\rho - h)^n} \times 2\pi \rho = \frac{2\pi M n g(h)}{\rho^{n-1} (\rho - h)^n},$$

where

$$g(h) := \max \left| \sum_{i=0}^{n-1} \binom{n}{i} (z - z_0)^i h^{n-i} \right|.$$

Since $\lim_{h \rightarrow 0} g(h) = 0 \implies \lim_{h \rightarrow 0} \frac{2\pi M n g(h)}{\rho^{n-1}(\rho-h)^n} = 0$. Therefore, the second integral vanishes in the limit as $h \rightarrow 0$, and so

$$\lim_{h \rightarrow 0} \frac{f^{(n-1)}(z_0 + h) - f^{(n-1)}(z_0)}{h} = \lim_{h \rightarrow 0} \frac{n!}{2\pi i} \int_{\Gamma} \frac{f(z)}{(z - z_0)^{n+1}} dz = \frac{n!}{2\pi i} \int_{\Gamma} \frac{f(z)}{(z - z_0)^{n+1}} dz$$

as desired. ■

3. SAMPLE PROBLEM 3: CONVERGENCE OF DIRICHLET SERIES

Question 3.1. Given a Dirichlet series $\sum_{n=1}^{\infty} \frac{a_n}{n^s}$, let

$$\sigma_c = \left\{ \sigma : \sum_{n=1}^{\infty} \frac{a_n}{n^s} \text{ converges for some } s \text{ with } \Re s = \sigma \right\}.$$

Prove that if $\Re s > \sigma_c$, then $\sum_{n=1}^{\infty} \frac{a_n}{n^s}$ converges. We call σ_c the abscissa of convergence.

Proof. Define $S_c = \{\sigma : \exists s \in \mathbb{C} : \sum_{n=1}^{\infty} a_n n^{-s} \text{ converges and } \Re(s) = \sigma\}$; note that $\sigma_c = \inf(S)$. Let $s \in \mathbb{C}$ be such that $\Re(s) > \sigma_c$. Then there exists a $\delta \in \mathbb{C}$ such that $\Re(\delta) \in S$ and $\Re(s) > \Re(\delta)$. We must show that for all $\varepsilon > 0$, there exists a $N \in \mathbb{Z}^+$ such that for all $M_1 > M_2 \geq N$, we have

$$\left| \sum_{N=M_1}^{M_2} a_N N^{-s} \right| = \left| \sum_{N=M_1}^{M_2} a_N N^{-\delta} \cdot N^{-(s-\delta)} \right| < \varepsilon.$$

Taking cue from this, define $x_n = a_n n^{-\delta}$ and $y_n = n^{-(s-\delta)}$. Abel's summation lemma states that:

$$\sum_{j=M_1}^{M_2} x_j y_j = (S_{M_2} y_{M_2} - S_{M_1-1} y_{M_1}) + \sum_{j=M_1}^{M_2-1} S_j (y_j - y_{j+1}),$$

where $S_j = \sum_{\mu=1}^j x_{\mu} = \sum_{\mu=1}^j a_{\mu} \mu^{-\delta}$. We take the absolute value of both sides and apply the triangle inequality to the *RHS* to get:

$$\left| \sum_{j=M_1}^{M_2-1} x_j y_j \right| \leq |S_{M_2} y_{M_2}| + |S_{M_1-1} y_{M_1}| + \left| \sum_{j=M_1}^{M_2-1} S_j (y_j - y_{j+1}) \right|.$$

Note that $(S_n)_{n=1}^{\infty}$ is a Cauchy sequence, so it is bounded. Thus, there exists a $C \in \mathbb{R}^+$ such that $|S_n| < C$ for all n . Hence,

$$L = \left| \sum_{j=M_1}^{M_2-1} S_j (y_j - y_{j+1}) \right| \leq \sum_{j=M_1}^{M_2-1} |S_j| |y_j - y_{j+1}| \leq C \sum_{j=M_1}^{M_2-1} |y_{j+1} - y_j|.$$

The last term hints at the mean value theorem. Indeed, define $f_j : [j, j+1]$ by $f_j(x) = x^{-(s-\delta)}$ for all $x \in [j, j+1]$. Then, by the mean value theorem, we have

$$|f(j+1) - f(j)| = |y_{j+1} - y_j| \leq |f'(\xi_j)| \leq |s - \delta| j^{-(\Re(s-\delta)+1)},$$

for some $j \leq \xi_j \leq j+1$. Thus, $L \leq C' \sum_{j=M_1}^{M_2-1} j^{-(\Re(s-\delta)+1)}$ where C' is a constant that depends only on s and δ . By the p -series test, we know that the series $\sum_{n=1}^{\infty} n^{-(\Re(s-\delta)+1)}$ converges as $\Re(s-\delta) > 0$. Therefore, for every $\varepsilon > 0$, there exists a N such that for all $M_2 > M_1 \geq N$, $\left| \sum_{j=M_1}^{M_2-1} j^{-(\Re(s-\delta)+1)} \right| < \varepsilon$ and so $L \leq C' \varepsilon$.

Next, note that

$$|S_{M_2} y_{M_2}| + |S_{M_1-1} y_{M_1}| \leq C(|y_{M_2}| + |y_{M_1}|) = C \left(M_1^{-\Re(s-\delta)} + M_2^{-\Re(s-\delta)} \right) \leq C,$$

where the last inequality follows as the function $h(x) := x^{-\Re(s-\delta)}$ is monotonically decreasing. In conclusion, we have $\left| \sum_{N=M_1}^{M_2} x_j y_j \right| < K\varepsilon$ for all $M_1 > M_2 \geq N$ where K is a constant that is dependent only on s and δ , which completes the proof. ■