

Modular Forms: E_2 and the Modular Discriminant

Siddharth Kothari
spacersid1331@gmail.com

Euler Circle

2 June 2025

Modular Forms: The Definition.

Definition

Let $k \in \mathbb{Z}$. A **modular form** of weight k is a function $f : \mathbb{H} \rightarrow \mathbb{C}$ such that

Modular Forms: The Definition.

Definition

Let $k \in \mathbb{Z}$. A **modular form** of weight k is a function $f : \mathbb{H} \rightarrow \mathbb{C}$ such that

- f is holomorphic on \mathbb{H} ,

Modular Forms: The Definition.

Definition

Let $k \in \mathbb{Z}$. A **modular form** of weight k is a function $f : \mathbb{H} \rightarrow \mathbb{C}$ such that

- f is holomorphic on \mathbb{H} ,
- $f\left(\frac{a\tau+b}{c\tau+d}\right) = (c\tau+d)^k f(\tau)$ for all $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ and $\tau \in \mathbb{H}$,

Modular Forms: The Definition.

Definition

Let $k \in \mathbb{Z}$. A **modular form** of weight k is a function $f : \mathbb{H} \rightarrow \mathbb{C}$ such that

- f is holomorphic on \mathbb{H} ,
- $f\left(\frac{a\tau+b}{c\tau+d}\right) = (c\tau+d)^k f(\tau)$ for all $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ and $\tau \in \mathbb{H}$,
- $|f(\tau)|$ stays bounded as $\Im(\tau) \rightarrow \infty$.

Modular Forms: The Definition.

Definition

Let $k \in \mathbb{Z}$. A **modular form** of weight k is a function $f : \mathbb{H} \rightarrow \mathbb{C}$ such that

- f is holomorphic on \mathbb{H} ,
- $f\left(\frac{a\tau+b}{c\tau+d}\right) = (c\tau+d)^k f(\tau)$ for all $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ and $\tau \in \mathbb{H}$,
- $|f(\tau)|$ stays bounded as $\Im(\tau) \rightarrow \infty$.

What is the purpose of the so-called **modularity condition**? Why does $\mathrm{SL}_2(\mathbb{Z})$ show up? We'll take a step back and consider functions defined on lattices.

Homogeneous Functions on Lattices: Definition.

Recall the definition of a lattice, which generalizes the relationship of \mathbb{Z} as a subset of \mathbb{R} .

Definition

Let ω_1 and ω_2 be two complex numbers that are linearly independent over \mathbb{R} . The **lattice** generated by ω_1 and ω_2 , denoted by $\Lambda(\omega_1, \omega_2)$, is defined by the set $\{n_1\omega_1 + n_2\omega_2 : n_i \in \mathbb{Z}\}$.

Remark

Notice that when we say ω_1 and ω_2 are *linearly independent* over \mathbb{R} , we mean $\omega_1/\omega_2 \notin \mathbb{R}$.

Homogeneous Functions on Lattices: Definition.

Recall the definition of a lattice, which generalizes the relationship of \mathbb{Z} as a subset of \mathbb{R} .

Definition

Let ω_1 and ω_2 be two complex numbers that are linearly independent over \mathbb{R} . The **lattice** generated by ω_1 and ω_2 , denoted by $\Lambda(\omega_1, \omega_2)$, is defined by the set $\{n_1\omega_1 + n_2\omega_2 : n_i \in \mathbb{Z}\}$.

Remark

Notice that when we say ω_1 and ω_2 are *linearly independent* over \mathbb{R} , we mean $\omega_1/\omega_2 \notin \mathbb{R}$.

Now, we consider a special class of functions $F : \mathcal{L} \rightarrow \mathbb{C}$, where \mathcal{L} is the set of all lattices. Recall that we may scale lattices by a constant: $a\Lambda := \{a\lambda : \lambda \in \Lambda\}$.

Homogeneous Functions on Lattices: Definition.

Recall the definition of a lattice, which generalizes the relationship of \mathbb{Z} as a subset of \mathbb{R} .

Definition

Let ω_1 and ω_2 be two complex numbers that are linearly independent over \mathbb{R} . The **lattice** generated by ω_1 and ω_2 , denoted by $\Lambda(\omega_1, \omega_2)$, is defined by the set $\{n_1\omega_1 + n_2\omega_2 : n_i \in \mathbb{Z}\}$.

Remark

Notice that when we say ω_1 and ω_2 are *linearly independent* over \mathbb{R} , we mean $\omega_1/\omega_2 \notin \mathbb{R}$.

Now, we consider a special class of functions $F : \mathcal{L} \rightarrow \mathbb{C}$, where \mathcal{L} is the set of all lattices. Recall that we may scale lattices by a constant: $a\Lambda := \{a\lambda : \lambda \in \Lambda\}$.

Definition

Let $k \in \mathbb{Z}$. A function $F : \mathcal{L} \rightarrow \mathbb{C}$ is said to be homogeneous of weight k if

$$F(\alpha\Lambda) = \alpha^{-k}F(\Lambda),$$

for all $\Lambda \in \mathcal{L}$ and $\alpha \in \mathbb{C}$.

Homogeneous Functions on Lattices: Unpacking Homogeneity.

Working with functions defined on sets is a bit clunky, so instead given such a F , we define G_F to be $G_F(\omega_1, \omega_2) := F(\Lambda(\omega_1, \omega_2))$ for all pairs of linearly independent complex numbers (ω_1, ω_2) .

Homogeneous Functions on Lattices: Unpacking Homogeneity.

Working with functions defined on sets is a bit clunky, so instead given such a F , we define G_F to be $G_F(\omega_1, \omega_2) := F(\Lambda(\omega_1, \omega_2))$ for all pairs of linearly independent complex numbers (ω_1, ω_2) .

Remark

The homogeneity of F implies that

$$G(\alpha\omega_1, \alpha\omega_2) = F(\Lambda(\alpha\omega_1, \alpha\omega_2)) = F(\alpha\Lambda(\omega_1, \omega_2)) = \alpha^k G(\omega_1, \omega_2).$$

Taking α to be ω_1^{-1} , we have $G(1, \omega_2/\omega_1) = \omega_1^k G(\omega_1, \omega_2) = F(\Lambda(1, \omega_2/\omega_1))$, so G is completely determined by where F sends lattices generated by $\{1, \tau\}$ for some $\tau \in \mathbb{C}$.

Homogeneous Functions on Lattices: Unpacking Homogeneity.

Working with functions defined on sets is a bit clunky, so instead given such a F , we define G_F to be $G_F(\omega_1, \omega_2) := F(\Lambda(\omega_1, \omega_2))$ for all pairs of linearly independent complex numbers (ω_1, ω_2) .

Remark

The homogeneity of F implies that

$$G(\alpha\omega_1, \alpha\omega_2) = F(\Lambda(\alpha\omega_1, \alpha\omega_2)) = F(\alpha\Lambda(\omega_1, \omega_2)) = \alpha^k G(\omega_1, \omega_2).$$

Taking α to be ω_1^{-1} , we have $G(1, \omega_2/\omega_1) = \omega_1^k G(\omega_1, \omega_2) = F(\Lambda(1, \omega_2/\omega_1))$, so G is completely determined by where F sends lattices generated by $\{1, \tau\}$ for some $\tau \in \mathbb{C}$.

This allows us to consider a uni-variate function $f(\tau) := G(1, \tau) = F(\Lambda(1, \tau))$. However, there are constraints on f : given two complex numbers $\tau_1 \neq \tau_2$, we may have $\Lambda(1, \tau_1) = \Lambda(1, \tau_2)$, meaning we have $f(\tau_1) = f(\tau_2)$. And when does this happen?

Homogeneous Functions on Lattices: Unpacking Homogeneity.

Working with functions defined on sets is a bit clunky, so instead given such a F , we define G_F to be $G_F(\omega_1, \omega_2) := F(\Lambda(\omega_1, \omega_2))$ for all pairs of linearly independent complex numbers (ω_1, ω_2) .

Remark

The homogeneity of F implies that

$$G(\alpha\omega_1, \alpha\omega_2) = F(\Lambda(\alpha\omega_1, \alpha\omega_2)) = F(\alpha\Lambda(\omega_1, \omega_2)) = \alpha^k G(\omega_1, \omega_2).$$

Taking α to be ω_1^{-1} , we have $G(1, \omega_2/\omega_1) = \omega_1^k G(\omega_1, \omega_2) = F(\Lambda(1, \omega_2/\omega_1))$, so G is completely determined by where F sends lattices generated by $\{1, \tau\}$ for some $\tau \in \mathbb{C}$.

This allows us to consider a uni-variate function $f(\tau) := G(1, \tau) = F(\Lambda(1, \tau))$. However, there are constraints on f : given two complex numbers $\tau_1 \neq \tau_2$, we may have $\Lambda(1, \tau_1) = \Lambda(1, \tau_2)$, meaning we have $f(\tau_1) = f(\tau_2)$. And when does this happen?

Proposition

Two pairs $(1, \tau_1)$ and $(1, \tau_2)$ are such that $\Lambda(1, \tau_1) = \Lambda(1, \tau_2)$ if and only if there exists integers a, b, c and d such that $ad - bc = 1$ and

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \tau_1 \\ 1 \end{pmatrix} = \begin{pmatrix} \tau_2 \\ 1 \end{pmatrix}.$$

Homogeneous Functions on Lattices: Unpacking Homogeneity.

With this in mind, we have

Homogeneous Functions on Lattices: Unpacking Homogeneity.

With this in mind, we have

$$\begin{aligned} f(\tau_1) &= F(\Lambda(1, \tau_1)) = F(\Lambda(c\tau_1 + d, a\tau_1 + b)) \\ &= F\left((c\tau_1 + d)\Lambda\left(1, \frac{a\tau_1 + b}{c\tau_1 + d}\right)\right) \\ &= (c\tau_1 + d)^k F\left(\Lambda\left(1, \frac{a\tau_1 + b}{c\tau_1 + d}\right)\right) \\ &= (c\tau_1 + d)^k f\left(\frac{a\tau_1 + b}{c\tau_1 + d}\right), \end{aligned}$$

Homogeneous Functions on Lattices: Unpacking Homogeneity.

With this in mind, we have

$$\begin{aligned} f(\tau_1) &= F(\Lambda(1, \tau_1)) = F(\Lambda(c\tau_1 + d, a\tau_1 + b)) \\ &= F\left((c\tau_1 + d)\Lambda\left(1, \frac{a\tau_1 + b}{c\tau_1 + d}\right)\right) \\ &= (c\tau_1 + d)^k F\left(\Lambda\left(1, \frac{a\tau_1 + b}{c\tau_1 + d}\right)\right) \\ &= (c\tau_1 + d)^k f\left(\frac{a\tau_1 + b}{c\tau_1 + d}\right), \end{aligned}$$

meaning that f satisfies the modularity condition!

Creating Modular Forms Using the Weierstrass \wp Function.

Definition

Let ω_1 and ω_2 be two complex numbers linearly independent over \mathbb{R} . Then, the Weierstrass elliptic function corresponding to the lattice $\Lambda(\omega_1, \omega_2)$ generated by ω_1 and ω_2 , denoted by $\wp_{\Lambda(\omega_1, \omega_2)}$, is defined by

$$\wp_{\Lambda(\omega_1, \omega_2)}(z) = \frac{1}{z^2} + \sum_{\lambda \in \Lambda(\omega_1, \omega_2) \setminus \{0\}} \left(\frac{1}{(z + \lambda)^2} - \frac{1}{\lambda^2} \right),$$

for all $z \in \mathbb{C} - \Lambda(\omega_1, \omega_2)$.

Creating Modular Forms Using the Weierstrass \wp Function.

Definition

Let ω_1 and ω_2 be two complex numbers linearly independent over \mathbb{R} . Then, the Weierstrass elliptic function corresponding to the lattice $\Lambda(\omega_1, \omega_2)$ generated by ω_1 and ω_2 , denoted by $\wp_{\Lambda(\omega_1, \omega_2)}$, is defined by

$$\wp_{\Lambda(\omega_1, \omega_2)}(z) = \frac{1}{z^2} + \sum_{\lambda \in \Lambda(\omega_1, \omega_2) \setminus \{0\}} \left(\frac{1}{(z + \lambda)^2} - \frac{1}{\lambda^2} \right),$$

for all $z \in \mathbb{C} - \Lambda(\omega_1, \omega_2)$.

Remark

Out of the many relationships satisfied by the Weierstrass elliptic function, one of them is that $\wp_{\Lambda(\alpha\omega_1, \alpha\omega_2)}(\alpha z) = \alpha^{-2} \wp_{\Lambda(\omega_1, \omega_2)}(z)$ for all $\alpha \in \mathbb{C}^*$.

Creating Modular Forms Using the Weierstrass \wp Function.

Definition

Let ω_1 and ω_2 be two complex numbers linearly independent over \mathbb{R} . Then, the Weierstrass elliptic function corresponding to the lattice $\Lambda(\omega_1, \omega_2)$ generated by ω_1 and ω_2 , denoted by $\wp_{\Lambda(\omega_1, \omega_2)}$, is defined by

$$\wp_{\Lambda(\omega_1, \omega_2)}(z) = \frac{1}{z^2} + \sum_{\lambda \in \Lambda(\omega_1, \omega_2) \setminus \{0\}} \left(\frac{1}{(z + \lambda)^2} - \frac{1}{\lambda^2} \right),$$

for all $z \in \mathbb{C} - \Lambda(\omega_1, \omega_2)$.

Remark

Out of the many relationships satisfied by the Weierstrass elliptic function, one of them is that $\wp_{\Lambda(\alpha\omega_1, \alpha\omega_2)}(\alpha z) = \alpha^{-2} \wp_{\Lambda(\omega_1, \omega_2)}(z)$ for all $\alpha \in \mathbb{C}^*$.

Recall that the Laurent series expansion for $\wp_{\Lambda(\omega_1, \omega_2)}$ about $z = 0$ (where it has a double pole, as at every other lattice point), has the form

$$\frac{1}{z^2} + \sum_{k=1}^{\infty} c_{2k}(\omega_1, \omega_2) z^{2k},$$

where c_{2k} is a complex-valued function that takes as input two linearly independent complex numbers for each $k \in \mathbb{N}$.

Creating Modular Forms Using the Weierstrass \wp Function.

Putting the previous relationship into the Laurent series, we have

Creating Modular Forms Using the Weierstrass \wp Function.

Putting the previous relationship into the Laurent series, we have

$$\begin{aligned}\wp_{\Lambda(\alpha\omega_1, \alpha\omega_2)}(\alpha z) &= \frac{1}{(\alpha z)^2} + \sum_{k=1}^{\infty} c_{2k}(\alpha\omega_1, \alpha\omega_2)(\alpha z)^{2k} \\ &= \alpha^{-2} \left(\frac{1}{z^2} + \sum_{k=1}^{\infty} \alpha^{2k+2} c_{2k}(\alpha\omega_1, \alpha\omega_2) z^{2k} \right) \\ &= \alpha^{-2} \left(\frac{1}{z^2} + \sum_{k=1}^{\infty} c_{2k}(\omega_1, \omega_2) z^{2k} \right),\end{aligned}$$

Creating Modular Forms Using the Weierstrass \wp Function.

Putting the previous relationship into the Laurent series, we have

$$\begin{aligned}\wp_{\Lambda(\alpha\omega_1, \alpha\omega_2)}(\alpha z) &= \frac{1}{(\alpha z)^2} + \sum_{k=1}^{\infty} c_{2k}(\alpha\omega_1, \alpha\omega_2)(\alpha z)^{2k} \\ &= \alpha^{-2} \left(\frac{1}{z^2} + \sum_{k=1}^{\infty} \alpha^{2k+2} c_{2k}(\alpha\omega_1, \alpha\omega_2) z^{2k} \right) \\ &= \alpha^{-2} \left(\frac{1}{z^2} + \sum_{k=1}^{\infty} c_{2k}(\omega_1, \omega_2) z^{2k} \right),\end{aligned}$$

meaning that $c_{2k+2}(\alpha\omega_1, \alpha\omega_2) = \alpha^{-(2k+2)} c_{2k}(\omega_1, \omega_2)$ for all $\alpha \in \mathbb{C}^*$.

Creating Modular Forms Using the Weierstrass \wp Function.

Putting the previous relationship into the Laurent series, we have

$$\begin{aligned}\wp_{\Lambda(\alpha\omega_1, \alpha\omega_2)}(\alpha z) &= \frac{1}{(\alpha z)^2} + \sum_{k=1}^{\infty} c_{2k}(\alpha\omega_1, \alpha\omega_2)(\alpha z)^{2k} \\ &= \alpha^{-2} \left(\frac{1}{z^2} + \sum_{k=1}^{\infty} \alpha^{2k+2} c_{2k}(\alpha\omega_1, \alpha\omega_2) z^{2k} \right) \\ &= \alpha^{-2} \left(\frac{1}{z^2} + \sum_{k=1}^{\infty} c_{2k}(\omega_1, \omega_2) z^{2k} \right),\end{aligned}$$

meaning that $c_{2k+2}(\alpha\omega_1, \alpha\omega_2) = \alpha^{-(2k+2)} c_{2k}(\omega_1, \omega_2)$ for all $\alpha \in \mathbb{C}^*$.

Remark

We have that c_{2k} is a homogeneous function on lattices of weight $2k + 2$. From our previous work, we know that the function defined by $f_{2k}(\tau) = c_{2k}(1, \tau)$ for all $\tau \in \mathbb{H}$ is a modular form of weight $2k + 2$!

Eisenstein Series as a Modular Form: Definition.

Computing these coefficients, we get the following.

Proposition

Using the notation above, we have that $c_{2k}(\omega_1, \omega_2) = (2k+1)E_{2k+2}(\omega_1, \omega_2)$, where

$$E_n(\omega_1, \omega_2) := \sum_{\lambda \in \Lambda(\omega_1, \omega_2) - \{0\}} \frac{1}{\lambda^k},$$

for all $n \geq 3$.

Due to time constraints, we will not prove this here.

Definition

Let $k \geq 4$ be even. Define E_k , called the Eisenstein series of weight k , as a complex valued function defined on the upper half plane

$$E_k(\tau) := \sum_{\omega \in \Lambda(1, \tau) - \{0\}} \frac{1}{\omega^k},$$

for all $\tau \in \mathbb{H}$.

Eisenstein Series as a Modular Form: Convergence.

First, we tackle the definition of G_k .

Proposition

For $k \geq 4$, the function G_k is absolutely convergent.

Eisenstein Series as a Modular Form: Convergence.

First, we tackle the definition of G_k .

Proposition

For $k \geq 4$, the function G_k is absolutely convergent.

To prove this, we need a lemma.

Lemma

For each $\tau \in \mathbb{H}$, there is a $\delta_\tau \in (0, 1)$ such that

$$|m\tau + n| \geq \delta_\tau |mi + n|,$$

for all integers m and n .

Eisenstein Series as a Modular Form: Convergence.

Proof.

We may assume that $m \neq 0$, for if $m = 0$, then one can take δ_τ to be any element of $(0, 1)$ for all τ . Dividing both sides by $|m|$ and then $|i + n/m|$, our problem is now equivalent to finding a δ_τ such that

$$\left| \frac{\tau + n/m}{i + n/m} \right| \geq \delta_\tau,$$

for all integers n and $m \neq 0$.

Eisenstein Series as a Modular Form: Convergence.

Proof.

We may assume that $m \neq 0$, for if $m = 0$, then one can take δ_τ to be any element of $(0, 1)$ for all τ . Dividing both sides by $|m|$ and then $|i + n/m|$, our problem is now equivalent to finding a δ_τ such that

$$\left| \frac{\tau + n/m}{i + n/m} \right| \geq \delta_\tau,$$

for all integers n and $m \neq 0$. Taking cue from the above expression, define the real valued function $f_\tau(x) = \left| \frac{\tau - x}{i - x} \right|$ for all real numbers x . Note that f is continuous at all $x \in \mathbb{R}$, and as $x \rightarrow \pm\infty$, we have $f_\tau(x) \rightarrow 1$. Moreover, f_τ maps to the positive real numbers. This implies that there exists a $R_\tau > 0$ such that for $|x| > R_\tau$, we have $f_\tau(x) \geq 1/2$. On the other hand, as the image of a compact set under a continuous function is compact, there exists a $c(\tau) > 0$ such that $f_\tau(x) \geq c(\tau)$ for all $x \in [-R_\tau, R_\tau]$. Therefore, for all $x \in \mathbb{R}$, we have $f_\tau(x) \geq \delta$ where $\delta = \min\{c(\tau), 1/2\}$, as desired. ■

Eisenstein Series as a Modular Form: Convergence.

To finish off the proof the absolute convergence of the Eisenstein series, we use the following general result regarding the convergence of general lattice sums.

Proposition

Let d be a positive integer. For $s > 0$, the infinite series $\sum_{\mathbf{a} \in \mathbb{Z}^d - \{0\}} \frac{1}{\|\mathbf{a}\|^s}$ converges for $s > d$ and diverges for $0 < s \leq d$.

Eisenstein Series as a Modular Form: Convergence.

To finish off the proof the absolute convergence of the Eisenstein series, we use the following general result regarding the convergence of general lattice sums.

Proposition

Let d be a positive integer. For $s > 0$, the infinite series $\sum_{\mathbf{a} \in \mathbb{Z}^d - \{0\}} \frac{1}{\|\mathbf{a}\|^s}$ converges for $s > d$ and diverges for $0 < s \leq d$.

Now, we prove the absolute convergence of G_k .

Eisenstein Series as a Modular Form: Convergence.

To finish off the proof the absolute convergence of the Eisenstein series, we use the following general result regarding the convergence of general lattice sums.

Proposition

Let d be a positive integer. For $s > 0$, the infinite series $\sum_{\mathbf{a} \in \mathbb{Z}^d - \{0\}} \frac{1}{\|\mathbf{a}\|^s}$ converges for $s > d$ and diverges for $0 < s \leq d$.

Now, we prove the absolute convergence of G_k .

Proof.

We must show that $\sum_{(m,n) \in \mathbb{Z}^2 - \{(0,0)\}} \frac{1}{|m\tau + n|^k}$ converges for $k \geq 4$. By the previous lemma, we may choose a $\delta_\tau \in (0, 1)$ such that $|m\tau + n|^k \geq \delta_\tau^k |mi + n|^k$ and so $\frac{1}{|m\tau + n|^k} \leq \frac{1}{\delta_\tau^k |mi + n|^k}$, so $\sum_{(m,n) \in \mathbb{Z}^2 - \{(0,0)\}} \frac{1}{|m\tau + n|^k} \leq \delta_\tau^{-k} \sum_{(m,n) \in \mathbb{Z}^2 - \{(0,0)\}} \frac{1}{|mi + n|^k} = \delta_\tau^{-k} \sum_{(m,n) \in \mathbb{Z}^2 - \{(0,0)\}} \frac{1}{(m^2 + n^2)^{k/2}}$. From the previous theorem, we know that as $k > 2$, $\sum_{(m,n) \in \mathbb{Z}^2 - \{(0,0)\}} \frac{1}{(m^2 + n^2)^{k/2}}$ converges, which completes the proof. ■

A Quick Note About the Issue With G_2 .

Remark

Note that G_2 is not absolutely convergent, and is hence not a modular form.

A Quick Note About the Issue With G_2 .

Remark

Note that G_2 is not absolutely convergent, and is hence not a modular form. In fact, there are no modular forms of weight 2.

A Quick Note About the Issue With G_2 .

Remark

Note that G_2 is not absolutely convergent, and is hence not a modular form. In fact, there are no modular forms of weight 2.

Remark

The modularity condition is equivalent to

- $f(\tau + 1) = f(\tau)$,
- $f(-1/\tau) = \tau^k f(\tau)$.

A Quick Note About the Issue With G_2 .

Remark

Note that G_2 is not absolutely convergent, and is hence not a modular form. In fact, there are no modular forms of weight 2.

Remark

The modularity condition is equivalent to

- $f(\tau + 1) = f(\tau)$,
- $f(-1/\tau) = \tau^k f(\tau)$.

This is because the two maps $\tau \mapsto \tau + 1$ and $\tau \mapsto -\tau^{-1}$ generate all of the transformation that correspond to $\mathrm{SL}_2(\mathbb{Z})$.

The Weierstrass \wp Function Again: Definition of Δ .

Recall from class that $\wp(z)$ satisfies the following differential equation

$$(\wp'_{\Lambda(1,\tau)}(z))^2 = 4(\wp_{\Lambda(1,\tau)}(z))^3 - 60E_4(\tau)\wp_{\Lambda(1,\tau)} - 140E_6(\tau).$$

The Weierstrass \wp Function Again: Definition of Δ .

Recall from class that $\wp(z)$ satisfies the following differential equation

$$(\wp'_{\Lambda(1,\tau)}(z))^2 = 4(\wp_{\Lambda(1,\tau)}(z))^3 - 60E_4(\tau)\wp_{\Lambda(1,\tau)} - 140E_6(\tau).$$

- Notice this differential equation is of the form of a elliptic curve over \mathbb{C} : $y^2 = x^3 + ax + b$, where $(x, y) = (\wp_{\Lambda(1,\tau)}(z), \wp'_{\Lambda(1,\tau)}(z))$, $a = -15E_4(\tau)$ and $b = -35E_6(\tau)$.
- Indeed, this effectively sets up a bijection between elliptic curves and elliptic functions.
- The discriminant of a general elliptic curve is $\Delta_{\text{disc}} = 4a^3 + 27b^2$, and $\Delta_{\text{disc}} = 0$ if and only if the cubic equation $x^3 + ax + b = 0$ has repeated roots.

The Weierstrass \wp Function Again: Definition of Δ .

Recall from class that $\wp(z)$ satisfies the following differential equation

$$(\wp'_{\Lambda(1,\tau)}(z))^2 = 4(\wp_{\Lambda(1,\tau)}(z))^3 - 60E_4(\tau)\wp_{\Lambda(1,\tau)} - 140E_6(\tau).$$

- Notice this differential equation is of the form of a elliptic curve over \mathbb{C} : $y^2 = x^3 + ax + b$, where $(x, y) = (\wp_{\Lambda(1,\tau)}(z), \wp'_{\Lambda(1,\tau)}(z))$, $a = -15E_4(\tau)$ and $b = -35E_6(\tau)$.
- Indeed, this effectively sets up a bijection between elliptic curves and elliptic functions.
- The discriminant of a general elliptic curve is $\Delta_{\text{disc}} = 4a^3 + 27b^2$, and $\Delta_{\text{disc}} = 0$ if and only if the cubic equation $x^3 + ax + b = 0$ has repeated roots.

Definition

Define the modular discriminant, denoted by Δ , as a complex valued function defined on the upper half plane by

$$\Delta(\tau) = (-60G_4(\tau))^3 - 27(140G_6(\tau))^2 = -216000G_4^3(\tau) + 529200G_6^2(\tau) = 16\Delta_{\text{disc}}(\tau).$$

In terms of the normalized Eisenstein series, we have

$$\Delta(\tau) = \frac{1}{1728}(E_4^3 - E_6^2).$$

The Modular Discriminant.

Remark

From the properties of $\wp'_{\Lambda(1,\tau)}(z)$, we must have that $\Delta(\tau)$ is non-zero on its domain.

A key identity satisfied by Δ has to do with infinite products.

Theorem

Δ is a modular form of weight 12.

Theorem

We have that

$$\Delta(\tau) = q \prod_{n=1}^{\infty} (1 - q^n)^{24},$$

where $q = e^{2\pi i\tau}$.

We will not prove this here, but see my paper.

The Connection Between G_2 and Δ .

What can we do assuming the product representation for Δ ?

Taking the logarithmic derivative, we get

$$\log(\Delta(\tau)) = \log(q) + 24 \sum_{n=1}^{\infty} \log(1 - q^n),$$

The Connection Between G_2 and Δ .

What can we do assuming the product representation for Δ ?

Taking the logarithmic derivative, we get

$$\log(\Delta(\tau)) = \log(q) + 24 \sum_{n=1}^{\infty} \log(1 - q^n),$$

and so

$$\frac{d}{d\tau}(\log(\Delta(\tau))) = 2\pi i \left(1 - 24 \sum_{n=1}^{\infty} \frac{nq}{1 - q^n} \right).$$

By writing $\sum_{n=1}^{\infty} \frac{nq}{1 - q^n}$ as $\sum_{n=1}^{\infty} \sum_{m=1}^{\infty} nq^{mn}$ and switching the order of summation, we get that $\sum_{n=1}^{\infty} \frac{nq}{1 - q^n} = \sum_{n=1}^{\infty} \sigma_1(n)q^n$,

The Connection Between G_2 and Δ .

What can we do assuming the product representation for Δ ?

Taking the logarithmic derivative, we get

$$\log(\Delta(\tau)) = \log(q) + 24 \sum_{n=1}^{\infty} \log(1 - q^n),$$

and so

$$\frac{d}{d\tau}(\log(\Delta(\tau))) = 2\pi i \left(1 - 24 \sum_{n=1}^{\infty} \frac{nq}{1 - q^n} \right).$$

By writing $\sum_{n=1}^{\infty} \frac{nq}{1 - q^n}$ as $\sum_{n=1}^{\infty} \sum_{m=1}^{\infty} nq^{mn}$ and switching the order of summation, we get that $\sum_{n=1}^{\infty} \frac{nq}{1 - q^n} = \sum_{n=1}^{\infty} \sigma_1(n)q^n$, and so

$$\frac{\Delta'(\tau)}{\Delta(\tau)} = 2\pi i \left(1 - 24 \sum_{n=1}^{\infty} \sigma_1(n)q^n \right) = 2\pi i E_2(\tau).$$

The Connection Between G_2 and Δ .

As Δ is a modular form of weight 12, we have

$$\Delta(-1/\tau) = \tau^{12} \Delta(\tau).$$

Differentiating both sides with respect to τ , we get

$$\Delta'(-1/\tau) = 12\tau^{13} \Delta(\tau) + \tau^{14} \Delta'(\tau).$$

Then,

$$E_2(-1/\tau) = \frac{\Delta'(-1/\tau)}{2\pi i \Delta(-1/\tau)} = \frac{12\tau^{13} \Delta(\tau) + \tau^{14} \Delta'(\tau)}{2\pi i (\tau^{12} \Delta(\tau))} = \frac{6\tau}{\pi i} + \tau^2 E_2(\tau).$$

All in all,

$$E_2(-1/\tau) = \frac{6\tau}{\pi i} + \tau^2 E_2(\tau),$$

meaning E_2 is almost a modular form!

Remark

This property of E_2 is used to create a differential operator, which takes a modular form of weight k to one with weight $k + 2$, called the Serre operator.