

THE EULER CIRCLE COMBINATORIAL GAME THEORY SAMPLE PROBLEM

SIDDHARTH KOTHARI

This document contains my solutions to select The Euler Circle Combinatorial Game Theory Problems. The first one is same one as in the Poster.

1. SAMPLE PROBLEM 1: TINY ANALOGUE OF THE NUMBER AVOIDANCE THEOREM

Question 1.1. Suppose $a > b \geq 0$ are numbers. Find the canonical form for $\star_a \star_b$, i.e. the sum of a and b . Explain why this is a tiny analogue of the number avoidance theorem.¹

We start by adding together \star_a and \star_b , and simplifying the dominated options first. We have

$$\begin{aligned}\star_a + \star_b &= \{0 \mid \{0 \mid -a\}\} + \{0 \mid \{0 \mid -b\}\} \\ &= \{\star_a, \star_b \mid \{0 \mid -a\} + \star_b, \star_a + \{0 \mid -b\}\} \\ &= \{\star_b \mid \{0 \mid -a\} + \star_b, \star_a + \{0 \mid -b\}\},\end{aligned}$$

where the last equality follows since $b < a \implies \star_a < \star_b$, and so the left option \star_a is dominated, and so we can ignore it. Next, we must compare $\{0 \mid -a\} + \star_b$ to $\star_a + \{0 \mid -b\}$. We do this by putting the two in canonical form.

Lemma 1.2. Let $a > b > 0$. Then $\{0 \mid -a\} + \star_b = \{\star_b \mid \{-a \mid \{-a \mid -a - b\}\}\}$. Notice that we have $\{0 \mid -a\} + \star_b = \{\star_b \mid -a + \star_b\}$.

Proof. Adding the two together, we get

$$\{0 \mid -a\} + \{0 \mid \{0 \mid -b\}\} = \{\star_b, \{0 \mid -a\} \mid \{0 \mid -a\} + \{0 \mid -b\}, -a + \star_b\}.$$

First, it can be shown that $\{0 \mid -a\} < \star_b$ (this is because $\{0 \mid -a\} \in \mathcal{N}$ and $\star_b \in \mathcal{L}$), so we can ignore the $\{0 \mid -a\}$. Then, we must compare $\{0 \mid -a\} + \{0 \mid -b\}$ to $-a + \star_b$. Let us put the two in a slightly simpler form.

- We have that $\{0 \mid -a\} + \{0 \mid -b\} = \{\{0 \mid -a\}, \{0 \mid -b\} \mid \{-a \mid -a - b\}, \{-b \mid -a - b\}\}$. Clearly, $\{0 \mid -a\} < \{0 \mid -b\}$, and $\{-a \mid -a - b\} < \{-b \mid -a - b\}$, so we can write

$$\{0 \mid -a\} + \{0 \mid -b\} = \{\{0 \mid -b\} \mid \{-a \mid -a - b\}\}.$$

- We have that $-a + \star_b = \{-a \mid -a + \{0 \mid -b\}\} = \{-a \mid \{-a \mid -a - b\}\}$, by the translation theorem.

Thus, both have the same right option, but the left option of $\{0 \mid -a\} + \{0 \mid -b\}$ is $\{0 \mid -b\}$, whereas the left option of $-a + \star_b$ is $-a$. Thus, when left starts, right will win a larger margin in the latter game, as the former will end at $-b > -a$. Hence, we claim that $-a + \star_b < \{0 \mid -a\} + \{0 \mid -b\}$ (it can be shown that this is indeed the case), and so we can ignore the $\{0 \mid -a\} + \{0 \mid -b\}$ altogether, which completes the proof. ■

Next, we turn our attention to $\{0 \mid -b\} + \star_a = \{\star_a, \{0 \mid -b\} \mid \{0 \mid -a\} + \{0 \mid -b\}, -b + \star_a\}$. Again, since $\star_a \in \mathcal{L}$ and $\{0 \mid -b\} \in \mathcal{N}$, we have that $\{0 \mid -b\} < \star_a$, and so we can ignore the $\{0 \mid -b\}$. As for the right options, we must compare $\{0 \mid -a\} + \{0 \mid -b\}$ and $-b + \star_a$. We

¹Course Description: <https://eulercircle.com/classes/combinatorial-game-theory/>.

already have that $\{0 \mid -a\} + \{0 \mid -b\} = \{\{0 \mid -b\} \mid \{-a \mid -a - b\}\}$. Similarly, $-b + \star_a = -b + \{0 \mid \{0 \mid -a\}\} = \{-b \mid \{-b \mid -a - b\}\} = -\{\{a + b \mid b\} \mid b\}$. It turns out that the game $\{\{a + b \mid b\} \mid b\} + \{\{0 \mid -b\} \mid \{-a \mid -a - b\}\} \in \mathcal{R}$, so $\{0 \mid -a\} + \{0 \mid -b\} < -b + \star_a$, and we can ignore the $-b + \star_a$. Hence, $\{0 \mid -b\} + \star_a = \{\star_a \mid \{0 \mid -a\} + \{0 \mid -b\}\}$.

Lastly, we claim that $\{0 \mid -a\} + \star_b < \star_a + \{0 \mid -b\}$. That is, the game

$$\{a - \star_b \mid -\star_b\} + \{\star_a \mid \{0 \mid -a\} + \{0 \mid -b\}\} \in \mathcal{L}.$$

Proof. Right to start has the following options.

- He can move to $-\star_b$, to which left will respond by moving to $\{b \mid 0\}$ in $-\star_b$, leaving us with $\{b \mid 0\} + \{\star_a \mid \{0 \mid -a\} + \{0 \mid -b\}\}$. It is easy to show that right has no winning moves from here.
- Right can move to $\{0 \mid -a\} + \{0 \mid -b\}$, to which left will respond by moving to $a - \star_b$, which will result in $a - \star_b + \{0 \mid -a\} + \{0 \mid -b\}$, which we've already shown is a \mathcal{L} position.

Left to start will move to $a - \star_b$, which will result in $\{\{a + b \mid a\} \mid a\} + \{\star_a \mid \{0 \mid -a\} + \{0 \mid -b\}\}$. One can show that right has no winning moves from here. ■

Here are some the results that I used in my proof above.

Lemma 1.3. *Let $a > b$. Then $-a + \star_b < \{0 \mid -a\} + \{0 \mid -b\}$.*

Proof. We must show that $\{\{0 \mid -b\} \mid \{-a \mid -a - b\}\} + \{\{a + b \mid a\} \mid a\} \in \mathcal{L}$. Right to start has the following options.

- He may move to $\{-a \mid -a - b\}$, in which case left will respond by moving to $\{a + b \mid a\}$, which will result in the 0 game.
- He may move to a , to which left will respond by moving to $\{0 \mid -b\}$, which will result in $\{0 \mid -b\} + a = \{a \mid a - b\} \in \mathcal{L}$.

Left's to start will move to $\{0 \mid -b\}$, which will result in $\{0 \mid -b\} + \{\{a + b \mid a\} \mid a\}$, which is losing for right. ■

Lemma 1.4. *Let $a > b$. Then $\{0 \mid -a\} + \{0 \mid -b\} < -b + \star_a$.*

Proof. We must show that $\{\{0 \mid -b\} \mid \{-a \mid -a - b\}\} + \{\{a + b \mid b\} \mid b\} \in \mathcal{R}$. Left to start has the following options.

- She can move to $\{0 \mid -b\}$, to which right will respond by moving to $-b$ in $\{0 \mid -b\}$, leaving us with $\{\{a \mid 0\} \mid 0\} \in \mathcal{R}$.
- She can move to $\{a + b \mid b\}$, to which right will respond by moving to b in $\{a + b \mid b\}$, leaving us with $\{\{b \mid 0\} \mid \{b - a \mid -a\}\}$. Left has no other option but to move to the \mathcal{N} position $\{b \mid 0\}$.

Right to start will move to b . ■

In conclusion, we have the following result.

Proposition 1.5. *Let $a > b$ be numbers. Then $\star_a + \star_b = \{\star_b \mid \{0 \mid -a\} + \star_b\}$.*

2. SAMPLE PROBLEM 2: SUBTRACTION GAME AND PERIODIC GRUNDY VALUES

Question 2.1. *Let S be a finite set of positive integers. Consider the S -subtraction game, in which a move consists of removing stones from the pile, where s is some element in S . Show that the Grundy values of this game are eventually periodic, i.e. there is some integer p and some nonnegative integer n_0 so that $\mathcal{G}(n + p) = \mathcal{G}(n)$ for all integers $n \geq n_0$.*

Lemma 2.2. *Let S be a set of non-negative integers. Then $\text{mex}(S) \leq |S|$. In other words, the minimal excludant of a set cannot exceed the size of the set itself.*

Proposition 2.3. *There are only finitely many Grundy values of the S -subtraction game. In other words, the set $\{\mathcal{G}(n) : n \in \mathbb{Z}^+ \cup \{0\}\}$ is a finite set of non-negative integers.*

Proof. From the definition of Grundy values, for all $n \in \mathbb{Z}^+ \cup \{0\}$, we have $\mathcal{G}(n) = \text{mex}(\{\mathcal{G}(n-s) : s \in S \text{ and } n-s \geq 0\})$. Clearly, $|\{\mathcal{G}(n-s) : s \in S \text{ and } n-s \geq 0\}| \leq |S| \implies \mathcal{G}(n) \leq |S|$ (this follows from the lemma stated above). Thus, $\mathcal{G}(n)$ can only take on finitely values. ■

Remark 2.4. In fact, we have that the Grundy values are less than or equal to $|S|$, the maximum number of legal moves available to a player. Thus, $\mathcal{G}(n)$ can take on a maximum of $|S| + 1$ distinct values ($\mathcal{G}(n) \in \{0, 1, \dots, |S|\}$).

Proof. For any positive integer m , there exist non-negative integers $n_0(m)$ and $n_1(m)$ such that $n_0(m) < n_1(m)$ and $\mathcal{G}(n_0(m) - s + i) = \mathcal{G}(n_1(m) - s + i)$ for all $s \in S$ and for $0 \leq i \leq m$. This follows from the pigeonhole principle. To see how, for each non-negative integer n we associate the $(m+1)$ by k matrix given by $A(n)_{ij} = \mathcal{G}(n + i - s_j)$ for $0 \leq i \leq m$ and $1 \leq j \leq k$. Now, each Grundy value of the S -subtraction game is between 0 and k (both inclusive), so we have $(k+1)^{(m+1)k}$ choices for $A(n)_{ij}$ for each n , which is clearly finite. Thus, somewhere down the line of integers, two integers must share the same matrix.

Now, let us take m to be $s_k - 1$. We denote the corresponding $n_0(m)$ by n_0 and $n_1(m)$ by n_1 . Note that we have $\mathcal{G}(n_0 + k) = \mathcal{G}(n_1 + k)$ for all $0 \leq k \leq m$. We will show that $\mathcal{G}(n_0 + m + j) = \mathcal{G}(n_1 + m + j)$ for all $j > 0$. We will use induction on j .

For the base case, that is, when $j = 1$, we must show that $\mathcal{G}(n_0 + m + 1) = \mathcal{G}(n_1 + m + 1)$. Note that $\mathcal{G}(n_0 + m + 1) = \text{mex}\{\mathcal{G}(n_0 + m + 1 - s) : s \in S\} = \text{mex}\{\mathcal{G}(n_0 + s_k - s) : s \in S\}$. At this point, observe that $0 \leq s_k - s$ and $1 \leq s \implies -1 \geq -s \implies s_k - 1 \geq s_k - s \implies m \geq s_k - s$ for all $s \in S$. Thus, $0 \leq s_k - s \leq m$, and so by the induction hypothesis, we have that $\mathcal{G}(n_0 + s_k - s) = \mathcal{G}(n_1 + s_k - s)$ for all $s \in S$, meaning that $\mathcal{G}(n_0 + m + 1) = \mathcal{G}(n_1 + m + 1)$.

Next, assume that $\mathcal{G}(n_0 + m + x) = \mathcal{G}(n_1 + m + x)$ for all $0 \leq x \leq j$, which implies $\mathcal{G}(n_0 + m + x) = \mathcal{G}(n_1 + m + x)$ for all $-m \leq x \leq j$. We must show that $\mathcal{G}(n_0 + m + j + 1) = \mathcal{G}(n_1 + m + j + 1)$. Observe that $\mathcal{G}(n_0 + m + j + 1) = \text{mex}\{\mathcal{G}(n_0 + m + j + 1 - s) : s \in S\}$. Note that $1 \leq s \implies 1 - s \leq 0 \implies j + 1 - s \leq j$ and $s \leq s_k \implies s - j \leq s_k \implies j - s \geq -s_k \implies 1 + j - s \geq 1 - s_k \implies j + 1 - s \geq -m$. Thus, $-m \leq j + 1 - s \leq j$ for all $s \in S$. Hence, by the induction hypothesis, $\mathcal{G}(n_0 + m + j + 1 - s) = \mathcal{G}(n_1 + m + j + 1 - s)$ for all $s \in S$ which means that $\mathcal{G}(n_0 + m + j + 1) = \mathcal{G}(n_1 + m + j + 1)$.

In conclusion, we have that $\mathcal{G}(n_0 + x) = \mathcal{G}(n_1 + x)$ for all $x \geq 0$. Since $n_1 > n_0$, we may write $n_1 = n_0 + p$ where $p > 0$. Thus, $\mathcal{G}(n_1 + x) = \mathcal{G}(n_0 + x + p) = \mathcal{G}(n_0 + x)$. Setting $n = n_0 + x$, we get that $\mathcal{G}(n) = \mathcal{G}(n + p)$ for all $n \geq n_0$. ■

Remark 2.5. In fact, this proof gives us a bound on p . Namely, $p \leq (k+1)^{k(m+1)} = (k+1)^{ks_k} = (|S| + 1)^{|S| \times \sup(S)}$.

3. SAMPLE PROBLEM 3: A PROOF

Question 3.1. *Show that if $G = \{G^{L_1}, \dots \mid G^{R_1}, \dots\}$ is a game born on day n and $s, t \geq n$, then we may replace G^{L_1} by $\{s \mid \{G^{L_1} \mid -t\}\}$ without changing the value.*

Lemma 3.2. *Let G be a game born on day n . Then $G \leq n$.*

Proof. We will use induction on the birthday of G . That is, assuming that $G' \leq g(G')$ for all games G' such that $g(G') < g(G)$, we will show that $G \leq n$. In other words, we must show that $n - G = \{n - 1 \mid \emptyset\} + \{-G^R \mid -G^L\} \in \mathcal{L}$.

If left starts, her strategy will be to move to a $-g^R \in -G^R$, which will reduce the game to $n - g^R$. By the induction hypothesis, since $g(g^R) < g(G) = n$ we must have that $g^R \leq g(g^R) < n \implies g^R < n \implies n - g^R \in \mathcal{L}$, so left has a winning strategy in $n - g^R$. If right starts, he has no other

option but to move to a $-g^L \in -G^L$, which will reduce the game to $n - g^L$. Just like last time one can show that $n - g^L \in \mathcal{L}$, which completes the proof. ■

Lemma 3.3. *Let G be a game born on day n , and let $x \geq n$ be an integer. Then $-x - g^R \in \mathcal{R}$ for all right options g^R of G .*

Proof. Recall that $-g^R$ is a option of $-G$, which is a game born on day n , so $-g^R$ is born before day n . Hence, we know that $-g^R \leq n$ by the previous lemma. Since we are given that $n \leq x$, we must have that $-g^R < x \implies -x - g^R \in \mathcal{R}$ by transitivity. ■

Proof. We must show that the game $\{\{s \mid \{G^L \mid -t\}\}, G^{L_i} \mid G^{R_i}\} + \{-G^{R_i} \mid -G^{L_i}\}$ is a \mathcal{P} position. If left starts, she has the following options.

- She may choose to move to $\{s \mid \{G^L \mid -t\}\}$ in $\{\{s \mid \{G^L \mid -t\}\}, G^{L_i} \mid G^{R_i}\}$ which leaves us with $\{s \mid \{G^L \mid -t\}\} + \{-G^{R_i} \mid -G^{L_i}\}$. Right will respond by moving to $\{G^L \mid -t\}$ in $\{s \mid \{G^L \mid -t\}\}$, which will reduce the game to $\{G^L \mid -t\} + \{-G^{R_i} \mid -G^{L_i}\}$. Now, if left responds by moving to G^L in $\{G^L \mid -t\}$, right will move to $-G^L$ in $\{-G^{R_i} \mid -G^{L_i}\}$, which will reduce the game to 0, which is losing for left. On the other hand, if left moves to $-G^{R_i}$ in $\{-G^{R_i} \mid -G^{L_i}\}$, right will respond by moving to $-t$ in $\{G^L \mid -t\}$ which will reduce the game to $-t - G^{R_i}$. By the lemma mentioned above, we know that this is a \mathcal{R} position.
- She may choose to move to G^{L_i} for $i \geq 2$ in $\{\{s \mid \{G^L \mid -t\}\}, G^{L_i} \mid G^{R_i}\}$. Right will respond by moving to $-G^{L_i}$ in $\{-G^{R_i} \mid -G^{L_i}\}$, which will make the game 0.
- She may choose to move to $-G^{R_i}$ in $\{-G^{R_i} \mid -G^{L_i}\}$. Right will respond by moving to G^{R_i} in $\{\{s \mid \{G^L \mid -t\}\}, G^{L_i} \mid G^{R_i}\}$, which will make the game 0 again.

In all the cases, left loses. If right starts, he has the following options.

- He may choose to move to $-G^L$ in $\{-G^{R_i} \mid -G^{L_i}\}$ which leaves us with $\{\{s \mid \{G^L \mid -t\}\}, G^{L_i} \mid G^{R_i}\} - G^L$. Left will respond by moving to $\{s \mid \{G^L \mid -t\}\}$ in the first game, after which we have $\{s \mid \{G^L \mid -t\}\} - G^L$. Right has two options. If he moves to $\{G^L \mid -t\}$ he will loose, as left will respond by moving to G^L in $\{G^L \mid -t\}$, which will reduce the game to 0. On the other hand, if he moves to some right option of $-G^L$, call it $-G^{LR}$, left will respond by moving to s in $\{s \mid \{G^L \mid -t\}\}$, which will reduce the game to $s - G^{LR}$. Note that this game is of the form $s - h$, where h is a left option of a left option of G . Thus, we know that h is born before G , and so by the above lemma we have that $h \leq n$ and since $n < s$ we have that $h < s \implies s - h \in \mathcal{L}$, which means that left has a winning strategy in $s - G^{LR}$ so she wins. ■