

CMS data and their interpretation using the phenomenological MSSM

Maurizio Pierini

CERN

Maria Spiropulu


1 Introduction

With the extremely successful performance of both machine and detectors at $\rho_{\bar{s}} = 7$

4. 3 trilinear couplings: $A_{b,t,\tau}$,
5. 1 pseudo-scalar Higgs mass: m_A .
6. 1 parameter.

The parameters m_A and $\tan\beta$ can be swapped for the Higgs mass parameters m_{H_u} and

We weight every pMSSM point with the product of the three likelihoods, one for each of the three analyses. Since the pMSSM points were sampled from a (bounded) uniform 19-dimensional density, the set of weights



d.

Figure 1: Ratios of profile likelihood L_p to maximum likelihood L_{max} shown for the squark mass parameters at SUSY scale. The colored and shaded histograms show the distributions before and after the inclusion of the CMS results.




Figure 2: Ratios of profile likelihood L_p to maximum likelihood L




Figure 3: Ratios of profile likelihood L_p / L




Figure 4: Ratios of profile likelihood L_p to maximum likelihood L_{max} shown for trilinear couplings at SUSY scale. The colored and shaded histograms show the distributions before and after the inclusion of the CMS results.





Figure 6: Ratios of profile likelihood L_p to maximum likelihood L



Figt57c666.492 00e866 155-351(Ratios0e866 of0e861(profil00e866 lik)10(elihood))TJ/F1f 70.866 15515827068837c666L))TJ/F10 0.7




Figure 10: Ratios of profile likelihood L_p to maximum likelihood L_{max} shown for chargino masses. The colored and shaded histograms show the distributions before and after the inclusion of the CMS results.




Figure 11: Ratios of profile likelihood L_p to maximum likelihood L_{max} shown for the Higgs masses. The colored and shaded histograms show the distributions before and after the inclusion of the CMS results.




Figure 12: Ratio of profile likelihood L_p to maximum likelihood L_{max} shown for lightest neutralino dark matter relic density. The colored and shaded histograms show the results before and after the inclusion of the CMS results.



Figure 14: Ratios of profile likelihood L_p to maximum likelihood L_{max} shown for predictions for weak scale observables as calculated by superiso - I. The colored and shaded histograms show the distributions before and




Figure 15: Ratios of profile likelihood L_p to maximum likelihood L_{max} shown for predictions for weak scale observables as calculated by superiso - II. The colored and shaded histograms show the distributions before and after the inclusion of the CMS results.

Appendix

