

SUSY discovery potential using dileptons at $\sqrt{s} = 7 \text{ TeV}$

D. Barge, C. Campagnari, P. Kalavase, D. Kovalskyi, V. Krutelyov, J. Ribnik

University of California, Santa Barbara

W. Andrews, D. Evans, F. Golf, J. Mülmenstädt, S. Padhi, Y. Tu, F. Würthwein, A. Yagil

University of California, San Diego

L. Bauerdick, I. Bloch, K. Burkett, I. Fisk, O. Gutsche

Fermi National Accelerator Laboratory, Batavia, Illinois

Abstract

The SUSY discovery potential of CMS detector at the LHC is evaluated in different regions of the mSUGRA parameter space. The sensitivities studies are performed for inclusive searches involving same and opposite sign dileptons. These searches for new physics is characterized by large \cancel{E}_T and significant hadronic activity. The study shows significant sensitivity in several regions in the parameters space with 100 pb^{-1} and 1 fb^{-1} of integrated luminosity.

- 1 Introduction**
- 2 Data Samples**
- 3 Event Selection**
- 4 mSUGRA discovery reach**
- 5 Same Sign Dileptons**
- 6 Opposite Sign Dileptons**
- 7 Conclusion**

References

- [1] <http://www.ge.infn.it/~tosi/cms/topMC.html>.
- [2] “Prospects for measuring the WW production cross section in pp collisions at $\sqrt{s}=10$ TeV”, CMS AN-2009/042 and PAS EWK-09-002.
- [3] “Expectations for observation of top quark pair production in the dilepton final state with the early CMS data”, CMS AN-2009/050 and PAS TOP-09-002.
- [4] <https://hypernews.cern.ch/HyperNews/CMS/get/muon/258.html>.
- [5] “Muon Identification in CMS”, CMS AN-2008/098.
- [6] “Data-driven methods to estimate the electron and muon fake contributions to lepton analyses”, CMS AN-2009/041.
- [7] “Data driven background study for new physics searches with same sign dileptons at $\sqrt{s}=10$ TeV”, CMS AN-2009/138.
- [8] “Data driven background estimate for a new physics search with opposite sign dileptons”, CMS AN-2009/130.