In [215…
```
import numpy as np
import pandas as pd
import matplotlib.pyplot as plt
from scipy import odr
```

Lab Portion

Calibration

In [216…
```
centroids = np.array([680.33, 769.49, 215.34, 392.4])
centroid_fwhm = np.array([29.9564, 31.1873, 16.8211, 28.4043])
energies = np.array(
    [
        1173.2,
        1332.5,
        356.0,
        661.6,
    ]
)
```

In [217…
```
df = pd.DataFrame(
    {
        "Centroids": centroids[: len(energies)],
        "Energies": energies,
    }
)

df["Centroid_unc"] = centroid_fwhm / 2
df["Energy_unc"] = 0  # For the purposes of ODR
df
```

Out[217…

	Centroids	Energies	Centroid_unc	Energy_unc
0	680.33	1173.2	14.97820	0
1	769.49	1332.5	15.59365	0
2	215.34	356.0	8.41055	0
3	392.40	661.6	14.20215	0

In [218…
```
data = odr.RealData(
    df["Centroids"],
    df["Energies"],
    sx=df["Centroid_unc"],
)


def regres_model(p, x):
    return p[0] * x + p[1]
```

```
model = odr.Model(regres_model)
odr_fit = odr.ODR(data, model, beta0=[1, 0])
out = odr_fit.run()

slope = out.beta[0]
intercept = out.beta[1]
slope_std_err = out.sd_beta[0]
intercept_std_err = out.sd_beta[1]
# print("Errors:", slope_std_err, intercept_std_err)
```

In [219…

```
def channel_to_energy(channel):
    return slope * channel + intercept


def channel_to_energy_unc(channel, channel_unc):
    """
    Obtains uncertainties associated with an energy
    found using the linear regression model.
    """

    # Propagate uncertainties through product
    m_x = slope * channel
    m_perc = slope_std_err / slope
    channel_perc = channel_unc / channel
    m_x_unc = m_x * np.sqrt(m_perc**2 + channel_perc**2)

    # Propagate uncertainties through sum
    outer_err = np.sqrt(m_x_unc**2 + intercept_std_err**2)

    return outer_err
```

In [245…

```
plt.figure(figsize=(10, 6))
# plt.scatter(
#     df["Centroids"],
#     df["Energies"],
#     color="blue",
#     marker="o",
#     label="Channel-Energy Known Points",
# )
plt.errorbar(
    df["Centroids"],
    df["Energies"],
    color="blue",
    fmt="o",
    label="Channel-Energy Known Points",
    xerr=df["Centroid_unc"],
    capsize=3
)
plt.xlabel("Centroids")
plt.ylabel("Energies (keV)")
plt.title("Centroids vs Energies")
plt.grid(True)
```

```
c_linspace = np.linspace(0, 800)
e_linspace = channel_to_energy(c_linspace)

line = slope * df["Centroids"] + intercept
plt.plot(
    c_linspace,
    e_linspace,
    color="red",
    linestyle="--",
    label="Regression line",
)
plt.legend()

plt.tight_layout()
plt.show()
```



Unknown-Sample Data

In [221…

```
unknown_centroids = np.array([135.37, 349.16, 488.61])
unknown_fwhm = np.array([29.2727, 2.5797, 28.4532])
```

Analysis

Question 1

In [222…

```
step3 = np.array([149, 118, 137, 128, 128, 135, 112, 133, 133, 121])
```

In [223…
```
sample_std = step3.std(ddof=1)
step3_mean = step3.mean()
theor_std = np.sqrt(step3_mean)
print(sample_std / theor_std)
```

```
0.9296017254842825
```

The ratio of the sample standard deviation to the theorical standard deviation (square root of the sample mean) is about 0.93, well within the 0.85 to 1.15 range. This provides evidence that the data is reasonably close to following a Gaussian distribution.

Question 2

In [224…
```
step3_sdom = sample_std / np.sqrt(len(step3))
step3_time = 20
step3_time_unc = 0.01
step3_rate = step3_mean / step3_time
sdom_perc = step3_sdom / step3_mean
time_perc = step3_time_unc / step3_time
step3_rate_unc = step3_rate * np.sqrt(sdom_perc**2 + time_perc**2)
print(step3_rate, step3_rate_unc)
```

```
6.470000000000001 0.16723044214662458
```

Using the data from step 3, the channel-400 peak's count rate is (6.47 ± 0.17) counts per second.

Question 3

In [225…
```
channel_to_energy(unknown_centroids)
```

Out[225…
```
array([214.00420105, 590.3972175 , 835.90919721])
```

In [226…
```
channel_to_energy_unc(unknown_centroids, unknown_fwhm / 2)
```

Out[226…
```
array([26.02420863,  4.93065895, 25.55707506])
```

The energies corresponding to the three peaks found are:

- 210 $\pm$ 30 keV
- 590 $\pm$ 5 keV
- 840 $\pm$ 30 keV

Question 4

Mn-54.

Question 5

The absolute shortest half life I would expect would be on the order of a month or two months; these elements do not remain radioactive for long enough to be of use in our labs. The absolute largest half life I would expect is on the order of thousands of years; past this point, it seems unlikely that the element is radioactive enough to be considered interesting for analysis.

Mn-54 is the most likely candidate for this unknown element. Although two lower-energy peaks were included in the energies above, these two peaks were not nearly as prominent as the 835-keV peak in the spectrum collected from the sample; I focused on this peak. The reference table provided includes a data point for Mn-54 with an energy of 834.8, well within 1 σ of my energy measurement. Further, there are no other energies listed for Mn-54 in the reference table, in agreement with the single peak observed on the spectrum. The half-life listed for Mn-54 is 312.2 days, which aligns with the expectation of a half life on the order of a few years.

Written Portion

Question 3

In [227…
```
meas = np.array([1.59, 1.64, 1.63, 1.52, 1.69, 1.58]) * 1e-19
```

In [228…
```
meas.mean()
```

Out[228…
```
np.float64(1.6083333333333334e-19)
```

In [229…
```
sdom = meas.std(ddof=1) / np.sqrt(len(meas))
sdom
```

Out[229…
```
np.float64(2.386303510546059e-21)
```

In [230…
```
sdom * 1e3
```

Out[230…
```
np.float64(2.386303510546059e-18)
```

In [231…
```
masses = np.array(
    [
        9.25,
        9.17,
        9.22,
        9.10,
        9.19,
        9.05,
```

```
    ]
)
```

In [232…
```
masses.mean()
```

Out[232…
```
np.float64(9.163333333333334)
```

In [233…
```
masses.std() / np.sqrt(len(masses))
```

Out[233…
```
np.float64(0.028054180384339072)
```

In [234…
```
# def unc_e_sq(e, unc_e):
#     return 2 * e * unc_e
#
#
# def unc_m_cu(m, unc_m):
#     return 3 * (m**2) * unc_m


def dBde(e, m):
    return 4 * e / m**3


def dBdm(e, m):
    return -6 * e**3 / m**4


def unc_total(e, m, unc_e, unc_m):
    t1 = dBde(e, m) * unc_e
    t2 = dBdm(e, m) * unc_m
    return np.sqrt(t1**2 + t2**2)
```

In [235…
```
unc_total(1.61e-19, 9.16e-31, 0.02e-19, 0.03e-31)
```

Out[235…
```
np.float64(1.675827998510116e+51)
```

In [236…
```
e = 1.61e-19
m = 9.16e-31
2 * (e**2 / m**3)
```

Out[236…
```
6.745207694003217e+52
```

Question 4

In [237…
```
plates = np.array([5.16, 4.47, 5.26, 4.50, 5.18, 5.12])
plates_unc = np.array([0.01] * 6)

thickness = [plates[: i + 1].sum() for i in range(len(plates))]
thickness_unc = [plates_unc[: i + 1].sum() for i in range(len(plates))]
```

In [238…
```
net_area = np.array([4454, 2556, 2250, 1432, 1086, 672])
net_area_unc = np.sqrt(net_area)
```

In [239…

```
dfp4 = pd.DataFrame()
dfp4["x"] = thickness
dfp4["x_unc"] = thickness_unc
dfp4["A"] = net_area.astype(float)
dfp4["A_unc"] = np.sqrt(net_area)
dfp4
```

Out[239…

	x	x_unc	A	A_unc
0	5.16	0.01	4454.0	66.738295
1	9.63	0.02	2556.0	50.556899
2	14.89	0.03	2250.0	47.434165
3	19.39	0.04	1432.0	37.841776
4	24.57	0.05	1086.0	32.954514
5	29.69	0.06	672.0	25.922963

In [240…

```
def lnA_unc(A, A_unc):
    return (A**-1) * A_unc


dfp4["lnA"] = np.log(dfp4["A"])
dfp4["lnA_unc"] = lnA_unc(dfp4["A"], dfp4["A_unc"])
dfp4
```

Out[240…

	x	x_unc	A	A_unc	lnA	lnA_unc
0	5.16	0.01	4454.0	66.738295	8.401558	0.014984
1	9.63	0.02	2556.0	50.556899	7.846199	0.019780
2	14.89	0.03	2250.0	47.434165	7.718685	0.021082
3	19.39	0.04	1432.0	37.841776	7.266827	0.026426
4	24.57	0.05	1086.0	32.954514	6.990257	0.030345
5	29.69	0.06	672.0	25.922963	6.510258	0.038576

In [241…

```
data = odr.RealData(
    dfp4["x"],
    dfp4["lnA"],
    sx=dfp4["x_unc"],
    sy=dfp4["A_unc"],
)


def regres_model(p, x):
    return p[0] * x + p[1]


model = odr.Model(regres_model)
odr_fit = odr.ODR(data, model, beta0=[1, 0])
```

```
out = odr_fit.run()

slope_p4 = out.beta[0]
intercept_p4 = out.beta[1]
slope_std_err_p4 = out.sd_beta[0]
intercept_std_err_p4 = out.sd_beta[1]
```

In [242…

```
plt.figure(figsize=(10, 6))
plt.xlabel("Lead Thickness (cm)")
plt.ylabel("Log of Net Area in ROI")
plt.title("Log of Net Area in ROI vs. Shielding Lead Thickness")
plt.grid(True)

x_linspace = np.linspace(0, 30)
A_linspace = slope_p4 * x_linspace + intercept_p4
plt.plot(
    x_linspace,
    A_linspace,
    color="red",
    linestyle="--",
    label="Regression line",
)
plt.errorbar(
    dfp4["x"],
    dfp4["lnA"],
    xerr=dfp4["x_unc"],
    yerr=dfp4["lnA_unc"],
    fmt="o",
    capsize=3,
    label="Area vs. Thickness Data Points",
)
plt.legend()
plt.tight_layout()
plt.show()
```



In [243…

```
print(slope_p4)
print(intercept_p4)
print(slope_std_err_p4)
print(intercept_std_err_p4)
```

```
-0.0727015488593651
8.70231989095447
0.004774243977844703
0.11144923418905714
```

In [ ]:

Amptek DppMCA live_data.mca

File View MCA Display Analyze DPP Help



MCA8000D (s/n	392)
Tag:	live_data
Mode	MCA
Channels	1,024
LLD Thresh	0.50% FS
Input Range	10V
Preset Mode	Live
Preset	200.000
Real Time	33.23
Live Time	33.12
Total Count	24,439
Total Rate	737.89
Start:	12/02/2025 15:15:19
Status:	stopped

Peak Information:

Centroid (N)	504.50
FWHM (N)	28.1598
Net Area	4,598
Uncertainty	1.69
Net Rate	138.83
Gross Area	5,313

Options Option PA

	Cursor	Range		LOG Scale
Channel	505	0	1023	
Count	139	0	31744	

Ready WinUSB NUM