










Shawn Pan - AM205 Homework 5 Code is attached as separate files.

Problem 1

Method	Starting Point	Steps	Final Point
Steepest Descent	$(-1, 1)$	2000	$(0.962, 0.926)$
Steepest Descent	$(0, 1)$	2000	$(0.967, 0.934)$
Steepest Descent	$(2, 1)$	980	$(1, 1)$
Newton	$(-1, 1)$	3	$(1, 1)$
Newton	$(0, 1)$	6	$(1, 1)$
Newton	$(2, 1)$	6	$(1, 1)$
BFGS	$(-1, 1)$	124	$(1, 1)$
BFGS	$(0, 1)$	38	$(1, 1)$
BFGS	$(2, 1)$	45	$(1, 1)$







Problem 2

(a)


Some convenient intermediate terms:

$$\begin{aligned}\frac{dx}{ds} &= \frac{L}{R} + \sum_{k=1}^{20} \frac{\pi k c_k}{R} \cos \frac{\pi k s}{R} \\ \frac{dy}{ds} &= \sum_{k=1}^{20} \frac{\pi k d_k}{R} \cos \frac{\pi k s}{R} \\ \frac{\partial}{\partial c_i} \frac{dx}{ds} &= \frac{\pi i}{R} \cos \frac{\pi i s}{R} \\ \frac{\partial}{\partial d_i} \frac{dy}{ds} &= \frac{\pi i}{R} \cos \frac{\pi i s}{R} \\ \frac{\partial y}{\partial d_i} &= \sin \frac{\pi i s}{R} \\ A &= \sqrt{\left(\frac{dx}{ds}\right)^2 + \left(\frac{dy}{ds}\right)^2}\end{aligned}$$

Gradient components (expressed with intermediate terms above):

$$\begin{aligned}\frac{\partial r}{\partial c_i} &= \int_0^R \mu(2) (A-1) \frac{1}{2A} (2) \left(\frac{dx}{ds}\right) \left(\frac{\partial}{\partial c_i} \frac{dx}{ds}\right) ds \\ &= \int_0^R 2\mu \frac{A-1}{A} \left(\frac{dx}{ds}\right) \left(\frac{\partial}{\partial c_i} \frac{dx}{ds}\right) ds \\ \frac{\partial r}{\partial d_i} &= \int_0^R \left[2\mu \frac{A-1}{A} \left(\frac{dy}{ds}\right) \left(\frac{\partial}{\partial d_i} \frac{dy}{ds}\right) - 2\rho\omega^2 y \left(\frac{\partial y}{\partial d_i}\right) \right] ds\end{aligned}$$

(b/c)



$\mu = 2000$ appears to converge to the same solution, while $\mu = 20$ and $\mu = 200$ appear to converge to alternate solutions reflected across the x -axis. Because all y terms in the residual function are squared, the problem is symmetric and the reflected solutions (shown in grey) have the same energy.

Problem 3

(a)


We will use the centered finite difference approximation:



$$\frac{\partial^2 \Psi}{\partial x^2} \approx \frac{\Psi(x-h) - 2\Psi(x) + \Psi(x+h)}{h^2}$$

We can express the equation as $H\Psi = E\Psi$, where H is the following matrix with v_i as the potential evaluated on the grid point i .

$$\begin{pmatrix} \frac{2}{h^2} + v_0 & \frac{-1}{h^2} & 0 & 0 & \dots & 0 & 0 \\ \frac{-1}{h^2} & \frac{2}{h^2} + v_1 & \frac{-1}{h^2} & 0 & \dots & 0 & 0 \\ 0 & \frac{-1}{h^2} & \frac{2}{h^2} + v_2 & \frac{-1}{h^2} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & \frac{-1}{h^2} & \frac{2}{h^2} + v_{1920} \end{pmatrix}$$

Note that truncating the first and last rows is conveniently equivalent to enforcing the zero Dirichlet boundary conditions. (Technically, this enforces $\Psi(-12-h) = 0$ and $\Psi(12+h) = 0$, but shifting one step doesn't change the numerical result.)





(b)

Probability particle lies between 0 and 6 for v_2 .

Energies [0.39068 1.04243 1.40185 1.60332 1.94743]

Probability [3.07578e-04 2.99542e-02 7.86516e-01 3.99204e-01 5.33140e-01]