Net contents: Equivalent to 4 grams amoxicillin.

Directions for mixing: Tap bottle until all powder flows freely. Add approximately 1/3 total amount of water for reconstitution (total = 55 mL): shake vigorously to wet powder. Add remaining water; again shake vigorously. Usual Adult Dosage: 250 to 500 mg every 8 hours. Usual Child Dosage: 20 to 40 mg/kg/day in divided doses every 8 hours, depending on age, weight, and infection severity. See accompanying prescribing information.

Store dry powder at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Keep tightly closed. Shake well before using. Refrigeration preferable after reconstitution but not required. Discard suspension after 14 days.

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

Mfd. For: Dava Pharmaceuticals, Inc., a Subsidiary of Qualitest Pharmaceuticals. Huntsville, AL 35811 By: Sandoz GmbH in Austria, Product of Spain



NDC 67253-148-45

AMOXICILLIN

FOR ORAL SUSPENSION, USP



When reconstituted, each 5 mL (1 teaspoonful) will contain amoxicillin trihydrate equivalent to 250 mg amoxicillin

80 mL (when reconstituted)

Rx only



Rev. 12-14M

46152660

EXP. 107



NDC 70954-**058**-10 **100 Tablets**

PredniSONE Tablets, USP

5 mg

Each tablet contains 5 mg of prednisone, USP.



Rx only

Usual Dosage: See accompanying package insert for complete prescribing information.

Store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature].

Protect from moisture.

Dispense in a tight, light-resistant container as defined in the USP Keep container tightly closed.

This package is child-resistant.

Keep this and all medications out of the reach of children.

Manufactured by: **Novitium Pharma LLC**

70 Lake Drive, East Windsor New Jersey 08520

Revised: 03/19

LB4088-01



660401

DiphenhydrAMINE Hydrochloride

Injection, USP

50 mg per mL

HIGH POTENCY

For IM or IV use. Protect from light. Keep from freezing.

1 mL Single Dose Vial Rx only

ш







