

Singapore Migrant Worker Dormitories

Team 3-Irises: Sparsh Gupta, Belén
Hutchins, Swasti Jain




How do different policies, movements, and interactions as factors of social connectivity in migrant workers' dorms affect the transmission of disease?

What makes it interesting and important?


- Based on the case-study of migrant workers' dorms in Singapore during COVID-19
- Factors of social connectivity
- Government policy changes (example: isolation areas)

Explanatory Question



Agent-based SIR Model

The social connectivity digram for the podded-agent-based SIR model



The pods used in the model are:

- D_1: overcrowded dorm
(higher infectivity)
- D_2: normal dorm
(normal infectivity)
- C_1: outbreak-center construction site
(higher infection rates)
- C_2: construction site
(normal infection rate)
- M: mall
(higher infection rates.)
- I: isolated area/site
(lowest infectivity)

Update state Equations

$$I = I \mid v_{infect} \ \& \ (\sim R)$$

$$R = R \mid (I \ \& \ v_{recover})$$

$$I = I \ \& \ (\sim R)$$

$$\text{dim} = \text{length}(I_{v_0})$$

$$I_h = \text{zeros}(\text{dim}, T)$$

$$R_h = \text{zeros}(\text{dim}, T)$$

$$S_h = \text{ones}(\text{dim}, T) - I_h - R_h$$

Edge weights used in the model:



- $n_{(D_1)}$: very high infection rate
- $n_{(D_2)}$: normal infection rates of COVID-19
- $n_{(C_1)}$: high infection rate
- $n_{(C_2)}$: normal infection rates of COVID-19
- $n_{(M)}$: high infection rate
- $n_{(I)}$: very low infection rate

Model Values:

D_1 = 1000; n_d1 = 10; D_2 = 390; n_d2 = 5; C_1 = 150; n_c1 = 7;







C_2 = 200; n_c2 = 6; M = 160; n_m = 6; I = 100; n_i = 1;

Results - Parameter Sweeps





Edge weights vs Total infections (per pod)

Interesting trend: When the cases were increasing and the edge weights are increased, the maximum infections start to decrease after a peak threshold value. For example, in pods D_1, D_2, and I.




Results - Model

Graph of
podded model



- Mean Total Infections
- Mean infected individuals in each pod
- 100 simulations and 100 replications



Interpretation

Results:

- Maximum intensity of peak infections is affected directly by adjusting the infection rate (probabilistic), recovery rate (probabilistic), and edge weights within the pods
- Desired results achieved by using unequal pod sizes & different edge weights

Limitations:

- Interactions with only individuals inside the migrant workers' model
- Not having unequal edge weights for individuals across different pods
- Individuals bound to be in the same pod all throughout the simulation

Next steps:

- Include pods for the general population
- Hybrid model